



Posts and Telecommunication Institute of Technology
Faculty of Information Technology 1

Introduction to Artificial Intelligence

Bayesian network

Dao Thi Thuy Quynh

Contents

- ▶ Definition and construction of Bayesian network
- ▶ Inference in Bayesian network

Problem in representing probability

- ▶ Inference problem:
 - Give evidence E_1, E_2, \dots, E_n
 - Need to define requirement Q by computing $P(Q|E_1, E_2, \dots, E_n)$
- ▶ If there are all simultaneous probabilities
 - Can compute conditional probability above
- ▶ Simultaneous probability table has size that increases exponentially of the number of variables
 - Too big in reality

Need a more realistic represent and inference


Example (1 / 2)

- ▶ Problem: A person comes home from work, need to guess if there is someone in the house?
- ▶ Know that:
 - If family members get away, the yard lights are often (not always) turned on
 - When there is no one at home, a dog is tied outside
 - If being sick, the dog is also tied outside
 - If the dog is outside, family members can hear the barking


Example (2/2)

- ▶ Define the following 5 random variables:
 - *O* : there is no one at home
 - *L* : yard lights is turned on
 - *D* : the dog is tied outside
 - *B* : the dog is sick
 - *H* : can hear the barking

The relations between nodes



Bayesian networks




Definition of Bayesian networks

- ▶ Bayesian network includes 2 parts:
 - The first part is **directed acyclic graph**, in which each node corresponds to a random variable, and each (directed) edge represents the relation between the root node and the targeted node.
 - The second part is **the conditional probability table** including conditional probability of child node when knowing the combinations of values of parent node

Independence probability in Bayes Network


- ▶ Bayes network allows to represent briefly all of simultaneous probabilities
 - The reduction by using the independence feature of probability in network
- ▶ Independence probability
 - Each node V is independent of all nodes that are not descendants of V , if knowing the values of parent nodes of V
 - Example: H is conditional independent with L, O, B if knowing D



Compute simultaneous probability in Bayes Network

► Example:

$$\begin{aligned}
 &P(H, \neg L, D, \neg O, B) \\
 &= P(H \mid \neg L, D, \neg O, B) P(\neg L, D, \neg O, B) \\
 &= P(H \mid D) P(\neg L, D, \neg O, B) \\
 &= P(H \mid D) P(\neg L \mid D, \neg O, B) P(D, \neg O, B) \\
 &= P(H \mid D) P(\neg L \mid \neg O) P(D, \neg O, B) \\
 &= P(H \mid D) P(\neg L \mid \neg O) P(D \mid \neg O, B) P(\neg O, B) \\
 &= P(H \mid D) P(\neg L \mid \neg O) P(D \mid \neg O, B) P(\neg O) P(B) \\
 &= (0.3)(1 - 0.6)(0.1)(1 - 0.6)(0.3)
 \end{aligned}$$



Compute simultaneous probability in Bayes Network (General)

$$P(X_1 = x_1, \dots, X_n = x_n) = \prod_{i=1}^n P(X_i = x_i \mid \text{parents}(X_i))$$

Construct Bayes Network

- ▶ 2 ways to build bayes network:
 - By hand (by human)
 - Base on the knowledge of human about the problem
 - Include 2 steps: define the structure of graph and fill values in conditional probability table
 - Machine learning from data: in case there are data of combinations of variables
 - Distributing probabilities presented by network best fits the frequency of occurrence of values in the data set

Construct Bayes Network (by hand)

1. Define the set related random variables
2. Choose the order for variables

Example: X_1, X_2, \dots, X_n

3. **for $i = 1$ to n do**


- a. Add a node for X_i
- b. Select $parents(X_i)$ is the smallest set of given nodes so that X_i is conditionally independent of all previous nodes if knowing $parents(X_i)$
- c. Add a directed arc from each node $parents(X_i)$ to X_i
- d. Add conditional probability values $P(X_i|parents(X_i))$ or $P(X_i)$ if $parents(X_i) = \emptyset$

Example (1 / 2)

- ▶ A person installed an anti-theft alarm system at home
- ▶ The system will alarm when there is a thief
- ▶ But, the system can alarm (inaccurately) if there is a tremor by an earthquake
- ▶ In case hearing the alarm, 2 neighbors Nam and Việt will call the house owner
- ▶ Due to many different reasons, Nam and Việt can announce inaccurately, for instance, due to noise they can not hear any alarm or vice versa, they mistake another sound for the alarm


Example (2/2)

- ▶ **Step 1:** Select variables: use 5 following variables:
 - T (having thief), \mathcal{D} (earthquake), C (the system alarm), N (Nam calls), V (Việt calls)
- ▶ **Step 2:** Variables are arranged by order: T, \mathcal{D}, C, N, V
- ▶ **Step 3:** Follow the steps in the figure, we can build the network (for simplicity, the figure only shows the structure and does not have the conditional probability table)




The impact of ordering nodes

- ▶ Building Bayes network in reality is not simple
 - Choosing the order of nodes to choose the set of father node having small size is difficult
- ▶ Assume that variables are arranged by different order:
 $N, V, C, T, Đ$



Generally independent feature of probability: Definition of d-seperation (1 / 5)

- ▶ If the value of node C is not known:
 - According to the independent feature of Bayes Network N and V are independent (unconditional)
- ▶ If the value of node C is known
 - Are N and V still independent?



The knowledge learned does not allow to answer this question

Generally independent feature of probability:

Definition of d-seperation (2 / 5)

- ▶ Definition *d-seperation* answer the question about the independence of set of nodes X with set of nodes Y when knowing the set of nodes E on a Bayes network
 - Nodes X and nodes Y are called as being *d-seperated* by nodes E if X and Y are independent when knowing E
 - Nodes X and nodes Y are *d-connection* if they are not *d-seperated*
- ▶ To define *d-seperation* of sets X and Y , we first define *d-seperation* between 2 single nodes x of X and y of Y
 - 2 sets of nodes will be independent if each node in one set is independent of all nodes in the other

Generally independent feature of probability: Definition of d-seperation (3/5)

- ▶ **Principle 1:** Node x and y are *d-connected* if there is an unblocked path *between 2 nodes*. In contrast, if there is no such path, x and y are *d-seperated*

- A path is a sequence of contiguous arcs, regardless of the direction of the arcs
- An unblocked path is a path on which no 2 adjacent arcs are directed at each other
- Nút có hai cung hướng vào như vậy gọi là nút *xung đột*




- The connection and seperation features following **Principle 1** is *unconditional* and so the independence of probability is defined by **Principle 1** is unconditionally independent.

Generally independent feature of probability:


Definition of d-separation (4/5)

- ▶ **Principle 2:** node x and y are *d-conditional connected* when knowing the set of nodes E if there exists unblocked path (not include any conflict node) and does not pass any node of E . In contrast, if there is no such path, we say that x and y are *d-seperated* by E . In other words, every paths between x and y (if any) are blocked by E .
 - When knowing the value of some nodes (set of nodes E), the *independence* or *dependence* between remanining nodes can be changed
 - the *independence* or *dependence* in this case is called as *d - conditional seperation* by the set of variables E



Generally independent feature of probability: Definition of d-seperation (5/5)

- ▶ **Principle 3:** If a **conflict node** is member of set E , or having descendant in set E , so that node does not block paths through it
 - Assume that we know an event is caused by 2 or more causes, if we already know 1 cause is true then the probability of the other causes is reduced, if we know 1 cause is false then the probability of other causes increases



s and y are d-connection, x and u are d-seperation

Contents


- ▶ Definition and construction of Bayesian network
- ▶ Inference in Bayesian network

Lessons learned (recall)

- ▶ How to build a bayes network (by hand)
- ▶ Bayes network allows to reduce the representation
 - No need to save the entire simultaneous probability table
- ▶ Can compute simultaneous probability of any combination of values of variables
- ▶ Therefore, to compute every posterior probability needed for inference


Example of posterior probability

► Compute $P(L | B, \neg H)$



Example of posterior probability

► $P(L|B, \neg H) = \frac{P(L, B, \neg H)}{P(B, \neg H)}$



Step 1: compute $P(L, B, \neg H)$

Step 2: compute $P(\neg L, B, \neg H)$

Step 3: compute

$$\frac{P(L, B, \neg H)}{P(L, B, \neg H) + P(\neg L, B, \neg H)}$$

Simultaneous probability
is computed as the
previous lesson

General case

$$P(E_1|E_2) = \frac{P(E_1 \wedge E_2)}{P(E_2)}$$

$$= \frac{\text{The sum of simultaneous probabilities including } E_1 \text{ and } E_2}{\text{The sum of simultaneous probabilities including } E_2}$$

► Problem



- Require to list simultaneous probabilities having E_1, E_2
- The number of simultaneous probabilities increases exponentially by the number of variables \Rightarrow unrealistic

► General case in bayes network is complete-binary problem ☹ ☹ ☹

Inference in reality

► Inference for a particular case

- When network is in form of single connection (poly tree): there is no more than 1 path between any 2 nodes




- There exists an algorithm with linear complexity for poly tree

► Approximate inference by sampling

Inference for a particular case


► The simplest case:

- When the evidence E and result Q have only one direct connection
- Distinguish 2 cases:
 - **Causal inference** (top to bottom): need to compute $P(Q|E)$ when E is the parent node of Q
 - **Diagnostic** (bottom to top): need to compute $P(Q|E)$ when E is the child node of Q



Causal inference (1 / 3)


► Example: compute $P(D|B)$ $P(O) = 0.6$ $P(B) = 0.3$



Causal inference (2/3)

► Example: compute $P(D|B)$ $P(O) = 0.6$ $P(B) = 0.3$

$$\begin{aligned}
 P(D|B) &= \frac{P(D, B)}{P(B)} \\
 &= \frac{P(D, B, O) + P(D, B, \neg O)}{P(B)} \\
 &= \frac{P(D|B, O)P(B, O) + P(D|B, \neg O)P(B, \neg O)}{P(B)} \\
 &= \frac{P(D|B, O)P(B)P(O) + P(D|B, \neg O)P(B)P(\neg O)}{P(B)} \\
 &= P(D|B, O)P(O) + P(D|B, \neg O)P(\neg O) \\
 &= (0.05)(0.6) + (0.1)(1 - 0.6) \\
 &= 0.07
 \end{aligned}$$



Diagnostic inference (3/3)

► Example: compute $P(D|B)$

$$P(D|B) = \frac{P(D, B)}{P(B)}$$

$$= \frac{P(D, B, O) + P(D, B, \neg O)}{P(B)}$$


$$= \frac{P(D|B, O)P(B, O) + P(D|B, \neg O)P(B, \neg O)}{P(B)}$$

$$= \frac{P(D|B, O)P(B)P(O) + P(D|B, \neg O)P(B)P(\neg O)}{P(B)}$$

$$= P(D|B, O)P(O) + P(D|B, \neg O)P(\neg O)$$

$$= (0.05)(0.6) + (0.1)(1 - 0.6)$$

$$= 0.07$$




Step 1: Convert conditional probability to simultaneous probability

Step 2: Use the independent feature of probability in Bayes network, rewrite simultaneous probability in form conditional probabilities of child node when knowing values of parent nodes

Step 3: Use probability values from conditional probability table to compute

Diagnostic inference (1 / 5)

- Example: compute $P(\neg B | \neg D)$




Diagnostic inference(2/5)

► Follow Bayes

$$P(\neg B|\neg D) = \frac{P(\neg D|\neg B)P(\neg B)}{P(\neg D)}$$

$$P(L|O) = 0.3$$

$$P(L|\neg O) = 0.6$$



compute $P(\neg D|\neg B)$ like the previous part $P(H|\neg D) = 0.8$

Diagnostic inference(3/5)

► Follow Bayes

$$P(\neg B|\neg D) = \frac{P(\neg D|\neg B)P(\neg B)}{P(\neg D)}$$


$P(L|O) = 0.3$
 $P(L|\neg O) = 0.6$

compute $P(\neg D|\neg B)$ as above

$$\begin{aligned}
 P(\neg D|\neg B) &= P(\neg D|O, \neg B)P(O) + P(\neg D|\neg O, \neg B)P(\neg O) \\
 &= (0.9)(0.6) + (0.8)(0.4) \\
 &= 0.86
 \end{aligned}$$

$$P(\neg B|\neg D) = \frac{(0.86)(0.7)}{P(\neg D)} = \frac{0.602}{P(\neg D)}$$

To compute $P(\neg D)$, we will compute $P(B|\neg D)$



Diagnostic inference (4/5)

$$\triangleright P(B|\neg D) = \frac{P(\neg D|B)P(B)}{P(\neg D)} = \frac{(1-0.07)0.3}{P(\neg D)} = \frac{0.279}{P(\neg D)}$$

► Use

$$P(\neg B|\neg D) + P(B|\neg D) = 1$$

$$\frac{0.602}{P(\neg D)} + \frac{0.279}{P(\neg D)} = 1$$

Then $P(\neg D) = 0.881$

Replace:

$$P(\neg B|\neg D) = \frac{0.602}{P(\neg D)} = \frac{0.602}{0.881} = 0.683$$

Diagnostic inference (5/5)

► Follow Bayes

$$P(\neg B|\neg D) = \frac{P(\neg D|\neg B)P(\neg B)}{P(\neg D)}$$

$P(L|O) = 0.3$
 $P(L|\neg O) = 0.6$

$P(O) = 0.6$ **O** $P(B) = 0.3$ **B**

Step 1: convert to causal inference using Bayes principle

$P(D|\neg O, \neg B) = 0.1$
 $P(D|\neg O, B) = 0.2$

$P(H|D) = 0.3$ **H**
 $P(H|\neg D) = 0.8$

compute $P(\neg D|\neg B)$ as above

$$\begin{aligned}
 P(\neg D|\neg B) &= P(\neg D|O, \neg B)P(O) + P(\neg D|\neg O, \neg B)P(\neg O) \\
 &= (0.9)(0.6) + (0.8)(0.4) \\
 &= 0.86
 \end{aligned}$$

Step 2: perform the same as causal inference

$$P(\neg B|\neg D) = \frac{(0.86)(0.7)}{P(\neg D)} = \frac{0.602}{P(\neg D)}$$

To compute $P(\neg D)$, we will compute $P(B|\neg D)$


General method

- ▶ Applies to both causal inference and diagnostic inference
 - **Step 1:** Convert conditional probability to simultaneous probability
 - **Step 2:** Use the independent feature of probability in Bayes network, rewrite simultaneous probability in form conditional probabilities of child node when knowing values of parent nodes
 - **Step 3:** Use probability values from conditional probability table to compute

Inference by sampling

- ▶ In general case: Inference in Bayes network is complete-binary (very complicated)
- ▶ Can infer approximately by sampling
- ▶ Generate sets of variables with the same simultaneous probabilities of network

Sampling(1 / 2)



- ▶ Randomly select S : $S = \text{true}$ with probability 0.3
- ▶ Randomly select M : $M = \text{true}$ with probability 0.6
- ▶ Randomly select L : probability $L = \text{true}$ depends on the values of S, M above
 - Assume that above steps generate $M = \text{true}, S = \text{false}, L = \text{true}$ with probability 0.1
- ▶ Randomly select R with probability depend on value of M
- ▶ Randomly select T with probability depend on value of L

Sampling(2/2)

- ▶ Assume the we need to compute: $P(R = \text{True} \mid T = \text{True}, S = \text{False})$
- ▶ Sampling many times as above, each generated set of values is called a sample
- ▶ Compute the number of occurrence of events:
 - N_c : number of samples having $T = \text{True}$ and $S = \text{False}$
 - N_s : number of samples having $R = \text{True}, T = \text{True}$ and $S = \text{False}$
 - N : Total number of samples
- ▶ If N is large enough:
 - N_c/N : (approximately) probability $P(T = \text{True} \text{ and } S = \text{False})$
 - N_s/N : (approximately) probability $P(R = \text{True}, T = \text{True}, S = \text{False})$
 - $P(R|T, \neg S) = P(R, T, \neg S)/P(T, \neg S) \approx N_s/N_c$

Generally sampling

- ▶ Need to compute $P(E_1|E_2)$
- ▶ Sample large enough quantity
- ▶ Compute quantity:
 - N_c : number of samples having E_2
 - N_s : number of samples having E_1 and E_2
 - N : Total number of samples
- ▶ If N is large enough, we have: $P(E_1|E_2) = \frac{N_s}{N_c}$



Exercise

- ▶ Do some exercises in the textbook