

Danh mục: medication

Type 2 diabetes in adults: management

Type 2 diabetes in adults: management

NICE guideline

Reference number:

NG28

Published:

02 December 2015

Last updated:

18 February 2026

Download guidance (PDF)

Overview

This guideline covers care and management for adults (aged 18 and over) with type 2 diabetes. It focuses on education, dietary advice, managing cardiovascular risk, managing blood glucose levels, and identifying and managing long-term complications.

For information on related topics, see our
cardiometabolic disease prevention and treatment summary page

Last reviewed:

18 February 2026

We reviewed the evidence on medicines for type 2 diabetes, and made and updated recommendations for people with no relevant comorbidities and for people with common comorbidities. We also made changes to recommendations on insulin, in the context of the withdrawal of insulin products and known brand shortages. For more details, see
update information

This guideline updates and replaces:

NICE guidelines CG66 (published May 2008) and CG87 (published May 2009)

NICE technology appraisal guidance 203 and 248

NICE evidence summaries ESNM20, ESNM26 and ESNM59.

For recommendations on hypertension in people with type 2 diabetes, see the
NICE guideline on hypertension in adults

Next review:

This guideline will be reviewed if there is new evidence that is likely to change the recommendations.

How we prioritise updating our guidance

Decisions about updating our guidance are made by NICE's prioritisation board. For more information on the principles and process, see

NICE-wide topic prioritisation: the manual

For information about individual topics, including any decisions affecting this guideline, see the
summary table of prioritisation board decisions

Recommendations

This guideline includes recommendations on:

individualised care

and

Danh mục: medication

person-centred medicine

managing blood glucose levels

, including

HbA1c measurement and targets

initial medicines

reviewing medicines

further medication

insulin-based treatments

complications

See the

2-page visual summary on factors to take into account when choosing, reviewing and changing medicines

Who is it for?

Healthcare professionals

Commissioners and providers

Adults with type 2 diabetes, and their families and carers

Guideline development process

How we develop NICE guidelines

Your responsibility

The recommendations in this guideline represent the view of NICE, arrived at after careful consideration of the evidence available. When exercising their judgement, professionals and practitioners are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or the people using their service. It is not mandatory to apply the recommendations, and the guideline does not override the responsibility to make decisions appropriate to the circumstances of the individual, in consultation with them and their families and carers or guardian.

All problems (adverse events) related to a medicine or medical device used for treatment or in a procedure should be reported to the Medicines and Healthcare products Regulatory Agency using the

Yellow Card Scheme

Local commissioners and providers of healthcare have a responsibility to enable the guideline to be applied when individual professionals and people using services wish to use it. They should do so in the context of local and national priorities for funding and developing services, and in light of their duties to have due regard to the need to eliminate unlawful discrimination, to advance equality of opportunity and to reduce health inequalities. Nothing in this guideline should be interpreted in a way that would be inconsistent with complying with those duties.

Commissioners and providers have a responsibility to promote an environmentally sustainable health and care system and should

assess and reduce the environmental impact of implementing NICE recommendations

wherever possible.

Related quality standards

Type 2 diabetes in adults