

Danh mục: medication

Type 2 diabetes in adults: management

Type 2 diabetes in adults: management

NICE guideline

Reference number:

NG28

Published:

02 December 2015

Last updated:

18 February 2026

Download guidance (PDF)

Dietary advice and interventions

1.3 Dietary advice

1.3.1

Provide individualised and ongoing nutritional advice from a healthcare professional with specific expertise and competencies in nutrition.

[2009]

1.3.2

Provide dietary advice in a form sensitive to the person's needs, culture and beliefs, being sensitive to their willingness to change and the effects on their quality of life.

[2009]

1.3.3

Encourage adults with type 2 diabetes to follow the same healthy eating advice as the general population, which includes:

eating high-fibre, low-glycaemic-index sources of carbohydrate, such as fruit, vegetables, wholegrains and pulses

choosing low-fat dairy products

eating oily fish

controlling their intake of saturated and trans fatty acids.

[2009]

1.3.4

For recommendations on low-energy and very-low-energy diets for the management of type 2 diabetes, follow the:

NHS Type 2 diabetes Path to Remission Programme

NICE guideline on overweight and obesity management

[2026]

1.3.5

Integrate dietary advice with a personalised diabetes management plan, including other aspects of healthy living such as increasing physical activity and losing weight (see the

NHS Better Health website

[2009, amended 2026]

1.3.6

Individualise recommendations for carbohydrate and alcohol intake, and meal patterns. Make reducing the risk of

Danh mục: medication

hypoglycaemia a particular aim for people using insulin or an insulin secretagogue.

[2009]

1.3.7

Advise adults with type 2 diabetes that they can substitute a limited amount of sucrose-containing foods for other carbohydrates in the meal plan but should take care to avoid excess energy intake.

[2009]

1.3.8

Discourage adults with type 2 diabetes from using foods marketed specifically for people with diabetes.

[2009]

1.3.9

When adults with type 2 diabetes are admitted as inpatients to hospital or any other care setting, implement a meal planning system that provides consistency in the carbohydrate content of meals and snacks.

[2009]

1.3.10

For recommendations on wellbeing advice, see

NICE's guidelines on overweight and obesity management

physical activity: brief advice for adults in primary care

, and

tobacco

[2015, amended 2026]

1.4 Bariatric surgery

1.4.1

For recommendations on bariatric surgery for people with recent-onset type 2 diabetes, see the section on surgical interventions in NICE's guideline on overweight and obesity management

[2015]