Exact Sectors Beyond the Pure-Top Conjecture

Nonlinear composition, protected canonical transport and sharp analytic stopping rules

Nonlinear Hamiltonian research programme

22 October 2026 - internally reviewed research draft

Contents

1 Exact sectors beyond the pure-top conjecture **1**

2 Pure total tops with unequal perfect-power transverse tops **4**

3 A complete parabolic quadratic-composition family **7**

4 The exact six-dimensional core of a squared shear coordinate **9**

5 Exact observable closure inside the six-dimensional shear-square core **11**

6 Nonlinear shear compositions: a complete exact-core phase diagram **13**

7 Three equations determine the analytic dynamic core **16**

8 A rational stopping theorem and an algebraic persistent cubic **21**

9 Protected nonlinear canonical transport of exact dynamic sectors **25**

10 Balanced-top audit and a minimal proof repair **28**

11 Independent adversarial assessment **30**

12 Bounded review of Claude's subsequent near-edge theorem **34**

13 Source and priority boundary **36**

1 Exact sectors beyond the pure-top conjecture

Research overlap with `claude/adoring-ritchie-1hq1kw`, inspected at `e91e14d` on 22 October 2026. All results use one canonical pair and the compendium's fixed flat Weyl convention. Written proofs were reviewed independently inside this research session; historical priority and external peer review remain open.

1.1 The consequential finding

The blanket pure-top part of conjecture P003 is false. The simplest active counterexample obtained here has degree four:

$$Q = p + q^2, \qquad H = Q^2, \qquad \mathcal{D}_\infty(H) = K[Q] + \operatorname{span}_K\{q, qQ, q^2\}.$$

The failure is structural. With `Q=p+bq^2+cq+d`, `b!=0`, the complete family `H=P(Q)+epsilon q` has core `K[Q]+Kq` for `deg P>=3`, and the displayed larger core for `deg P=2`. If `epsilon!=0`, the whole tower `K[Q]` is active. Consequently a general edge-polynomial finiteness theorem must exclude these families. A Newton edge alone does not imply a finite core.

A second, different phenomenon has a finite three-dimensional active core:

$$J = qp + q^3, \qquad H = qJ^2, \qquad \{H, J\} = H, \qquad \mathcal{D}_\infty(H) = \operatorname{span}\{1, H, J\}.$$

Its extension is a complete classification of all pure total tops with unequal perfect-power transverse top `alpha p^s(q+gamma)^r, r,s>=2`. For `s>=3` the core is `span{1,H}`. For `s=2`, precisely the explicitly recognized nonlinear shear family has a third generator.

There is also a complete six-dimensional family. If `Q=p+u(q), deg u>=3`, and `H=AQ^2+BQ+C+epsilon q`, `A!=0`, the core is exactly `span{1,q,Q,q^2,qQ,Q^2}`. Third-gate tests at Hamiltonian depths zero, one, and two suffice, and depth two is necessary. Thus even finite pure cores need not have dimension two or three.

This six-dimensional sector also connects directly to the compendium's observer machinery. Its exact derivation matrix has Jordan blocks `3,2,1` when the forcing vanishes and `5,1` otherwise. The minimal derivative closures of the targets `q`, `q^2`, and their joint family change from dimensions `2,3,5` to `3,5,6`. The existing independent observer checker replays this reduction, and the closed evolution is polynomial in time.

Claude's near-top completion in commit `268d553` is incorporated on the review branch. The edge theorem and the new pure exceptions have distinct scopes. Independent review supplied two proof clarifications, without changing the theorem or its determinant certificates.

1.2 Other proved overlaps

For `J=qp+u(q)` with `u'''!=0`, composition has a sharp phase diagram: `H=P(J)` has the strip of transverse degree at most two when `deg P=1`, the stationary span of `1,J,J^2` when `deg P=2`, and `span{1,H}` when `deg P>=3`. The distinction from `Q=p+q^2` is essential.

For analytic coefficients `A=C^omega(I)` and finite degree in `q`, write `D={H,·}` and `C_k=T_k(H,·)`. For every analytic `F,V`,

$$\mathcal{D}_\infty(qF) = \ker C_3 \cap \ker(C_3 D) \cap \ker C_5,$$
$$\mathcal{D}_\infty(qF + V) = \ker C_3 \cap \ker C_5 \cap \ker(C_3 D) \cap \ker(C_3 D^2).$$

The bounds are sharp. For every polygamma `F`, and also `F=log Gamma`, an arbitrary analytic `V` still gives the stronger equality `D_infinity(qF+V)=ker C3 intersect ker(C3D)=A+qA`. A meromorphic pole or the logarithmic Gamma singularity rules out every higher sector at once.

Outside that rigid class, the explicit algebraic profile `F=a p^(-1/4)+b p` has the active cubic `G=q^3p^(-3/4)`, with `D G=-(15b/4)G`. All iterated third-order tests can nevertheless miss a fifth-order obstruction in other fractional-power profiles, so the polynomial branch's third-gate arguments must retain their coefficient hypotheses.

The nonlinear canonical mechanism is proved by an exact shear formula. Quantum and classical shear pullbacks agree on transverse degree at most two. At degree three the correction is `-hbar^2 w''/4`, explaining both the protected finite core and the failure of unrestricted nonlinear covariance.

1.3 Proofs and reproduction

The polynomial proofs treat unequal transverse powers, parabolic compositions, six-dimensional cores, and shear compositions. The analytic stopping rules and fractional family connect to the

polygamma work. The canonical transport proof explains the protected sector. The balanced audit repairs the harmless `s<mu` setup restriction; no upper bound is needed. The independent review records the proof assessments.

Run `python3 -B research/core_overlap/verify_all.py` from the repository root. All numerical receipts are supplementary to the proofs. The declared coefficient-box census covers 3,888 exceptional families and 3,888 broken closure perturbations, with additional exact affine transport checks. The existing source populations and sealed baseline proofs are untouched. The portable bundle contains this research package and the required extension code; replay of the retained compendium archives requires the full repository.

1.4 What remains open

The complete classification of arbitrary pure perturbations is still open. The original P003 is disproved, so unresolved pure cases are reported as unclassified rather than assigned the former conjectural core. Claude's subsequent commit `268d553` completes the near-top question; its edge theorem is integrated with two proof clarifications recorded in the follow-up review. Recognition of the new composition families is sufficient for their routes; it is not claimed to recognize every possible exception.

No Nijenhuis, Haantjes, or Moyal operation is claimed to be newly invented. The useful research claims here are the fixed-H sector classifications, their explicit counterfamilies, and the finite stopping theorems.

2 Pure total tops with unequal perfect-power transverse tops

This note uses the flat Weyl convention of the compendium, `D_H G = H_q G_p - H_p G_q`, and the polynomial dynamic core `D_infinity(H) = intersection ker T_k(H,D_H^j ·)` for odd `k >= 3`, `j >= 0`. The coefficient field has characteristic zero. The statements below concern these fixed canonical coordinates; arbitrary nonlinear canonical substitutions are not claimed to preserve Weyl exactness.

2.1 A counterexample to the remaining pure-top conjecture

Put

$$G = qp + q^3, \qquad H = qG^2 = q^3p^2 + 2q^5p + q^7.$$

The total top of `H` is the pure seventh power `q^7`. Its transverse, or `p`-degree, top is `q^3 p^2`, precisely one of the perfect-power exceptions left open in Theorem T. Nevertheless

$$\{H, G\} = H, \qquad T_k(H, G) = 0 \quad (k \geq 3).$$

For the gate identity, every derivative of `G` of total order at least three is purely in `q`, whereas `H` has `p`-degree two. Thus the sole possibly nonzero summand of `T_k(H,G)` contains `partial_p^k H=0`. Antisymmetry gives `T_k(H,H)=0`, and `{H,H}=0`; consequently `G` survives every repeated bracket. In particular, this is an **active** exact observable, not only a stationary partner. The all-degree dimension argument below proves

$$\boxed{D_\infty(H) = \operatorname{span}_K\{1, H, G\}.}$$

This disproves the unrestricted pure/near two-root conjecture as it was stated in `DYNAMIC_CORE_RESULTS.md` at commit `e91e14d`. It does not contradict its proved Theorems B, Q, Q', or the restricted Theorem T.

2.2 Complete classification of the unequal perfect-power exception

Theorem. Suppose the total top of a polynomial `H(q,p)` is `c q^d`, with `d >= 3`, and the `p`-degree top is

$$F = \alpha\, p^s (q + \gamma)^r, \qquad \alpha \neq 0, \quad r, s \geq 2, \quad r \neq s.$$

Set `t=q+gamma`. Then the following classification holds.

If `s >= 3`, the core is `span{1,H}`. If `s=2` (necessarily `r>=3`), the core is also `span{1,H}` except for the following explicitly recognizable family:

$$J = tp + u(t),\ H \ = t^{r-2}(a_2 J^2 + a_1 J + a_0) + \beta J + \delta, \qquad a_2 \neq 0, \quad u \in K[t], \quad \deg u \geq 3.$$

An additive constant of `J` may be removed and absorbed into the displayed parameters. For this family,

$$D_\infty(H) = \operatorname{span}_K\{1, H, J\}, \qquad \{H, J\} = (r-2)(H - \beta J - \delta).$$

The parameters do not impose an artificial degree bound. This is a family of arbitrarily high total degree and a complete classification under the stated unequal perfect-power transverse hypothesis. Historical priority is not claimed.

2.2.1 Proof

Transverse filtration. Filter by degree in p. The top of D_H has p-degree shift s-1; the top of T_k(H,·) has shift s-k because the two arguments receive exactly k derivatives in p in each summand. Therefore the transverse top of a core element lies in D_infinity(F). This remains valid for the nonnegative weight (0,1); finite-dimensional filtered pieces are not required. All polynomials have finite transverse degree, and the usual leading-component comparison applies separately to each identity.

After translating q, F=alpha t^r p^s. The already proved unequal monomial classification (Theorem M) gives

$$D_\infty(F) = \operatorname{span}_K\{1, tp, F\}.$$

It follows directly, by subtracting a multiple of H and descending in transverse degree, that dim D_infinity(H)<=3. At transverse degree zero the core consists only of constants. If a third element exists, normalize its transverse top and subtract a multiple of H to write it

$$J = tp + u(t), \qquad u \in K[t].$$

The core is invariant under D_H, so

$$\{H, J\} = aH + bJ + c.$$

The coefficient of p^s t^r on the left is (r-s)alpha; hence a=r-s !=0.

The gate removes all s>=3 cases. Direct differentiation gives

$$T_3(H, J) = -u'''(t)\,\partial_p^3 H.$$

If s>=3, this forces deg u<=2. But if deg u<=2, J is quadratic. Since the total top of H is c q^d, comparison of total degree d in {H,J}=aH+bJ+c gives a=d: the top term is q H_q=d c q^d, and all contributions from lower total components of H have smaller degree. On the other hand the pure total top implies d>r+s, because F has a nonzero mixed monomial of degree r+s. This contradicts a=r-s. Thus no third element exists for s>=3. The same comparison proves deg u>=3 whenever a third element exists for s=2.

Integrate the closure relation for s=2. Here a=r-2>=1. On the Laurent polynomial algebra K[t,t^{-1},J], the derivation

$$H \longmapsto \{H, J\} = tH_t - (p + u')H_p$$

is t partial_t at fixed J. Thus

$$H + \frac{b}{a}J + \frac{c}{a} = t^a P(J), \qquad P \in K[J].$$

Since H has p-degree two, P has degree two and nonzero leading coefficient. Set beta=-b/a, delta=-c/a. This is the displayed family. Conversely, its closure identity follows by direct differentiation. Every odd gate kills J since H has p-degree two; every odd gate kills H by antisymmetry. Therefore the span of 1,H,J is an invariant exact sector. The transverse upper bound makes it the whole core. This proves the theorem.

2.3 An exact decision test in the original polynomial coefficients

Write `H=A(t)p^2+B(t)p+C(t)` with `A=alpha t^r, r>=3`, and put `a=r-2`. Choose the harmless normalization `u(0)=0`. The coefficient of `p` in the closure identity is equivalent to

$$2\alpha t^r u' = tB' - (a+1)B - bt.$$

Its coefficient of `t` fixes `b=-a[t]B`. The numerator must be divisible by `t^r`; if so, integration uniquely determines the polynomial `u` with `u(0)=0`. Finally test whether

$$tC' - Bu' - aC - bu$$

is constant. This finite calculation is necessary and sufficient. It neither fits a bounded ansatz nor infers an all-degree claim from a finite matrix. The accompanying checker replays the closure identity and every nonzero odd gate using independent PBW operator arithmetic.

2.4 Why the coordinate mechanism is subtle

The nonlinear triangular substitution `p -> p+u(q)/q` explains the classical eigenvector form when `u(0)=0`. Its Weyl exactness in this sector follows from the separate derivative argument above, rather than from a claim that the Moyal product is covariant under nonlinear canonical maps. For example, the cubic choice `u=q^3` is outside affine covariance and creates a pure total top from an unequal transverse monomial.

3 A complete parabolic quadratic-composition family

Use the flat Weyl convention and polynomial dynamic core of the compendium. Let the coefficient field have characteristic zero, and set

$$Q = p + bq^2 + cq + d, \qquad b \neq 0, \qquad H = P(Q) + \varepsilon q, \qquad P \in K[z].$$

For `deg P>=2` the total top of `H` is a pure power of `q`. These are edge polynomials with constant transverse leading coefficient; no nonlinear covariance of the Moyal product is assumed.

Theorem. The full polynomial core in all degrees is

$$\boxed{D_\infty(H) = \begin{cases} K[q,p], & \deg P \leq 1, \\ K[Q] + \operatorname{span}_K\{q, qQ, q^2\}, & \deg P = 2, \\ K[Q] + Kq, & \deg P \geq 3. \end{cases}}$$

For `epsilon=0`, `K[Q]` is stationary and `q` is active, with `{H,q}=-P'(Q)`. For `epsilon!=0`, even the tower `K[Q]` is active: `{H,f(Q)}=epsilon f'(Q)`. Thus these families disprove the unrestricted pure-top collapse conjecture both through finite active extensions and through an infinite active exact sector. Historical priority is not claimed.

3.1 Proof

The polynomial change of variables `t=q`, `z=Q` is used only to analyze the defining differential operators. Its inverse is `p=z-bt^2-ct-d`, so it is a polynomial algebra isomorphism. Chain rule gives

$$D_H = \varepsilon\partial_z - P'(z)\partial_t,$$

and, by expanding the third transvectant,

$$\boxed{C_3 = -P'''(z)(\partial_t^3 + 6b\partial_t\partial_z) - 6bP''(z)\partial_t\partial_z^2.}$$

The linear summand `epsilon q` contributes no gate. The coefficient `c` and the additive constant `d` cancel in the displayed chain-rule identity.

Exactness of the tower. With the standard Moyal parameter `hbar`,

$$Q \star f(Q) = Qf(Q) - \frac{b\hbar^2}{4} f''(Q) = f(Q) \star Q.$$

Every star power of `Q` is its ordinary power plus lower-degree polynomials in `Q`. Induction therefore identifies `K[hbar][Q]` with the star algebra generated by `Q`. It is commutative. In particular every odd transvectant between `P(Q)` and `f(Q)` vanishes separately, since each is the coefficient of a distinct power of `hbar`. The same holds for the added `epsilon q`, whose gates vanish. This proves exactness of all of `K[Q]` without claiming covariance under a nonlinear canonical transformation.

Degree at least three. Let a core element have positive `t`-degree `N`:

$$G = \sum_{j=0}^{N} t^j f_j(z), \qquad f_N \neq 0.$$

The coefficient of `t^{N-1}` in `C_3G=0` is `-6bN(P'' f_N')'=0`. Since `P''` is nonconstant, a polynomial `f_N` can satisfy this only if `f_N` is constant. If `N>=2`, the next coefficient similarly forces `f_{N-1}` constant:

the partial_t^3 term cannot reach t^{N-2}. In D_HG, the coefficient of t^{N-1} is consequently -N P'f_N; the possible term epsilon f_{N-1}' is zero. The coefficient of t^{N-2} in C_3D_HG is then

$$6bN(N-1)f_N\big((P'')^2\big)',$$

which is nonzero. Hence N<=1; when N=1, the first comparison shows G=f_0(z)+a t. Conversely K[z]+Kt is invariant under the displayed D_H, and its generators pass every gate: the tower was proved exact and q is linear. This proves the degree-at-least-three case using only C_3 and one further bracket.

Degree two. Write P'(z)=kz+l, k!=0. The third gate is now C_3=-6bk partial_t partial_z^2, and no higher odd gate occurs because deg H=4. Its polynomial kernel consists of

$$G = A(z) + zB(t) + C(t).$$

The decisive identities are exactly

$$C_3D_HG = 12bk^2B'', \qquad C_3D_H^2G = -12bk^3C''' \quad \text{once } B'' = 0.$$

They force B''=0 and C'''=0. The forcing term epsilon partial_z contributes no term to either decisive identity. Thus the only extra generators are t,tz,t^2. Their brackets are

$$\begin{aligned} D_Ht &= -P'(z),\\ D_H(tz) &= \varepsilon t - zP'(z),\\ D_H(t^2) &= -2tP'(z). \end{aligned}$$

These lie in the asserted sector, whose third gate is zero. All later brackets are therefore exact. If deg P<=1, H has degree at most two and every gate vanishes, giving the whole polynomial ring. This completes the proof.

3.2 Concrete lowest-degree counterexample

For Q=p+q^2 and H=Q^2=q^4+2q^2p+p^2,

$$D_\infty(H) = K[p+q^2] + \operatorname{span}\{q, q(p+q^2), q^2\}.$$

In particular q evolves exactly and nontrivially. This quartic is outside the previously listed pure-top strip and affine structures: its transverse second derivative is nonzero and its mixed term has degree three.

3.3 Exact recognition

Normalize the coefficient of p in Q to one and its constant term to zero. If mu=deg_p H>=2 and the coefficient of p^mu is the nonzero constant a_mu, the coefficient B(q) of p^{mu-1} must satisfy

$$V(q) = \frac{B(q) - B(0)}{\mu a_\mu} = bq^2 + cq, \qquad b \neq 0.$$

This fixes Q=p+V(q). Substitute p=z-V(q) into H; recognition succeeds exactly when the result is a polynomial in z plus epsilon q. Thus the adjunct decision rule uses exact identities and never a fitted finite-degree ansatz. Its independent checks compare the transformed third gate against PBW arithmetic and directly test the asserted finite-degree dimensions.

4 The exact six-dimensional core of a squared shear coordinate

Let `u in K[q]` have degree at least three, with `K` a field of characteristic zero, and put `Q=p+u(q)`. Use the original flat Moyal product in `(q,p)` and the convention `{H,G}=H_q G_p-H_p G_q`.

Theorem SS. For any quadratic polynomial

$$H = AQ^2 + BQ + C + \varepsilon q, \qquad A \neq 0,$$

the full polynomial dynamic core is exactly

$$\boxed{\mathcal{D}_\infty(H) = \operatorname{span}_K\{1, q, Q, q^2, qQ, Q^2\}.} \tag{SS1}$$

It therefore has dimension six in all degrees. These pure-top Hamiltonians are outside the conjectured two-dimensional collapse. Their positive structure is an elementary polynomial canonical shear, but covariance of the flat Moyal product under arbitrary nonlinear canonical maps is not assumed in the proof.

Proof. Polynomial substitution identifies `K[q,p]` with `K[q,Q]`. The Hamiltonian derivation and the third gate, expressed in this polynomial coordinate ring, are exactly

$$D = -(2AQ + B)\partial_q + \varepsilon\partial_Q, \qquad C_3 = (2AQ + B)u'''(q)\partial_Q^3 - 6Au''(q)\partial_q\partial_Q^2. \tag{SS2}$$

For clarity, the derivatives in (SS2) keep `Q` fixed. To check the gate, use `H_qqq=(2AQ+B)u'''+6Au''`, `H_qqp=2Au''`, and `H_qpp=H_ppp=0`, and then `partial_q(old)=partial_q(new)+u' partial_Q`. The terms involving `u'u''` cancel exactly.

Let `V` be the polynomials of total degree at most two in `(q,Q)`. Equation (SS2) kills `V`, and `D` preserves `V`. Every element of `V` has degree at most two in the original variable `p`, as does `H`, so all odd transvectants of order at least five vanish. Thus `V subset D_infty(H)`.

Suppose a core element `G` lies outside `V`, and let `G_N` be its highest component in total degree, with `N>=3`. Write `m=deg u>=3` and let `u_m` be the leading monomial of `u`. The top parts of the operators for this total-degree filtration are

$$D_0 = -2AQ\partial_q, \qquad C_0 = 2AQu_m'''\partial_Q^3 - 6Au_m''\partial_q\partial_Q^2.$$

The weighted shifts are zero for `D_0` and `m-5` for `C_0`. The other terms have strictly smaller shifts; this includes the `B` term of `C3`. Hence `C3 D^rG=0` implies `C_0 D_0^rG_N=0` for every `r>=0`. Let `a` be the largest `q` exponent occurring in `G_N`. By homogeneity,

$$D_0^a G_N = c_0 Q^N, \qquad c_0 \neq 0.$$

But then

$$C_0 D_0^a G_N = 2Ac_0(N)_3 u_m''' Q^{N-2} \neq 0,$$

because `N>=3`, `A c_0!=0`, and `u_m'''!=0` in characteristic zero. This contradiction proves (SS1), including the constant-force term `epsilon q`. The argument gives a finite failure witness for every proposed polynomial outside the core, not a finite-degree extrapolation.

Sharp uniform stopping rule. In fact,

$$\boxed{\mathcal{D}_\infty(H) = \ker C_3 \cap \ker(C_3 D) \cap \ker(C_3 D^2).} \tag{SS3}$$

To prove the uniform depth bound, write `u_m=alpha q^m`. The top operator `C_0D_0^r` sends `q^aQ^c` to a single monomial, with coefficient

$$2A\alpha m(m-1)(-2A)^r(a)_r(c+r)_2\big[(m-2)(c+r-2)-3(a-r)\big]. \tag{SS4}$$

For fixed `r` the output shift is independent of `(a,c)`, so distinct input monomials cannot cancel. Every monomial with `a+c>=3` fails one of the checks `r=0,1,2`. If `c>=2` and the `r=0` bracket is zero, then `a>=1` and the `r=1` bracket is `m+1`, nonzero. If `c=1`, then `a>=2` and `r=1` gives `-3(a-1)`, nonzero. If `c=0`, then `a>=3` and `r=2` gives `-3(a-2)`, nonzero. All the relevant falling factorials are nonzero. Thus the joint top kernel has total degree at most two. The same total-degree filtration used in the proof lifts this bound to the actual operators, proving (SS3).

Two bracket steps are necessary for this gate: the cubic `q^3` satisfies

$$C_3q^3 = C_3Dq^3 = 0, \qquad C_3D^2q^3 = -288A^3u'' \neq 0. \tag{SS5}$$

So discarding the last condition in (SS3) gives a strictly larger space. The theorem settles both the all-degree core and the exact uniform bracket depth for detecting it using `T3`.

4.1 Recognition in the original coordinates

Suppose a supplied rational Hamiltonian has degree two in `p`, its `p^2` coefficient is a nonzero constant `A`, and its `p` coefficient is `b(q)`. Set

$$B = b(0), \qquad u(q) = \frac{b(q)-B}{2A}, \qquad Q = p + u(q).$$

If `deg u>=3` and `H-AQ^2-BQ` is affine in `q`, Theorem SS applies. This test uses exact coefficient arithmetic. It rejects a nonconstant quadratic coefficient and rejects residual terms of degree at least two scope. The canonical shear may subsequently be composed with affine symplectic substitutions, under which the original flat Moyal structure is covariant.

The qualification `deg u>=3` matters: when `u'''=0`, the last contradiction does not apply. For instance if `deg u=2`, all polynomials in `Q` pass every gate and remain in that space under `D`, giving an infinite core. Such an input must not be assigned a six-dimensional bound by this theorem.

`analytic_shear_square.py` supplies the exact recognizer and a direct transvectant checker. It verifies the transformed gate on monomials, all six basis elements under every nonzero odd correction, invariance under `D`, and finite-box dimensions with full output retained. The proof above establishes the all-degree result.

No literature-priority claim is made. The result is a scoped classification for the fixed original affine Moyal product and an explicit correction to the pure-top collapse conjecture.

5 Exact observable closure inside the six-dimensional shear-square core

Take Theorem SS with `Q=p+u(q)`, `deg(u)>=3`, and

$$H = AQ^2 + BQ + C + \varepsilon q, \qquad A \neq 0.$$

Its exact sector has the ordered basis `(1,q,Q,q^2,qQ,Q^2)`. This supplies a direct bridge to Claude's finite target-observer closure machinery: repeated Hamiltonian derivatives of any target in the sector become an exact finite rational matrix problem. No truncation of the ambient polynomial algebra is required.

With coefficient vectors written as columns, the matrix of `D={H,.}` is

$$M = \begin{pmatrix} 0 & -B & \varepsilon & 0 & 0 & 0 \\ 0 & 0 & 0 & -2B & \varepsilon & 0 \\ 0 & -2A & 0 & 0 & -B & 2\varepsilon \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -4A & 0 & 0 \\ 0 & 0 & 0 & 0 & -2A & 0 \end{pmatrix}.$$

In particular `Dq=-(2AQ+B)` and `DQ=epsilon`. The exact flows in the Hamiltonian-time convention `exp(tD)` are

$$Q(t) = Q + \varepsilon t, \qquad q(t) = q - (2AQ + B)t - A\varepsilon t^2.$$

Every basis observable follows by substituting these two expressions; for example `q^2(t)=q(t)^2`. The original classical momentum coordinate is `p(t)=Q(t)-u(q(t))`; this nonlinear reconstructed observable generally lies outside the exact sector, so it is not asserted to be its flat-Moyal symbol. The time convention here is the compendium's observable derivation convention; changing the sign of that convention reverses time.

5.1 Nilpotency and the role of a constant force

For `epsilon=0`, `D^3=0` on the sector and the ranks of `D,D^2,D^3` are `3,1,0`. Its Jordan block sizes are `3,2,1`. Indeed `D=-(2AQ+B)partial_q` on quadratic polynomials in `(q,Q)`, so its kernel has dimension three and the image of its square has dimension one.

For `epsilon!=0`, the ranks of `D,D^2,D^3,D^4,D^5` are `4,3,2,1,0`; the Jordan block sizes are `5,1`. The displayed flow shows `D^5=0`, and

$$D^4 q^2 = 24A^2\varepsilon^2 \neq 0.$$

Thus there is a size-five block, and the remaining one dimension is a stationary block. A constant force changes the organization of the exact sector without changing its dimension.

5.2 Minimal derivative closures for targets

The span generated by a target and all its repeated derivatives is its Krylov closure. For all parameter choices in the theorem,

Target family	epsilon=0	epsilon nonzero
`q`	2	3
`q^2`	3	5
`q` and `q^2` together	5	6

To see the last row explicitly, let `L=2AQ+B`. For zero force the joint closure is `span{q,L,q^2,qL,L^2}`. For nonzero force the `q^2` chain spans `q^2,qL,L^2-2A epsilon q,L,1`; adjoining the `q` target fills the sixth dimension. These are minimal closure dimensions, not a claim about every possible sensing model or noisy measurement design.

`audit_sector_dynamics.py` checks the matrix, closed forms, ranks and target dimensions exactly. It also feeds the transposed matrix into the existing `extensions/observer/observation_closure_reference.py` compiler and replays its independent `observer-closure-v1` certificate checker. Transposition is essential: that backend acts on target rows and the state consists of basis observable values. This is an application of the existing closure machinery, with no external-priority claim.

6 Nonlinear shear compositions: a complete exact-core phase diagram

Use the fixed flat Weyl convention `D_H={H,.}=H_q partial_p-H_p partial_q`. Work over any field of characteristic zero. Put

$$J = qp + u(q), \qquad u \in K[q], \quad u''' \neq 0, \qquad H = P(J), \quad \deg P = \mu \geq 1.$$

Then the entire polynomial dynamic exact core is

$$\boxed{D_\infty(P(J)) = \begin{cases} K[q] \oplus pK[q] \oplus p^2K[q], & \mu = 1, \\ \operatorname{span}_K\{1, J, J^2\}, & \mu = 2, \\ \operatorname{span}_K\{1, H\}, & \mu \geq 3. \end{cases}}$$

The last two sectors are stationary. The total homogeneous top of `H` is a pure power of `q`, of degree `mu deg(u)`. Its transverse, or `p`-degree, top is a balanced power `(qp)^mu`. Thus these are the balanced **transverse** exceptions in §6.6 of Claude commit `e91e14d`, not counterexamples to its balanced **total-top** theorem B.

For example,

$$J = qp + q^3, \quad H = J^2, \qquad D_\infty(H) = \operatorname{span}\{1, J, H\}.$$

This is a second counterexample to P003. It is stationary, unlike the active unequal-transverse family in `pure_perfect_power.md`. It also shows why nonlinear canonical covariance cannot be assumed: a quadratic Hamiltonian in `J` retains only a quadratic stationary sector, while `T3(J^2,J^3)=72q^3J` is nonzero.

6.1 A closed composition identity

For arbitrary univariate polynomials `F` and `G`, direct differentiation gives

$$\boxed{T_3(F(J), G(J)) = q^3u'''(q)(F'(J)G'''(J) - F'''(J)G'(J)).}$$

Here no nonlinear covariance is being used. To verify the identity, put `A=p+u'`, `B=q`, `C=u''`, `E=u'''`. The four third derivatives of `F(J)` are

$$\begin{aligned} F(J)_{qqq} &= F'''A^3 + 3F''AC + F'E, \\ F(J)_{qqp} &= F'''A^2B + F''(CB + 2A), \\ F(J)_{qpp} &= F'''AB^2 + 2F''B, \\ F(J)_{ppp} &= F'''B^3. \end{aligned}$$

Insert these and their `G` counterparts into the alternating third transvectant. All terms cancel except `B^3E(F'G'''-F'''G')`. In particular,

$$T_3(J^\mu, J^s) = \mu s(s-\mu)(s+\mu-3)q^3u'''J^{\mu+s-4},$$

whenever the exponent on `J` is nonnegative; the remaining low-index cases vanish directly. For nonzero `u'''`, this formula already bounds the static exact powers of `J`. The next argument proves that arbitrary polynomial core elements, including those not assumed to be functions of `J`, obey the same finite bound.

6.2 Proof for mu at least two

Multiplying `H` by a nonzero scalar or adding a constant does not change the core. Write its leading coefficient as one, and set

$$P(z) = z^{\mu} + bz^{\mu-1} + \cdots, \qquad v(q) = u(q) + \frac{b}{\mu}, \qquad Q = qp.$$

The first two pieces in decreasing `p`-degree are

$$H_0 = Q^{\mu}, \qquad H_1 = \mu v(q) Q^{\mu-1}.$$

The transverse top of any core element belongs to the core of `Q^mu`. By the monomial theorem, that core is `K[Q]`. Therefore a nonconstant core element of transverse degree `s` can be normalized to have the first pieces

$$A_0 = Q^{s}, \qquad A_1 = f(q)p^{s-1}, \qquad s \geq 1.$$

At transverse degree zero the core contains only constants.

One bracket determines the first correction. The top term of `{H,A}` vanishes since `{Q^mu,Q^s}=0`. Its coefficient at the next possible `p`-degree, `mu+s-2`, is

$$\mu q^{\mu-1}\big((s-1)f - qf' + sq^{s}v'\big).$$

The bracket is again a core element. Hence this coefficient, if nonzero, must be a multiple of `q^(mu+s-2)`. After dividing by `mu q^(mu-1)`, the right-hand side must be a multiple of `q^(s-1)`. But the Euler operator `f -> (s-1)f-qf'` has zero coefficient at that very monomial, and `q^s v'` also has zero coefficient there. The putative multiple is therefore zero. Solving the resulting polynomial Euler equation gives exactly

$$f = sq^{s-1}v + cq^{s-1}, \qquad c \in K.$$

This argument does not require a degree bound for `f` or `u`.

The first gate bounds the transverse degree. The leading `p`-degree piece of `T3(H,A)` vanishes for two powers of `Q`. At the next possible degree, `mu+s-4`, only the pairs `(H_0,A_1)` and `(H_1,A_0)` contribute. The diagonal summand `cQ^(s-1)` gives zero. Direct falling-factorial expansion of the two remaining terms gives

$$\boxed{[p^{\mu+s-4}]\,T_3(H,A) = \mu s(s-\mu)(s+\mu-3)q^{\mu+s-1}u'''(q).}$$

To check the identity for arbitrary `v`, it suffices by linearity to take `v=q^n`. The coefficient on the left becomes

$$s\,T_3[(\mu,\mu);(s-1+n,s-1)] + \mu\,T_3[(\mu-1+n,\mu-1);(s,s)] = \mu s(s-\mu)(s+\mu-3)n(n-1)(n-2).$$

Here `T3[(a,b);(c,d)]` means the scalar multiplying `q^(a+c-3)p^(b+d-3)` for the two monomials. The symbolic identity is verified independently by `audit_balanced_transverse.py` using a polynomial ring in the three integer parameters. The exceptional index `mu=2,s=1` has formal output degree `-1`; the scalar is identically zero and no nonexistent polynomial output is used in the argument.

Since `u'''` is nonzero, when `mu=2` the only allowed positive values of `s` are `1,2`. The transverse associated graded core has at most one dimension at each allowed degree `0,1,2`; therefore the

entire core has dimension at most three. Both `J` and `J^2=H` survive. Indeed every derivative of `J` of order at least three is in `q` only, whereas `H` has `p`-degree two, so all odd gates kill `J`; antisymmetry kills `H`; their Poisson bracket is zero. This proves the quadratic case.

When `mu>=3`, the factor `s+mu-3` is positive for every integer `s>=1`. The only allowed positive degree is `s=mu`. The same filtration argument bounds the core by two dimensions, and `1,H` attain that bound. This proves the remaining case.

6.3 Linear compositions

For `H=aJ+b`, `a!=0`, the third gate is `T3(H,.)=-a u''' partial_p^3`. Its polynomial kernel consists exactly of the polynomials of `p`-degree at most two. Every higher odd gate kills this space, and

$$D_H = a\big((p+u')\partial_p - q\partial_q\big)$$

preserves it. Thus the static and dynamic exact spaces coincide and equal the stated strip.

If `u'''=0`, this phase diagram does not apply: `J` is quadratic. For `mu>=2`, Theorem B gives the full stationary tower `K[J]`; for `mu=1`, all polynomials are exact. The condition `u'''!=0` therefore identifies a real change in exact observable sectors.

6.4 Scope of the result

The proof is all-degree and applies to every coefficient choice satisfying the hypotheses. Numerical degree cutoffs are not used. The nonlinear substitution `p -> p+u(q)/q`, after removing `u(0)`, motivates the family classically; every assertion about Weyl exactness is proved in the original flat canonical coordinates. No historical-priority claim is made.

7 Three equations determine the analytic dynamic core

This is a finite-determination bridge between the polynomial dynamic-core programme on `claude/adoring-ritchie-1hq1kw` and the analytic polygamma calculations. It concerns the existing flat Weyl–Moyal gates, not a new torsion operation or a literature-priority claim.

Let `I` be a connected open real interval, let `A=C^omega(I,R)`, and suppose `F in A` has `F''` nowhere zero. Work in `A[q]`, with finite degree in `q`. Write

$$H = qF(p), \qquad D = \{H, \cdot\}, \qquad C_j = T_j(H, \cdot),$$

with the compendium's affine canonical coordinates and transvectant convention. The Hamiltonian derivation is `D(q^c g)=q^c(Fg'-cF'g)`.

Theorem FD. The full dynamic core is determined by three equations:

$$\boxed{\mathcal{D}_\infty(qF) = \ker C_3 \ \cap\ \ker(C_3 D) \ \cap\ \ker C_5.} \tag{FD1}$$

Thus no gate above fifth order, and no Hamiltonian iterate above the first, is needed to certify membership. This is an equality in all finite `q` degrees, not a degree cutoff or a numerical stabilization criterion. The three equations are individually indispensable across this class of Hamiltonians.

In particular, with `r=cuberoot(F'')` and

$$K_3 = \frac{FF'''}{3F''} - 3F',$$

the answer is

$$\mathcal{D}_\infty(qF) = \begin{cases} A \oplus qA \oplus \mathbb{R}q^3 r, & K_3 \text{ constant}, \\ A \oplus qA, & K_3 \text{ nonconstant}. \end{cases} \tag{FD2}$$

The constant case permits an active cubic eigenpartner. The theorem does not replace the analytic hypotheses by arbitrary smoothness, nor classify infinite `q` series.

Analytic continuation extension. The three-equation equality (FD1) holds for every real-analytic `F` on a connected interval, including zeros of `F''`. If `F''` vanishes identically, `H` is quadratic and both sides are all of `A[q]`. Otherwise choose any open subinterval where `F''` is nonzero. The proof below excludes every degree other than zero, one and three on that subinterval, hence everywhere by analyticity. A surviving cubic is a local eigenvector of `D`; its analytic identity `DG=lambda*G` extends to the whole interval, and all higher gates kill it by `q` degree. This proves the equality without dividing by `F''` at a zero.

For the explicit formula (FD2) across those zeros, a nonzero cubic coefficient must be a global analytic cube root of a constant multiple of `F''`, and must obey the eigenvector identity. Such a cube root exists exactly when every zero of `F''` has multiplicity divisible by three. Thus the three-equation test is global, while a local cubic formula may fail to extend through a zero.

7.1 Proof by independent degree sectors

For odd `j`, direct expansion gives

$$C_j(q^c g) = (c)_{j-1} q^{c-j+1} \left[jF^{(j-1)} g' - (c-j+1) F^{(j)} g \right]. \tag{FD3}$$

At fixed `j`, different input `q` degrees have different output degrees. The derivation `D` also preserves each degree. Therefore all three equations in (FD1) split by `q` degree.

Degrees zero and one are unrestricted, killed by every gate, and preserved by `D`. For `c>=2`, `C_3(q^c g)=0` is

$$3F''g' - (c-2)F'''g = 0.$$

Put `B=|F''|>0`, `delta=sign(F'')`, and `alpha=(c-2)/3`. A nonzero solution is `g=constant*B^alpha`, and it never vanishes. Then

$$D(q^c g) = q^c g K_c, \qquad K_c = \alpha\frac{FB'}{B} - cF'. \tag{FD4}$$

The equation `C_3D(q^c g)=0` is exactly `K_c'=0`. If all gates kill this generator, a constant `K_c` makes it an eigenvector of `D` and automatically proves persistence under every iterate.

At degree two, `K_2=-2F'` is not constant. At degree three, there are no gates above `C_3`, so precisely the constant-`K_3` possibility survives. It remains to exclude every `c>=4` using `C_5`.

For `c=4`, fifth order gives `B'B''=0`. Analyticity on the connected interval implies `B''=0`, so `F` has degree at most three. For any `c>=4` the latter case is impossible under constant `K_c`: if `F` is quadratic then `K_c=-cF'` has nonzero derivative, and if `F=f_3p^3+...` then its rational expression has leading term `-(8c+2)f_3p^2/3`, which is not constant. An identity on the interval would be a rational identity, so this leading-term argument applies even on bounded intervals away from the zero of `F''`.

For `c>=5`, fifth order gives

$$BB''' = \beta B'B'', \qquad \beta = \frac{5(c-2)}{3(c-4)} > 1.$$

Hence `B''=kappa*B^beta` for a constant `kappa`. If `kappa=0`, the polynomial case above concludes. Suppose `kappa!=0`. A constant `K_c` would give

$$FB' - aF'B = kB, \qquad a = \frac{3c}{c-2} > 0 \tag{FD5}$$

for some real `k`. Apply the following incompatibility lemma, using

$$1 + a(\beta - 2) = \frac{8(c+1)}{(c-4)(c-2)} > 0. \tag{FD6}$$

This excludes every higher degree and proves (FD1) and (FD2).

7.2 Scalar incompatibility lemma

Suppose `B>0`, `F''=delta*B`, `delta in {1,-1}`, `B''=kappa*B^beta`, `kappa!=0`, `beta>1`, `a>0`, and `1+a(beta-2)!=0`. Then `FB'/B-aF'` is not constant.

Write `u=B'`, `v=B''`, `w=F'`. If the expression equalled `k`, then

$$Fu - awB = kB.$$

Differentiate twice, use `v'=beta*vu/B` and `w'=delta*B`, and eliminate `F`. The exact result is

$$Lwv = (3a-1)\delta Bu - (\beta-1)kv, \qquad L = a(\beta-1) + 2 > 0. \tag{FD7}$$

Since `v` is nowhere zero, solve this for `w` and differentiate. Using `w'=delta*B` gives

$$[a(\beta-4)+3]Bv=(3a-1)(1-\beta)u^2. \tag{FD8}$$

There is an open subinterval with `u!=0`, since `v` never vanishes. Differentiate (FD8) there, cancel the nonzero `uv`, and obtain

$$(\beta+1)[a(\beta-4)+3]=2(3a-1)(1-\beta).$$

Their difference factors as

$$(\beta+5)[1+a(\beta-2)]=0,$$

contrary to the hypotheses. The argument makes no division by `u` at a zero, by `F`, by a first-integral constant, or by either factor in (FD8).

7.3 Sharpness and the distinction from static gate determination

The examples below are on `p>0`, and every power is its positive real analytic branch.

The first-step condition is necessary. For `F=p^2`, `G=q^2` passes every static gate, while

$$C_3DG=-48.$$

The fifth-order condition is necessary even after arbitrarily many third-order tests. Take `F=p^{-2/5}` and `G=H^4=q^4p^{-8/5}`. Then `DG=0` and `C_3G=0`, but

$$C_5G=-\frac{548352}{625}p^{-7}\neq 0.$$

Thus testing `C_3D^jG=0` for every `j>=0` still does not determine the true dynamic core.

The initial third-order condition is necessary. For `F=p^2` and `G=H^2=q^2p^4`, the equation `DG=0` makes `C_3DG=0`, and `C_5G=0`, whereas

$$C_3G=48p^3\neq 0.$$

Finally, static exactness has a different universal bound: the full static partner space is `ker C_3 intersect ker C_5 intersect ker C_7`, by the complete analytic partner classification. Seventh order really is required for that static statement: with `F=p^{8/7}` and `G=q^6p^{-8/7}`,

$$C_3G=C_5G=0,\qquad C_7G=-\frac{1940889600}{117649}p^{-7}\neq 0.$$

Dynamic persistence lowers the necessary maximum gate from seven to five. This explains a concrete way in which the analytic coefficient space and the polynomial programme can share an exact finite stopping rule even though their static partner spaces differ.

7.4 A sharper two-equation certificate for gamma and polygamma

Corollary FP. For `F=log Gamma` or `F=psi^(m)`, `m>=0`, on `p>0`,

$$\boxed{\mathcal{D}_\infty(qF)=\ker C_3\cap\ker(C_3D)=A\oplus qA.} \tag{FP1}$$

This eliminates the fifth-order equation as well. It proves the dynamic result directly, without first classifying the static partner space. It still uses all finite `q` degrees with analytic coefficients.

More generally, suppose `F` analytically continues to a genuine meromorphic pole of integer order `t>=1`,

$$F(p) = C_0 p^{-t} + \text{a lower-order Laurent part}, \qquad C_0 \neq 0,$$

after shifting the pole to zero. A hypothetical sector with `c>=2` that passes both equations would make `K_c` constant. But

$$K_c(p) \sim \frac{2C_0}{3}[(c+1)t - c + 2]p^{-t-1}. \tag{FP2}$$

Its coefficient does not vanish because

$$(c+1)t - c + 2 = (c+1)(t-1) + 3 > 0.$$

The meromorphic identity theorem transports a supposed constant identity from the original interval to a punctured neighborhood of the pole. Equation (FP2) contradicts that identity. No continuation of the partner coefficient is required. Every polygamma has such a pole, with `t=m+1` and `C_0=(-1)^(m+1)m!`.

For the logarithmic template `F=-a log p+b(p), a!=0`, with `b` analytic at zero,

$$K_c(p) = \frac{2a(c-2)}{3}\frac{\log p}{p} + \left[ca - \frac{2(c-2)b(0)}{3}\right]\frac{1}{p} + O(|\log p| + 1). \tag{FP3}$$

At `c=2`, the coefficient `2a/p` is nonzero. At every `c>=3`, the logarithmic coefficient is nonzero. Thus `K_c` is never constant, which covers `log Gamma` using its `-log p` singularity. As above, the potential must have the stated continuation; this is not an assertion about a real-analytic function with no access to that singularity.

The two-equation conclusion for these pole and logarithmic classes also holds for `H=qF(p)+V(p)` with **arbitrary analytic `V`**. The highest `q`-degree of a hypothetical partner still satisfies exactly the two equations above: the `V` terms lower `q` degree in both `C_3` and `D`. Thus its degree cannot exceed one. Conversely, `A+qA` is invariant under `D` and killed by every gate for this affine-in-`q` Hamiltonian. This extends the polygamma result to an arbitrary analytic additive potential without a new asymptotic assumption on that potential.

The real integer pole restriction has content: the fractional power `F=p^{-2/5}` in the sharpness example admits a stationary fourth-degree sector through both third-order tests and fails the fifth gate. Thus the two-equation corollary cannot be extended to all algebraic singularities.

7.5 Arbitrary additive potentials: a sharp depth-two theorem

Theorem FA. Let `F,V` be arbitrary real-analytic functions on a connected open interval, and let `H=qF(p)+V(p)`. With the same finite `q`-degree coefficient algebra,

$$\boxed{\mathcal{D}_\infty(H) = \ker C_3 \cap \ker C_5 \cap \ker(C_3 D) \cap \ker(C_3 D^2).} \tag{FA1}$$

Thus an arbitrary analytic additive potential requires at most two Hamiltonian steps and gates no higher than fifth order. Two steps really are necessary in this larger class.

Proof. First suppose `F''` is not identically zero. Choose an open subinterval `J` where it is nowhere zero. Let `S_1` be the common kernel of the first three operators in (FA1). The highest `q`-degree sector of an element of `S_1`, restricted to `J`, satisfies exactly the three equations of Theorem FD for `qF`: every `V` contribution has strictly smaller `q` degree. The proof of FD therefore forces its

degree to be at most three. Analyticity extends the vanishing of all higher coefficients from `J` to the whole interval.

The strip `W=A+qA` is invariant under `D` and killed by every gate. Moreover,

$$\dim_{\mathbb{R}}(S_1/W) \le 1. \tag{FA2}$$

Indeed, the leading coefficient of a cubic is a scalar multiple of `cuberoot(F'')` on `J`, hence has at most one free real scalar globally. The difference of two candidates with equal cubic coefficient has degree at most two. Its quadratic coefficient is constant by `C_3G=0`, and `C_3DG=-12(F'')^2` on the generator `q^2` excludes that coefficient. Thus the difference belongs to `W`.

Now let `G` pass all four equations in (FA1). Then `DG` belongs to `S_1`: the two third-order equations follow from the last two conditions on `G`, and `C_5DG=0` automatically because `DG` has `q` degree at most three. If `G` is not in `W`, (FA2) gives

$$DG = \lambda G + w, \qquad \lambda \in \mathbb{R}, \quad w \in W.$$

As `W` is invariant, every iterate stays in `S_1`, where the third gate vanishes; every higher odd gate vanishes by `q` degree. If `G` is in `W`, persistence already holds. This proves the sufficient direction. Necessity follows directly from the definition of the core.

If `F''` is identically zero and `V'''` is not, the operator is `C_3=-V'''*partial_q^3`. Hence its analytic kernel has `q` degree at most two, is preserved by `D`, and is killed by every higher gate. If both derivatives vanish identically, `H` is quadratic and every gate vanishes on all of `A[q]`. These cases finish the proof.

Exact sharpness witness. On `p>0`, take

$$\begin{aligned} H &= qp^{-1/4} + p^{7/4}, \\ G &= q^3 p^{-3/4} - \frac{21}{25} q^2 p^{5/4}. \end{aligned}$$

Direct rational-power differentiation gives

$$\begin{aligned} C_3 G = C_5 G = C_3 DG = 0, \\ DG = -\frac{168}{25} q^2 + \frac{147}{50} q p^2, \\ C_3 D^2 G = \frac{63}{8} p^{-9/2} \neq 0. \end{aligned} \tag{FA3}$$

This partner passes every static gate and remains static after one Hamiltonian bracket, but escapes at the second. Thus the one-step bound for `qF` cannot be carried unchanged to `qF+V`. The corollary for gamma and polygamma is stronger because its singularity excludes the cubic sector before this obstruction can arise.

7.6 Checkers and status

`depth_verify.py` independently expands the differential operators on sparse fractional-power monomials using rational arithmetic. It checks the sector formula, all four sharpness examples, the parameter identity (FD6), a grid of the scalar factors, and the exact pole coefficient in (FP2), as well as the sharp depth-two additive-potential example. The separate adversarial audit checks the formal differential elimination in the lemma.

8 A rational stopping theorem and an algebraic persistent cubic

This note connects the polynomial affine strips in `programmes/DYNAMIC_CORE_RESULTS.md` with the analytic `qF(p)` results of the polygamma programme. All statements use the original affine coordinates `(q,p)`, the constant-Poisson Moyal product, and finite degree in `q`. They do not assert covariance of this same product under a nonlinear canonical substitution. Historical priority is unassessed.

Let `I` be a connected open real interval, `A=C^omega(I,R)`, `H=qF(p)+B(p)`, and `D={H,·}` with `{H,G}=H_q G_p-H_p G_q`. Let `C3=T3(H,·)`, using the transvectant convention of the compendium. Formal Moyal exactness requires every odd transvectant of order at least three to vanish. Put

$$\mathcal{D}_\infty(H) = \{G \in A[q] : T_k(H, D^r G) = 0 \quad (k \geq 3 \text{ odd},\ r \geq 0)\}.$$

8.1 Rational coefficients: one gate and one bracket suffice

Theorem A. Suppose `F` is rational with real coefficients, has no pole on `I`, and `F''` is nowhere zero on `I`. Let `B` be arbitrary in `A`. Then, in all finite `q` degrees,

$$\boxed{\ker C_3 \cap \ker(C_3 D) = \mathcal{D}_\infty(H) = A \oplus qA.} \tag{A1}$$

Thus the full infinite family of odd gates and repeated brackets has a uniform stopping rule in this coefficient class: `T3(H,G)=0` and `T3(H,{H,G})=0` already characterize the core. No degree cutoff on the observable is present. The polynomial specialization is the affine strip already classified by Claude's theorem; the uniform two-check certificate and arbitrary analytic potential `B(p)` are the refinement.

Proof. The strip `A+qA` is invariant under `D` and all its elements pass every odd gate, because the `q` degrees of both `H` and a strip element are at most one. It remains to exclude `G` of `q` degree `c>=2` from the left side of (A1).

Write its highest coefficient as `q^c g(p)`, with `g` not identically zero. The highest `q` coefficient of `T3(H,G)` is

$$c(c-1)q^{c-2}\,[3F''g' - (c-2)F'''g]. \tag{A2}$$

The term `B` contributes only to lower `q` degrees. Let `r=cuberoot(F'')`, the nonzero real-analytic real branch. Equation (A2) forces `g=C r^(c-2)` with a nonzero real constant `C`; in particular `g` is nowhere zero. The highest `q` coefficient of `DG` is

$$Fg' - cF'g = K_c g, \qquad K_c = \frac{c-2}{3}\frac{FF'''}{F''} - cF'. \tag{A3}$$

Applying (A2) to `DG` gives `3c(c-1)F'' g K_c'=0`, whether or not `DG` still has degree `c`. Consequently `K_c` would have to be constant. We prove this impossible for rational `F` and every integer `c>=2`.

If `F` has a finite complex pole of order `m>=1`, write `F=a t^(-m)+O(t^(-m+1))`, where `t=p-p0` and `a!=0`. Then

$$K_c = \frac{2}{3}\big[(c+1)m - c + 2\big]a\,t^{-m-1} + O(t^{-m}). \tag{A4}$$

The bracket is at least three. Hence `K_c` has a pole and is not constant. This includes nonreal poles: a rational function constant on the real interval would be the same constant rational identity over

the complex numbers. If `F` has no finite complex poles, it is a polynomial. Its degree `n` is at least two since `F''!=0`. If its leading coefficient is `a`, then

$$K_c = -\frac{2}{3}\big[(c+1)n + c - 2\big]a\,p^{n-1} + O(p^{n-2}), \tag{A5}$$

again nonconstant. This contradiction proves (A1). Every division was by a function assumed or proved nonzero on the selected interval.

The numerator of the obstruction `K_c'` is particularly simple:

$$3(F'')^2K_c' = (c-2)\big[F'F'''F'' + FF''''F'' - F(F''')^2\big] - 3c(F'')^3. \tag{A6}$$

This is affine in the integer `c`; equations (A4)–(A5) prove its nonvanishing uniformly rather than searching over sectors.

Corollary A.1 (a meromorphic pole suffices). The same conclusion (A1) holds if, instead of rationality, `F` has a meromorphic extension to a connected complex domain containing `I` and at least one finite pole. The analytic potential `B` still needs to be defined only on `I`. Indeed, `K_c` is meromorphic on that domain. If it were constant on `I`, the identity theorem would make it constant meromorphically, in contradiction to the nonzero pole coefficient (A4). The pole may be nonreal or outside the interval. No asymptotic estimate at infinity is required.

Every polygamma `F=psi^(m)`, including digamma at `m=0`, satisfies this corollary on `p>0`: it is meromorphic with poles of integer order `m+1` at the nonpositive integers, and `F''=psi^(m+2)` has constant nonzero sign on the positive axis. Thus

$$\ker T_3(q\psi^{(m)} + B, \cdot) \cap \ker[T_3(q\psi^{(m)} + B, D\cdot)] = A \oplus qA \tag{A6a}$$

for arbitrary analytic `B(p)` and every `m>=0`. The standard polygamma definitions, recurrence and reflection identities are documented in NIST DLMF 5.15. Log-Gamma is not meromorphic; its separate asymptotic proof of the same conclusion is recorded in `depth_finite_determination.md`.

8.2 An explicit algebraic family across the boundary

Theorem B. On `p>0`, let

$$F(p) = ap^{-1/4} + bp, \quad a \neq 0, \qquad H = qF(p), \qquad X = qp^{-1/4}, \qquad G = X^3.$$

Then

$$\boxed{\frac{[H,G]_\star}{i\hbar} = \{H,G\} = -\frac{15b}{4}G, \qquad D^jG = \left(-\frac{15b}{4}\right)^j G.} \tag{A7}$$

In particular `b!=0` gives a completely explicit algebraic active persistent cubic outside the rational coefficient class of Theorem A. It replaces the earlier implicit autonomous-ODE examples with elementary fractional powers. Its full finite-`q`-degree core is

$$\boxed{\mathcal{D}_\infty(H) = A \oplus qA \oplus \mathbb{R}X^3.} \tag{A8}$$

Proof of persistence. Write `H=aX+bJ` with `J=qp`. The direct transvectant identity gives `T3(X,X^3)=0`; all higher odd gates vanish because their combined `q` degree is four. Also `J` is quadratic, so it contributes no gate, and

$$\{X, X^3\} = 0, \qquad \{J, X^3\} = -\frac{15}{4}X^3.$$

This proves (A7) independently of the complete analytic classification. Equivalently, `F''=(5a/16)p^(-9/4)` and `K3=FF'''/(3F'')-3F'=-15b/4`.

Proof of completeness. The complete analytic `qF` boundary theorem is not needed here. Since `B=0`, distinct `q` degrees cannot cancel in either `C3` or `C3D`. Equations (A2)–(A3) apply to each degree `c>=2` separately, and direct simplification gives

$$K_c = \frac{a(3-c)}{2}p^{-5/4} - \frac{b(7c-6)}{4}. \tag{A8a}$$

Because `a!=0`, this is constant exactly when `c=3`. Thus the common kernel of `C3` and `C3D` is exactly the strip plus `R X^3`. Persistence already proved in (A7) identifies it with the full dynamic core. Hence the stronger identity

$$\ker C_3 \cap \ker(C_3 D) = \mathcal{D}_\infty(H) = A \oplus qA \oplus \mathbb{R}X^3 \tag{A8b}$$

is self-contained and holds in every finite `q` degree. The exceptional cubic is selected directly by the factor `3-c`; no imported analytic classification or numerical truncation is necessary.

Why the cubic is a boundary, not a template for arbitrary degrees. For any integer `c>=4`, set

$$\rho = \frac{2-c}{c+1}, \qquad H = qp^\rho + bqp, \qquad G = (qp^\rho)^c.$$

Then `T3(H,G)=0` and `{H,G}=bc(rho-1)G`, so `T3(H,D^jG)=0` for every `j>=0`. Nevertheless, direct fifth-order arithmetic gives

$$T_5(H, G) = -8(c)_4(\rho)_4 q^{c-4}p^{-c-3} \neq 0. \tag{A8c}$$

Thus arbitrarily deep third-order checks cannot replace higher gates for general analytic coefficients. At `c=3`, `rho=-1/4` and the falling factorial `(c)_4` vanishes; this is the genuine cubic family above. The distinction matters when exporting a successful `T3` argument from the polynomial branch to a wider coefficient ring.

8.3 A sharp nonlinear canonical-coordinate obstruction

Set

$$Y = \frac{4}{5}p^{5/4}.$$

Then `\{X,Y\}=1` and

$$H = aX + \frac{5b}{4}XY. \tag{A9}$$

So this nonlinear Hamiltonian becomes quadratic in the algebraic canonical coordinates `(X,Y)`. Rebuilding the standard flat Moyal product in those new coordinates would make every observable exact. The original flat Moyal product in `(q,p)` has a different answer: although `X` and `X^3` belong to its dynamic core, `X^2` does not. Indeed,

$$
\begin{aligned}
T_3(H, X^2) &= -\frac{15a}{16} p^{-15/4}, \\
\frac{[H, X^2]_\star}{i\hbar} &= -\frac{5b}{2} X^2 + \frac{5a\hbar^2}{128} p^{-15/4}, \\
X \star X &= X^2 + \frac{\hbar^2}{64} p^{-5/2}.
\end{aligned}
\tag{A10}
$$

The star square is a quantum eigenobservable: `[H,X star X]_star/(i hbar)=-(5b/2)(X star X)`, by the derivation law. This does not put its ordinary leading term `X^2` in the classical dynamic core; its explicit correction cancels the anomaly in (A10). The core is a vector space, not an algebra under ordinary multiplication. The formulas exhibit the coordinate dependence directly, without mistaking a nonlinear change of coordinates for covariance of the same flat Moyal product.

The classical evolution is elementary as well. Put `Z=p^(5/4)`. Then `DZ=(5/4)(a+bZ)` and `DX=-(5b/4)X`. For `b!=0`, `Z(t)=exp(5bt/4)(Z(0)+a/b)-a/b`, as long as the trajectory stays in `p>0`, and `G(t)=exp(-15bt/4)G(0)` is exact under both evolutions. These are local-domain evolution statements, not assertions of a globally complete flow across the singular boundary `p=0`.

8.4 Relation to the branch and verification scope

Theorem A recovers Claude's polynomial affine strips and adds an observable-degree-independent gate-depth bound. Theorem B shows exactly why extending a polynomial-core collapse statement to arbitrary analytic coefficients would fail: a simple algebraic coefficient already adds an active cubic. Log-Gamma and all polygammas lie on the collapsing side by their separately proved analytic asymptotic/pole arguments.

`analytic_bridge_check.py` uses exact rational arithmetic and a sparse fractional-power derivative engine. It checks the gate identities, persistence, canonical relation, anomaly and star-square correction; it also tests Theorem A on finite polynomial/Laurent boxes by a direct matrix kernel calculation with no output projection. Those boxes cross-check conventions; the proof above, not their size, supplies the all-degree claims.

The distinction between affine covariance and nonlinear canonical changes belongs to established Weyl quantization theory. Primary background includes Mark Gotay, On the Groenewold–Van Hove problem for R^{2n}, arXiv:math-ph/9809015. No claim is made that a new geometric torsion operation or a new general quantization obstruction has been invented. The specific stopping theorem and explicit family here still require a focused priority check.

9 Protected nonlinear canonical transport of exact dynamic sectors

This note explains why a nonlinear canonical change can preserve a particular flat Weyl–Moyal dynamic sector even though it does not preserve flat Moyal exactness on every observable. It supplies a structural explanation for the pure-top counterfamilies in this programme. No historical priority claim is made for canonical transformations, Weyl ordering, or their quantum correction.

We use

$$\{A,B\} = A_q B_p - A_p B_q, \qquad \frac{[A,B]_\star}{i\hbar} = \sum_{r\ge 0} \frac{(-1)^r \hbar^{2r}}{2^{2r}(2r+1)!} T_{2r+1}(A,B).$$

All statements below are algebraic in characteristic zero. For polynomial inputs the relevant series terminate. Formal powers of `hbar` are retained; they are never fitted from numerical samples.

9.1 The protected shear lemma

Let `w(q)` be a polynomial, choose an antiderivative `U'(q)=w(q)`, and write `sigma_w A(q,p)=A(q,p+w(q))`. The quantum canonical automorphism

$$\mathcal{S}_w = \exp\left(\frac{\mathrm{ad}_\star U}{i\hbar}\right)$$

has the exact factorization

$$\boxed{\mathcal{S}_w = \sigma_w \exp\left(-\frac{\hbar^2}{24} w'' \partial_p^3 + \frac{\hbar^4}{1920} w^{(4)} \partial_p^5 - \cdots\right).} \tag{C1}$$

Indeed, `T_k(U,A)=U^(k) partial_p^k A`, and every coefficient in these operators depends on `q` alone. Consequently all the displayed differential operators commute. The exponential is finite on each polynomial because each nonzero application lowers its `p` degree. Since its generator is an inner star derivation, `S_w` preserves the star product.

In particular

$$\mathcal{S}_w A = \sigma_w A \qquad (\deg_p A \le 2). \tag{C2}$$

This degree threshold is sharp when `w''` is nonzero:

$$\mathcal{S}_w(p^3) = (p+w)^3 - \frac{\hbar^2}{4} w''. \tag{C3}$$

Here is the resulting transport principle. Suppose `H_0` and a linear space `V` consist of polynomials of `p` degree at most two, every odd gate `T_k(H_0,v)` with `k>=3` vanishes on `V`, and `{H_0,V} subset V`. Then

$$\sigma_w V \subseteq \mathcal{D}_\infty(\sigma_w H_0). \tag{C4}$$

To prove it, apply the star automorphism `S_w` to `[H_0,v]_star/(i hbar)={H_0,v}`. Every member of this identity is in the protected subspace, so (C2) changes the quantum transport into the ordinary canonical pullback. Hence `[sigma_wH_0,sigma_wv]_star/(i hbar)={sigma_wH_0,sigma_wv}`. The same argument applies to every Hamiltonian iterate, because these remain in `V`. This proves (C4). It does not assert transport of the entire core under an arbitrary nonlinear change of coordinates.

9.2 A complete pure-top family

Take `a>=1`, `A!=0`, `w` of degree `t>=2`, and arbitrary constants `B,C,beta,delta`. Set

$$z = qp, \quad H_0 = q^a(Az^2 + Bz + C) + \beta z + \delta,$$

$$G = qp + qw(q), \qquad H = q^a(AG^2 + BG + C) + \beta G + \delta. \tag{C5}$$

Then

$$\boxed{\mathcal{D}_\infty(H) = \mathrm{span}\{1, H, G\}, \qquad \{H, G\} = a(H - \beta G - \delta).} \tag{C6}$$

The lower bound follows from the protected shear lemma. It can also be checked directly: `H` has `p` degree two and `G=qp+u(q)`. In every term of an odd gate of order at least three, either three `p` derivatives hit `H`, two hit `G`, or at least two `q` derivatives and one `p` derivative hit `G`. Each possibility vanishes. The bracket in (C6) follows by transporting the identity for `H_0` under `sigma_w`.

For completeness, filter by `p` degree. The transverse top of `H` is `Aq^(a+2)p^2`. Its all-degree core is `span{1,qp,Aq^(a+2)p^2}` by the unequal-monomial theorem, because `a+2>2`. The associated graded of the core of `H` embeds in this three-dimensional core. Therefore the core of `H` has dimension at most three; the three independent elements in (C6) give equality. The top total-degree term of `H` is a nonzero multiple of `q^(a+2t+2)`, so these are pure tops. The generator `G` has degree `t+1>=3`: its presence is not explained by a quadratic core element. The `p^2` coefficient `Aq^(a+2)` is nonconstant, so the family is outside the previously proved affine and constant-quadratic strip structures.

The smallest example of this form is

$$H = q^7 + 2q^5p + q^3p^2 = q(qp + q^3)^2, \quad G = qp + q^3, \quad \{H, G\} = H. \tag{C7}$$

Thus a nonlinear shear of a three-dimensional exact sector is a mechanism for retaining active polynomial exact dynamics beneath a pure top. It also explains why a conjecture based only on affine normal forms can miss this family.

9.3 The complementary `qF(p)` chart

Let `F` be real analytic and nonzero on an interval `I`, and put

$$Q = qF(p), \qquad P = S(p), \qquad S' = 1/F.$$

Then `{Q,P}=1`, and also `[Q,P]_star=i hbar` exactly. The latter holds because `Q` is affine in `q` and `P` depends only on `p`. Nevertheless ordinary pullback fails already at the square:

$$T_3(Q, Q^2) = 12FF'F'', \quad \frac{[Q, Q^2]_\star}{i\hbar} = -\frac{\hbar^2}{2}FF'F'', \quad \{Q, Q^2\} = 0, \tag{C8}$$

$$Q \star Q = Q^2 + \frac{\hbar^2}{4}(F')^2. \tag{C9}$$

For a polygamma `F=psi^(m)`, the obstruction in (C8) is `12 psi^(m) psi^(m+1) psi^(m+2)`; it is nowhere zero for `m>=1` on the positive axis. For digamma one restricts to an interval avoiding its zero. For `F=log Gamma`, the same expression is `12 log Gamma psi psi^(1)` and is not identically zero on any open interval avoiding the zeros of `F`. These are ordinary consequences of the derivative identities; no numerical special-function evaluation is used in (C8).

The exact quantum pullback of a finite polynomial in `Q` with analytic `P` coefficients is

$$\mathcal{W}_F(Q^n a(P)) = J_Q^n(a(S(p))), \qquad J_Q(A) = \tfrac{1}{2}(Q \star A + A \star Q). \tag{C10}$$

Here `Q` inside the star product denotes `qF(p)`. The commutation relation `[Q,a(S)]_star/(i hbar)=a'(S)` verifies the usual Weyl relations. Formula (C10) is therefore a star-algebra homomorphism from the flat `(Q,P)` algebra to the original one. It is invertible as a formal triangular map: its leading term is ordinary canonical pullback, and its corrections lower `q` degree by positive even integers. Each finite `Q` degree has only finitely many corrections.

Writing `a` now for `a(S(p))`, its first two nontrivial cases are

$$\mathcal{W}_F(Q^2 a) = q^2 F^2 a + \frac{\hbar^2}{4} F'(F'a + Fa'), \tag{C11}$$

$$\mathcal{W}_F(Q^3 a) = q^3 F^3 a + \frac{\hbar^2 q}{4}\big[(5F(F')^2 - F^2F'')a + 3F^2F'a'\big]. \tag{C12}$$

For every integer `n>=2`, the coefficient of `hbar^2` is

$$\frac{n(n-1)}{24} q^{n-2} F^{n-2}\big[(2n-1)(F')^2 a + 3FF'a' - (n-2)FF''a\big]. \tag{C13}$$

For an all-`n` verification, write the `hbar^2` term as `q^(n-2)b_n(p)`. The defining Jordan product gives `b_1=0` and

$$b_{n+1} = Fb_n + \frac{n}{8}\left[2nF^{n-1}(F')^2 a + 2F^n F'a' - (n-1)F^n F''a\right].$$

Summing `sum j^2`, `sum j`, and `sum j(j-1)` gives (C13). Thus its arbitrary-degree statement is an induction identity, rather than an extrapolation of the companion checker's finite fixtures. No coordinate covariance of the ordinary flat Moyal product is assumed.

Finally suppose `F''` is nowhere zero and

$$K_3 = \frac{FF'''}{3F''} - 3F' = \lambda.$$

Then `r=cuberoot(F'')` satisfies `r/F^3=c_0 exp(lambda S)` for a nonzero constant `c_0`. The classical persistent cubic `q^3r` is the ordinary pullback of `c_0 Q^3 exp(lambda P)`. Its first integral is

$$FF'' - 5(F')^2 - 3\lambda F' = C.$$

Substitution into (C12) gives the exact correction

$$\mathcal{W}_F(Q^3 e^{\lambda P}) = q^3 F^3 e^{\lambda S} - \frac{C\hbar^2}{4} q F e^{\lambda S}. \tag{C14}$$

Both terms are classical Hamiltonian eigenvectors of eigenvalue `lambda`. In the explicitly constructed active family `F'=aF^5+b`, `a!=0`, one has `lambda=-5b/3` and `C=0`. Thus its cubic eigenpartner is a second protected nonlinear canonical pullback: the classical and quantum Weyl pullbacks agree on this particular cubic, although the square obstruction (C8) is generally nonzero. The polynomial shear protection (C2) and the analytic cubic protection (C14) supply complementary, precisely scoped links between the two research programmes.

10 Balanced-top audit and a minimal proof repair

The balanced theorem at Claude commit `e91e14d` survives this audit. One step in its published argument needs a correction. Subtracting multiples of `H` does **not** justify requiring an extra element's top exponent to satisfy `s < mu`. For example, a hypothetical extra element can initially have top `(qp)^(mu+1)`. The dynamic core is a vector space, not an algebra; powers of `H` cannot be subtracted without first proving they belong to it.

No additional theorem is needed to repair the argument. Choose any element `G` outside `span{1,H}`. Its top is `c(qp)^s` by the top-weight lemma. If `s=mu`, subtract `cH`; if this leaves a nonconstant polynomial, its degree is smaller than `2mu` and its new exponent differs from `mu`. If it leaves only a constant, the original `G` was in the excluded span. Thus in every case, after rescaling, there is an extra `G` with `s >= 1` and `s != mu`. This is the only condition the rest of the proof uses.

The factor `s(s-mu)` is nonzero for every such exponent, including `s>mu`. The positivity estimate is symmetric in `s` and `mu`: a matched pair of monomials has indices `(x,y)` and `(x-mu+s,y-mu+s)`, all nonnegative. Consequently both `2mu >= k+|u|` and `2s >= k+|u|`, where `k=2mu-x-y` and `u=x-y`. All stated positivity conclusions remain valid unchanged.

There are two boundary cases worth stating explicitly. When the proposed partner monomial has a negative exponent, the bracket relation immediately forces the original off-diagonal coefficient to vanish. When `k` exceeds the degree of one of the two polynomials, this is the same argument with the missing homogeneous component set to zero. Thus induction continues through all components, without a restriction on the relative degrees of `H` and `G`.

The bracket's component at the first unremoved deficit has an off-diagonal support. It must also be diagonal because `{H,G}` lies in the dynamic core, whose top is in `K[qp]`. This justifies setting that component to zero; no assumption that the whole bracket vanishes is needed. Lower components of both polynomials are diagonal by induction and hence Poisson commute, so they do not introduce additional terms at that deficit.

The all-odd-order vanishing for two polynomials in `qp` follows from exchanging the two coordinates in the transvectant, or directly pairing its summands. The finite checks at orders 3, 5, and 7 in `check_balanced.py` are useful calibrations, while this antisymmetry argument supplies the statement for every odd order.

`audit_balanced.py` independently checks the scalar identity directly from falling-factorial transvectants, both relative exponent ranges, support boundaries, and every possible zero in the finite declared grid. It does not replace the symbolic identity or the positivity proof. Its purpose is to exercise the cases omitted by the unnecessary `s<mu` assertion.

Theorem T and Lemma F in §6.6 have no additional blocker found in this audit. Their transverse filtration may assign weight zero to `ell`: the associated graded embedding still implies finiteness when the entire top core is finite. For the pure part of T, the exclusion is specifically a **power of a linear form**, `phi=c(ell+gamma)^e`; the phrase "perfect power" should not be read as excluding, for example, a square of a nonlinear polynomial.

10.1 Independent review of the new unequal transverse classification

The theorem and proof in `pure_perfect_power.md` have been read independently. The transverse associated graded argument bounds the entire core by three dimensions. Its extra element must have the exact shape `J=(q+gamma)p+u(q+gamma)`, so the proof uses no bounded-degree guess for `u`. The identity `T3(H,J)=-H_ppp u'''` eliminates every nonlinear `u` when the transverse degree is at least three. A quadratic `J` is impossible under a pure total top: the top of `{H,J}` then has coefficient `d` against `H`, whereas its transverse top forces coefficient `r-s`, with `d>r+s`.

For transverse degree two, the Laurent-ring integration of the closure equation is valid because `J` is algebraically independent of `t=q+gamma` and the derivation is `t*d/dt` at fixed `J`. The eigenvalue `r-2` is a positive integer. This forces `H-beta J-delta=t^(r-2)P(J)` with `P` quadratic. The coefficient recognition equations have the correct signs in the stated Poisson convention. Their constant test

is necessary and sufficient, rather than only a necessary test obtained by projecting the image.

The family is polynomial in the original variables and its nonquadratic `J` passes every odd gate by the separate derivative argument. Hence it is a valid counterexample to P003, not a nonlinear covariance assertion.

There is a further simple stationary example outside the unequal scope of that theorem: `J=qp+q^3`, `H=J^2`. Here the total top is pure `q^6` and the transverse top is balanced `q^2p^2`. The span of `1,J,H` survives, while `T3(H,J^3)=72q^3J` is nonzero. This does not affect the balanced **total-top** theorem, whose leading homogeneous part is a nondegenerate quadratic power.

11 Independent adversarial assessment

This assessment concerns the overlap work based on Claude commit `e91e14d`. It is an internal mathematical review, not external peer review, formal proof verification, or a priority claim.

11.1 Pure-top counterexample and replacement stratum

The counterexample `J=qp+q^3`, `H=qJ^2` is valid. Direct differentiation gives `{H,J}=H`. Every derivative of `J` of order at least three involves only `q`, while `H` has degree two in `p`, so every odd gate vanishes. The span of `1,H,J` is therefore invariant and exact. Its dimension is three, because the three elements have different transverse degrees.

The all-degree upper bound does not follow from a finite search. It follows from the transverse filtration and the already proved monomial theorem: the transverse top `q^3p^2` has core `span{1,qp,q^3p^2}`. An additional core element of transverse degree zero must be constant; after subtracting a multiple of `H`, any other element has the uniquely normalized form `qp+u(q)`. Hence the full core is exactly three-dimensional. This is a counterexample to the unrestricted conjecture P003 at the inspected commit, not to any of the restricted proved statements B, Q, Q′, or T.

The proof of the complete unequal perfect-power classification in `pure_perfect_power.md` is sound. The transverse filtration is valid with weight `(0,1)` even though the filtered pieces are not finite-dimensional: each polynomial has finite transverse degree and the leading comparison is applied separately to each operator identity. Finite-dimensionality of the resulting associated graded core, rather than of the ambient filtered pieces, supplies the dimension bound.

If a third core element exists, it must be `J=tp+u(t)` and satisfy `{H,J}=(r-s)H+bJ+c`. The identity `T3(H,J)=-u'''H_ppp` forces `u` to be quadratic when `s>=3`. A quadratic `J` would force the top eigenvalue to be the total degree `d`, whereas the transverse comparison forces it to be `r-s`; pure total top implies `d>r+s`, a contradiction. For `s=2`, Euler integration in the Laurent coordinates `(t,J)` gives precisely `H=t^(r-2)P2(J)+beta J+delta`. The coefficient-based decision test has the correct signs, and its divisibility conditions account for the resonant coefficient of degree `r-1` in the linear-in-`p` part.

The independent checker `review_checks.py` uses its own direct polynomial differentiation. It verifies 120 parameterized family members, 2,760 odd gate identities, and brackets through depth five. For the basic degree-seven example, independent modular matrices have nullity three through total degree nine. Those matrices corroborate the example; the filtration proof is what establishes the all-degree statement.

11.2 Analytic finite determination

The theorem in `depth_finite_determination.md` is correct under its stated real-analytic, connected-interval, finite-`q`-degree hypotheses and the condition that `F''` is nowhere zero. The three tests `C3 G=0`, `C3 D G=0`, and `C5 G=0` determine the full core. Sector splitting is exact because `D` preserves `q` degree and each gate shifts every input sector by the same amount.

For degrees at least five the shorter incompatibility argument is an independent derivation of the obstruction. In the notation of the theorem, let `E=Fu-awB-kB`, with `F'=w`, `w'=delta B`, `B'=u`, `u'=v`, and `v'=beta vu/B`. Define `L=a(beta-1)+2` and

```
J = Lwv + (beta-1)kv + (1-3a)delta Bu,
R = [a(beta-4)+3]Bv - (3a-1)(1-beta)u^2.
```

The following three identities hold formally in a Laurent-polynomial ring:

```
E'' = J + beta v E/B,
v (J/v)' = delta R,
R' = uv (beta+5)[1+a(beta-2)].
```

The independent checker verifies these by exact symbolic arithmetic with its own Laurent-

polynomial derivation. There is no numerical differential approximation and no hidden division by a first-integral constant. Since `v` is nowhere zero and `u` is nonzero on some open subinterval, `E=0` would force the nonzero scalar `(beta+5)[1+a(beta-2)]` to vanish. The parameter identity shows the second factor is strictly positive for every integer sector degree at least five. The separate degree-two, degree-four, and `B''=0` cases are correctly treated in the theorem note.

The direct fractional-power examples in that note establish necessity of each of the three tests. In particular, testing third-order corrections at every bracket depth is still insufficient universally: the stationary fourth-power example passes all of those tests and fails the fifth-order gate.

11.3 Rational coefficients with arbitrary analytic potential

The highest-`q`-degree proof of the rational-coefficient extension is also valid: in `H=qF(p)+B(p)`, the additional potential contributes one lower `q` degree to both the third gate and the derivation. It cannot cancel the highest-sector obstruction. At a pole of integer order `m>=1`, the candidate sector obstruction has nonzero leading coefficient proportional to `(c+1)m-c+2`; if the rational function has no finite poles, its polynomial leading coefficient is proportional to `(c+1)n+c-2`, also nonzero. Thus no sector of degree at least two persists. This is an exact algebraic argument, not a finite-degree extrapolation. A meromorphic function with a finite pole gives the same contradiction when continuation to that pole is part of the explicitly stated domain assumptions.

11.4 Scope retained

No claim here transports the flat Moyal product through an arbitrary nonlinear canonical map. The special shear family's exactness is proved directly by derivatives. No claim identifies an analytic-function kernel with a rational polynomial Weyl-ideal receipt. Literature priority remains unresolved, and the remaining transverse balanced exception is a different stratum from the unequal exception classified here.

11.5 Nonlinear shear compositions with balanced transverse top

The theorem in `audit_balanced_transverse.md` is approved. Its key transverse Euler resonance is exact: for `H=P(J)`, `J=qp+u(q)`, and a hypothetical core element with first transverse pieces `Q^s+f(q)p^(s-1)`, one bracket forces `f=sq^(s-1)v+cq^(s-1)`. The coefficient at the resonant Euler monomial vanishes on the left, so a nonzero diagonal bracket top is impossible. No assumption that powers of a core element remain in the core is made in this reasoning.

The first nonzero third-gate coefficient is `mu s(s-mu)(s+mu-3)q^(mu+s-1)u'''`. The formal negative output degree in the exceptional pair `mu=2,s=1` has zero coefficient and is correctly excluded from the polynomial comparison. Consequently the three regimes are the width-two strip for linear compositions, the stationary space `span{1,J,J^2}` for quadratic compositions, and `span{1,H}` for degree at least three. This concerns a balanced transverse top and does not conflict with the already proved balanced total-top theorem.

Independent direct differentiation now checks this first-gate factor in 90 complete polynomial composition pairs, rather than only in formal monomial coefficients.

11.6 Parabolic quadratic compositions, including linear forcing

The theorem in `pure_quadratic_composition.md` is approved in its full stated scope `Q=p+bq^2+cq+d`, `b!=0`, `H=P(Q)+epsilon q`. The change of variables is a polynomial algebra isomorphism used to transform the operators; it is not an appeal to nonlinear covariance. Direct chain-rule expansion gives the stated third gate and `D=epsilon partial_z-P'(z)partial_t`.

For `deg P>=3`, the leading and second-leading `t` coefficients of a candidate core element are constants. Applying the third gate after one bracket then forces `t` degree at most one; the linear forcing contributes no surviving term to that decisive coefficient. For `deg P=2`, write `P'=kz+l` and

a static candidate as A(z)+zB(t)+C(t). The exact formulas are C3DG=12bk^2 B'' and, after B''=0, C3D^2G=-12bk^3 C'''. Hence the additional generators are exactly t,tz,t^2.

The infinite tower is proved without circularity. Since Q star f(Q)=Qf(Q)-b hbar^2 f''(Q)/4, star powers of Q form a triangular basis of its ordinary polynomial algebra. This algebra is commutative for the star product. Each odd transvectant therefore vanishes individually. The resulting sectors are invariant under the forced derivation as claimed. In particular the forcing makes the tower active, which is a substantial counterexample to unrestricted pure-top collapse.

The independent checker adds 48 direct comparisons of the transformed derivation and third gate with differentiation in the original coordinates, plus 262 odd-gate checks on tower and extension generators through two further brackets.

11.7 Arbitrary analytic additive potential: the second depth is sharp

The additive-potential finite-determination theorem in depth_finite_determination.md is also approved. For H=qF(p)+V(p), the first three tests C3G=0, C5G=0, and C3DG=0 force finite q degree at most three. Their solution space modulo the invariant affine strip has dimension at most one: cancellation of two cubic leading coefficients would leave a quadratic candidate, which the same third-gate and first-bracket tests exclude.

The additional equation C3D^2G=0 places DG in that same first-three-test space, since its q degree is at most three and fifth gates are then automatic. Thus DG=lambda G+strip, and all later iterates pass every gate. This proves the four-test stopping rule. The extension across isolated zeros of F'' follows by restriction to a subinterval where it does not vanish and then by analyticity. The identically affine F case is separately elementary, as stated in the note.

Independent sparse fractional-power differentiation verifies the sharp example H=q p^(-1/4)+p^(7/4) and G=q^3p^(-3/4)-(21/25)q^2p^(5/4): its first three tests vanish, while C3D^2G=(63/8)p^(-9/2) is nonzero. This is a genuine failure of a first-bracket stopping rule when the potential is arbitrary.

11.8 Explicit algebraic active cubic

The strengthened analytic_bridge.md is approved, including the family F=a p^(-1/4)+bp, a!=0, on the positive axis. Independent simplification gives the sector obstruction Kc=a(3-c)p^(-5/4)/2-b(7c-6)/4; precisely the cubic degree survives. The claimed eigenvalue, the nonlinear canonical quadratic representation, and the second-power Moyal anomaly all have the stated constants and signs. The general higher-degree fractional-power example has fifth-gate coefficient -8(c)_4(rho)_4, with rho=(2-c)/(c+1), so all-depth third-order exactness does not imply full Moyal exactness.

The gamma/polygamma specialization FP is approved as well. The integer-pole coefficient is nonzero in every integer degree at least two. For the logarithmic template F=-a log p+b(p), direct expansion of F'''/F'' gives the stated logarithmic-over-p leading term; at degree two the remaining 2a/p term is nonzero. The continuation requirement belongs to the coefficient function F, while the additive potential remains arbitrary analytic on the working interval.

11.9 Arbitrary polynomial shear and the exact six-dimensional square sector

The further shear-square theorem is approved. Let Q=p+u(q), deg u=n>=3, and H=AQ^2+BQ+C+epsilon q, A!=0. In coordinates (q,Q) the exact operators are

```
D  = epsilon partial_Q-(2AQ+B)partial_q,
C3 = (2AQ+B)u''' partial_Q^3-6A u'' partial_q partial_Q^2.
```

The proposed sector is every polynomial of total degree at most two in (q,Q), namely span{1,q,Q,q^2,qQ,Q^2}. The transformed derivation preserves it, and the third gate kills it.

Every higher odd gate kills it in the original coordinates as well: both arguments have degree at most two in p, so an odd gate of order at least five requires more than two p derivatives on one argument.

Completeness follows from the stated total-degree filtration in (q,Q). The derivation has top part D0=-2AQ partial_q, of degree shift zero. If a core element had homogeneous top G_N, N>=3, choose its highest q exponent a. Then D0^a G_N is a nonzero multiple of Q^N. The top third gate, of degree shift n-5, sends that monomial to a nonzero multiple of q^(n-3)Q^(N-2). Lower filtration pieces cannot cancel this component. This remains true at n=3, when the gate shift is negative: every lower-degree piece still has strictly lower output degree. Thus a core element cannot have transformed degree above two.

This proof uses repeated third gates only for the upper bound and direct derivative exactness for the six-dimensional lower bound. It never assumes nonlinear covariance or closure under ordinary multiplication. The linear forcing is fully included; it changes only the lower filtration part of the derivation. Direct independent checks in review_checks.py compare both transformed operators with original-coordinate differentiation and check the whole protected sector under repeated brackets.

The stronger uniform depth-two corollary is approved. On a leading monomial q^a Q^c, the top operator C0 D0^r has coefficient proportional to

(a)_r (c+r)_2 [(n-2)(c+r-2)-3(a-r)].

Its output shift is independent of (a,c) at fixed r, so cancellation between different input monomials is impossible. For total degree at least three, these coefficients cannot vanish simultaneously at r=0,1,2. For c>=3, a zero at r=0 forces a>=1, and the bracket at r=1 is n+1. For c=2, positive a already fails at r=0; for c=1, the required a>=2 fails at r=1; and for c=0, a>=3 fails at r=2. The full three-test kernel is therefore exactly the six-dimensional sector.

The depth is sharp for every allowed shear and every linear forcing: q^3 passes the first two tests, while D^2q^3=6q(2AQ+B)^2-6A epsilon q^2 and hence C3D^2q^3=-288A^3u'', which is nonzero. Independent original-coordinate differentiation checks this identity in twenty parameterized fixtures, alongside an exact grid of the monomial factors. This grid supplements the case argument above rather than replacing its all-degree proof.

12 Bounded review of Claude's subsequent near-edge theorem

This read-only follow-up inspected commit `268d553eb6a4434afeaf5fcb4689a6294e5ab688`, specifically Theorem NE in `programmes/DYNAMIC_CORE_RESULTS.md` section 6.7 and the new checker `extensions/hamiltonian/check_near_edges.py`. It supplements the overlap review; it is not a claim that the remaining pure-top conjecture is true.

Assessment: the near-top theorem is supported by the determinant identities and the filtration argument, subject to the two explicit proof clarifications below. Neither clarification supplies a counterexample to Theorem NE. The unrestricted pure-top conjecture still has the exact counterexamples recorded in the overlap package.

The checker was extracted, unchanged, with its PBW arithmetic dependency to a temporary directory. Its full replay passed: `E1`, `E2`, `E2_at1`, `E3`, and `E4`, including 150 direct operator grid checks. The run took approximately 0.44 seconds. No current worktree source file was overwritten during this review.

12.1 First-edge and lattice issues

The first edge has a strictly positive weight. A lower monomial of transverse exponent at most one cannot tie `m ell^(d-1)` as the ratio of the positive weights is decreased from one. A non-affine monomial has exponent at least two and ties at a positive ratio below one. Thus the primitive edge step `(a,-b)` satisfies `b>a>=1`.

A weight-homogeneous polynomial is supported on a single primitive lattice line. Its total-degree top lies in the near monomial core `K[ell]+mK[ell]`, so it has one of the two stated types. Different transverse congruence classes, including the two possibilities when the primitive step has transverse size greater than one, are precisely the separate alpha and beta cases. No half-integer step or unaccounted coset is needed.

The determinant at fixed edge parameters is a nonzero polynomial in the top ordinary degree `N`. Its displayed leading coefficient is nonzero for all admissible integer parameters. It therefore has only finitely many exceptional degrees. An explicit uniform bound in all edge parameters is not needed for the finiteness assertion. Since the core of the edge Hamiltonian is graded by a positive edge weight, a common bound on the ordinary degrees of its homogeneous elements bounds the entire core and makes it finite-dimensional.

The first nonzero edge coefficient may occur after a gap. This does not introduce additional terms into the first-order determinant: at step `t`, every term of `C3(E,G)` or `C3(E,D_E G)` uses only `G_0` and `G_t`, because all intermediate edge coefficients of `E` vanish. In the second-order exceptional case `a=t=1`, the four declared unknowns `y2,e1*y1,e2*g,e1^2*g` cover every contribution. If the formal second edge monomial does not exist, its coefficient `e2` is simply zero; an invertible four-column system still forces `e1^2*g=0`.

12.2 Clarification 1: intermediate coefficients need not all clear

The sentence asserting that the third-gate recurrence clears every coefficient between steps zero and `t` is stronger than necessary and is not always true. For alpha type with `a=1` and `t>=2`, the intermediate monomial `m ell^(N-b)` is in the static near strip and passes every gate of the unperturbed near monomial. It is not forced to vanish by that recurrence.

The first-order determinant remains valid because this intermediate coefficient cannot occur in either step-`t` equation. The sentence should be replaced by the decoupling argument in the preceding paragraph. The exceptional second-order calculation already includes its first-step coefficient explicitly.

12.3 Clarification 2: separate the cubic case in Lemma F

The proof of Lemma F, used after edge finiteness, should separate degree three. For the near cubic top `m ell^2`, the top derivation has the finite chain

```
m -> -2m ell -> 2m ell^2 -> 0.
```

Thus the lemma's stated unbounded-chain argument for every degree `N` outside `{2,d}` is not valid at `d=3,N=1`. For `d>=4`, the argument works: the only positive-degree chain starting points that can encounter the stationary top degree `d` are `N=2` and `N=d`.

Theorem NE at degree three is already Theorem Q. Every non-affine perturbation of a near cubic has a nonzero `m^2` term and otherwise terms affine in `m`; that is exactly the degree-three case covered by the retained binomial certificate. Stating "degree three follows from Theorem Q; apply Lemma F's chain argument for degree at least four" repairs the dependency without new computation or assumptions.

With these clarifications, no conceptual blocker was found in the near-top completion. This bounded review checked the proof mechanism and replayed its symbolic identities; it does not establish historical priority or replace external mathematical review.

13 Source and priority boundary

The source branch reviewed here is `claude/adoring-ritchie-1hq1kw`, pinned at `e91e14d`. The subsequent near-top completion at `268d553` is integrated and separately reviewed. The retained monomial theorem, affine covariance and associated-graded reduction are dependencies of the polynomial classifications. The new analytic arguments are supplied explicitly and do not substitute analytic coefficients into polynomial Weyl-ideal receipts.

The literature checks below establish background overlap, not priority for the fixed-H classifications in this package. No literature-priority claim has been made. These primary sources were inspected on 22 October 2026.

Mark J. Gotay, On the Groenewold-Van Hove problem for R^{2n}, J. Math. Phys. 40 (1999), 2107-2116, arXiv:math-ph/9809015. The quantization obstruction and quantizable polynomial Lie subalgebras are mature background. In particular the coordinate-affine algebra underlying the strip is not a new construction.

T. Dereli, T. Hakioglu and A. Tegmen, Quantum canonical transformations in Weyl-Wigner-Groenewold-Moyal formalism, arXiv:0904.3416, especially sections 2-4. Gauge and point transformations, star-exponential representations and nonlinear function transforms are existing theory. The protected-sector shear explanation in this package is an explicit application of that theory; it is not a claim to have invented quantum canonical transformations.

T. Hakioglu, Extended covariance under nonlinear canonical transformations in Weyl quantization, arXiv:quant-ph/0011076, version 3 (2001), section VI(b) and equations 62 and 84. The gauge-transform operator and its explicit higher-derivative quantum correction already appear in this source. The protected shear mechanism is therefore established background, not a novelty candidate. T. Hakioglu and A. Dragt, The Moyal-Lie Theory of Phase Space Quantum Mechanics, arXiv:quant-ph/0108081, provides a broader Lie-algebraic treatment of these quantum transformations.

Didier Robert, When Poisson and Moyal Brackets are equal?, arXiv:2207.08798, version 2 (2023), Theorem 1.1. Equality against every Schwartz observable under the paper's symbol hypotheses forces a quadratic Hamiltonian. Our fixed nonlinear H and restricted observable sectors have different quantifiers and do not contradict that theorem. The local analytic formal-gate setting here also differs from its global symbol assumptions.

The present claims concern algebraic/formal gate identities for every power of hbar separately, and repeated Hamiltonian brackets. Global self-adjoint operator domains, convergence of a star series for arbitrary smooth functions, spectral predictions, and experimental realizability are outside the claims. Finite-q-degree analytic observables have finitely many gates against H=qF+V, so no infinite gate summation is used there.