

Flavor mechanisms, vacuum alignment and predictive limits

Singlet dynamics, exact finite symmetries, canonical quark matching
and the input audit

PerfectPower research monograph
6 October 2026 (UTC)

The selected model admits controlled vacua near ± 66 degrees. Its initialized equations receive perturbative supersymmetric protection; its energy selection, physical coefficient readout and quark couplings remain separately specified interactions.

Exact rational results and numerical scenario results are distinguished throughout. The worked construction is conditional and does not claim an independently derived theory of flavor.


What this push accomplishes

The subsequent shared-mediator investigation is documented in B5_FLAVOR_MEDIATOR.md and included as the second part of the reader monograph. It derives exact canonical tree matching, establishes a scoped low-degree alignment obstruction, and tests an adjoint alternative. It transmits an initialized golden frame but does not derive that frame from symmetry.

The candidate now has an explicit renormalizable supersymmetric singlet sector, a holomorphic golden-coefficient extension, an actual auxiliary-field relaxation calculation, and a one-loop calculation including every scalar and fermion in that sector. The 34-field extension has 68 real scalar modes and 34 Weyl fermions. In the declared numerical examples, a small scalar energy selector prefers the branches near plus and minus 66 degrees, and the one-loop corrections leave that preference intact among all sixteen continued branches. The construction produces a quantitatively controlled realization of the nominated vacuum. It does not derive the initial superpotential coefficients or the energy-selector pattern from an independently specified symmetry, and it does not derive the nominated exact CKM relation from quark interactions.

The strongest negative result is exact: the scalar-only version of the potential is not closed under its own one-loop divergent counterterm. Polynomial division yields a nonzero torque remainder, and a Bezout identity shows that the remainder vanishes at none of the primitive real order-60 roots. Thus there is a specific radiative protection problem, even before quarks or gauge fields are introduced. The strongest constructive result is an initialized holomorphic constraint circuit whose supersymmetric zeros are perturbatively protected. Soft selection then produces small calculable departures instead of an exact unchanged angle.

The quark calculation exposes another independent issue. The declared flavor charges allow the required hierarchy of operators, but their coefficients are independent. Exact diagonalization of a selected triangular texture, even with its 1-3 coefficient initialized to the golden value, gives a different observable mixing strength. This is a constructive counterexample to assuming that the right spurion powers automatically imply the exact CKM formula.



The original exact carrier and energy winner

$$V'_{\text{lock}}(x) = P(x) [2\lambda P'(x) + \eta Q(x)]$$

Define $x = 2 \cos(\theta)$, $P(x) = x^8 - 7x^6 + 14x^4 - 8x^2 + 1$ and $Q(x) = -1 + 3x - x^3$. The previously constructed potential obeys $V'(x) = P(x)Q(x)$. Its six nonconstant angular terms are $\cos(2\theta) + (2/3)\cos(3\theta) + (2/5)\cos(5\theta) + (1/4)\cos(8\theta) - (2/9)\cos(9\theta) - (1/6)\cos(12\theta)$. V differs from that angular expression by the constant $11/12$. Exact rational Sturm isolation and energy enclosures selected $x = 2 \cos(66^\circ)$ on $[-2, 2]$, with more than 0.060

separation from the next competing stationary energy in the polynomial normalization.

The new family $V_{\text{lock}}(x) = \lambda P(x)^2 + \eta V(x)$, $\lambda \geq 0$ and $\eta > 0$, obeys $V_{\text{lock}}'(x) = P(x)[2\lambda P'(x) + \eta Q(x)]$. Its global winner is unchanged: the added square is nonnegative, vanishes at the old winner, and the original potential already has a unique global winner in x . This statement covers the full declared parameter orthant without discretizing λ or η . The two full-circle CP partners remain degenerate. This is a mathematical family, not an assertion that the interaction generating λ or η has been identified.

The rational stationary-family constraint is stronger than stationarity under arbitrary real couplings at just one chosen angle. Rational coefficients force Galois-conjugate roots to share the polynomial factor. An isolated real linear torque condition at a prescribed angle is not equivalent to the eight rational relations found for the twelve-harmonic carrier. The field model must explain which of those structures it actually enforces.

A precise scalar-loop obstruction

$$h(x) = (4 - x^2)V''(x) - xV'(x), \quad \gcd(P, (h^2)') = 1$$

Give the angle the kinetic term $f^2 (\partial \theta)^2/2$ and the physical potential $\Lambda^4 V(2 \cos(\theta))$. With $D(x) = 4 - x^2$, its dimensionless angular curvature is $h(x) = D(x)V''(x) - xV'(x)$. The physical field-dependent squared mass is $\Lambda^4 h(x)/f^2$. The one-loop scalar divergence contains a coefficient proportional to this mass to the fourth power. Consequently the question of preserving the stationary family reduces to whether P divides the x derivative of h^2 .

For the original potential, the exact remainder is

$R(x) = 5040 - 6880x - 31360x^2 + 43600x^3 + 24160x^4 - 30080x^5 - 4400x^6 + 4800x^7$. Its gcd with P is one. The stored rational Bezout multipliers satisfy $a(x)P(x) + b(x)R(x) = 1$. Since $\sin(\theta)$ is nonzero at every primitive order-60 angle, a nonzero x torque also means a nonzero angular torque. This establishes that the original family is not closed under this particular divergent scalar counterterm. It does not establish that every model containing additional fields must fail.

The coefficient field can also be included. Write $c(x) = 1 - S_{12}(x)$, where $S_n(2 \cos(\theta)) = 2 \cos(n\theta)$, and take $F(\theta, C) = V_{\text{lock}}(2 \cos(\theta)) + \kappa[C - c(x)]^2$. On its tree valley, with kinetic scales f and g , the polynomial proportional to $\text{Tr}(M^4)/\Lambda^8$ is $[h + 2\kappa D(c')^2]^2/f^4 + 8\kappa^2 D(c')^2/(f^2 g^2) + 4\kappa^2/g^4$. The derivative is again reduced modulo P . All sixteen declared angular-only and two-scalar parameter cases give nonzero remainders. The calculation includes the coefficient scalar instead of silently integrating it out before evaluating the loops.

Finite scalar-loop displacements

$$\Delta V = \frac{m^4}{64\pi^2} \left[\log \frac{m^2}{\mu^2} - \frac{3}{2} \right], \quad \Delta \theta = -\frac{\Delta U'}{U''}$$

The scalar-only numerical calculation uses the local positive-mass $\overline{\text{MS}}$ expression $\Delta V = m^4 [\log(m^2/\mu^2) - 3/2]/(64\pi^2)$. The renormalized additional finite Wilson coefficients are initialized to zero at the stated μ . Every example declares Λ/f , the lock coefficient and μ/m_{tree} . Resetting that boundary condition at another μ defines another example; it is not a claim that a fixed physical theory depends on the arbitrary renormalization scale after consistent running.

At $\theta_0=66$ degrees, $U''=12.2092424208$ and $U'''=-140.847974475$. The first-order angular displacement is $-\Delta U'(\theta_0)/U''(\theta_0)$. For $\Lambda/f=0.1$, no added lock, and $\mu=m_{\text{tree}}$, the numerical stationary point is approximately 65.997446321 degrees. The angle-based coefficient changes by about -0.2663 percent. The small-ratio displacement scales as $(\Lambda/f)^4$. The derivative and curvature are independently checked against finite differences, and the nonlinear stationary solution is compared with its linear prediction.

The sweep retains sixty angular examples. One leaves the physical positive-depth CKM chart; it remains in the results with a null CKM forecast rather than being dropped. Eighty of eighty-one declared two-scalar examples produce a controlled local stationary solution with a positive numerical Hessian. The unsuccessful case is retained with its parameters and failure reason. These calculations certify neither a global quantum minimum nor a cosmological occupation probability.

At the tree portal $C=1-2 \cos(12 \theta)$, $dC/d\theta=22.8253563911$ at the selected angle. A one-percent coefficient tolerance corresponds locally to about 0.00958804 degrees. Adding a large classical locking term is not automatically a radiative solution: it changes the mass and its derivatives, which also changes its own loop torque. The stored sweeps illustrate this effect directly.

A cubic supersymmetric constraint circuit

$$z_n = z^n, \quad \Phi_{60}(z) = z^{16} + z^{14} - z^{10} - z^8 - z^6 + z^2 + 1 = 0$$

Introduce sixteen gauge-singlet chiral fields $Z_1, \dots, Z_{16}$, with $Z_1=Z$, and sixteen drivers $D_2, \dots, D_{16}, D_{\text{star}}$. All Z fields have R charge zero; all drivers have R charge two. The declared CP action is complex conjugation and the chosen coefficients are real. A common real coupling can multiply the superpotential. With a real positive reference mass M , the base superpotential is $W = \sum_{n=2}^{16} D_n (M Z_n - Z_{n-1}) + M D_{\text{star}} [Z_{16} + Z_{14} - Z_{10} - Z_8 - Z_6 + Z_2 + M]$. Every term has mass dimension three.

At $D=0$, F -flatness gives $z_n = z^n$, where $z_n = Z_n/M$, and $\Phi_{60}(z) = z^{16} + z^{14} - z^{10} - z^8 - z^6 + z^2 + 1 = 0$. There are sixteen simple complex roots. They have unit modulus and are the primitive order-60 roots. This is an explicit renormalizable circuit implementing the algebraic equation; it avoids introducing a degree-sixteen superpotential interaction in the fundamental variables.

The constraint Jacobian J has nonzero determinant at every root. Its determinant divided by $\Phi_{60}'(z)$ is minus one in the declared ordering. Thus the chosen vacua are isolated. The full holomorphic mass matrix has blocks $[[0, J^T], [J, 0]]$, and every singular value of J occurs twice. With canonical kinetic terms, each pair has four real scalar modes and four fermionic spin modes with equal masses. The unbroken-SUSY one-loop contribution cancels at the F -flat zeros.

Perturbative protection here is the Wilsonian superpotential non-renormalization result for an initialized, globally supersymmetric sector, with a nonsingular positive Kahler metric. It does not force the initial coefficient choices. Kinetic corrections do not remove a zero of all F terms, but they change spectra and matter when supersymmetry is broken. This distinction follows the Wilsonian treatment of non-renormalization, rather than treating a generic 1PI potential as an unchanged superpotential [2].

What the declared symmetry actually permits

For the base singlet sector, the complete holomorphic superpotential basis through field degree three has 2,448 monomials. R charge forces one driver, followed by zero, one or two neutral fields: $16(1+16+136)$. The selected circuit uses 37 monomials. CP and the declared R symmetry allow many other coefficients. They protect neither the chosen sparse pattern nor its relative integer values by themselves. The general CP-even quadratic Kahler metric contains 272 real parameters in the two neutral/driver blocks; the numerical mass calculations initialize it canonically.

A real change of the D_{star} tadpole changes $\Phi_{60}(z)$ to $\Phi_{60}(z)+\epsilon$. This deformation is allowed by the declared symmetries and changes both phase and radius. It demonstrates freedom in the initialized theory. It is not a claim that perturbative supersymmetric loops generate that absent term in the Wilsonian superpotential.

For a separate one-complex-field comparison, all CP-even operators $\text{Re}(z^a \bar{z}^b)$, $a \geq b$, $1 \leq a+b \leq 12$, are enumerated. There are 48 operators when no rotation symmetry is imposed; 42 have a nonzero carrier torque. The six desired harmonic numbers have gcd one. Among all cyclic rotations tested, orders two through sixty cannot permit every desired harmonic. This is a precise obstruction for that one-field CP-plus-cyclic class, not for arbitrary multi-field finite groups.

A holomorphic golden-coefficient extension

$$T = 2 - z^4 - z^6 + z^{14}, \quad T^2 - 3T + 1 = 0$$

The inverse phase needed for the original coefficient can be represented in the carrier without complex conjugation. Since $z^{60}=1$ on $\Phi_{60}=0$, $z^{-12}=z^{48}$. Exact reduction gives $T=1-z^{12}-z^{48}=2-z^4-z^6+z^{14}$ modulo Φ_{60} . It obeys $T^2-3T+1=0$. At the root with phase 66 degrees, $T=(3-\sqrt{5})/2$, with zero imaginary part.

Two additional chiral fields T and D_{gold} implement this linearly: $W_{\text{gold}}=M D_{\text{gold}}[T-2M+Z_4+Z_6-Z_{14}]$. T has R charge zero and D_{gold} has charge two. The extended model has 34 chiral fields and remains renormalizable. Its general allowed singlet superpotential through degree three contains 2,907 monomials. The extension fixes a golden spurion value in the selected initialized vacuum, while preserving the sixteenfold supersymmetric degeneracy. It does not fix a quark Wilson coefficient or explain why the twelfth power is the desired coefficient operator.

Actual field relaxation under energy selection

To lift the degeneracy, the worked scalar deformation is $V_{\text{soft}}/M^4=g_W^2 \epsilon \sum a_n \text{Re}(z_n)$, with the six nonconstant Fourier coefficients a_n . This is an explicitly chosen supersymmetry-breaking scalar tadpole pattern. At exact circuit roots its leading energy is the original angular potential times $g_W^2 \epsilon$. The full fields are then allowed to relax; they are not constrained to remain equal to powers of a unit-modulus number.

For fixed z_1 , the other fifteen base neutral fields enter the F-term potential quadratically. They are eliminated by linear least squares, including the scalar tadpoles. The remaining complex z_1 is optimized in both its real and imaginary coordinates. All sixteen branches are continued for $\epsilon=10^{-8}, 10^{-6}, 10^{-4}, 10^{-3}$. The branches descended from root powers eleven and forty-nine have the lowest energies in every declared continuation. Every retained branch has a positive reduced tree Hessian. This is a comparison of local continued vacua, not a proof that the

full polynomial field potential has no distant competing vacuum or runaway.

At $\epsilon=10^{-4}$, the positive branch has phase 66.0005874177 degrees and radius 1.00002214970. The angle-only readout gives $C=0.382200029974$. The nonholomorphic real auxiliary readout $1-2 \operatorname{Re}(z_{12})$ gives 0.381955432748. The holomorphic extension gives $T=0.381669772701-0.000178406720 i$. These are different operators once the fields relax. A quark interaction must specify which operator it actually uses. Choosing a real auxiliary readout in a holomorphic superpotential would require a different construction; it cannot be substituted silently.

Full-circuit leading quantum calculation

$$A = J^\dagger J, \quad B = \sum_k \overline{F_k} H_k$$

The full quantum calculation includes the 34 chiral fields of the golden extension. On the R-symmetric $D=0$ slice, let F be the vector of driver constraints, J their Jacobian with respect to the seventeen neutral fields, and H_k their constant quadratic Hessians. Define $A=J^\dagger J$ and $B=\sum \text{conjugate}(F_k) H_k$. In canonical $\sqrt{2}$ real and imaginary neutral coordinates, the scalar squared-mass matrix is $[[\operatorname{Re}(A+B), -\operatorname{Im}(A)-\operatorname{Im}(B)], [\operatorname{Im}(A)-\operatorname{Im}(B), \operatorname{Re}(A-B)]]$. The driver scalar block has the corresponding unsplit singular-value spectrum.

The driver scalar trace cancels half of the fermion trace. The complete singlet one-loop potential therefore equals $g_W^4/(64 \pi^2)$ times $[\sum (34 \text{ neutral real modes}) b_i^2 (\log(g_W^2 b_i/\mu^2)-3/2)-2 \sum (17 \text{ Jacobian modes}) a_i^2 (\log(g_W^2 a_i/\mu^2)-3/2)]$. This expression accounts for all 68 real scalars and 34 Weyl fields, not just the light angular mode. It is evaluated in a DRbar example with canonical renormalized Kahler metric and specified scalar tadpoles at the reference scale. There are no gauge fields in this singlet-sector calculation.

The gradient uses eigenvector contractions with the analytic derivatives of A and B . It is independently checked against potential differences. At exact F-flat roots, the potential and gradient cancel to numerical roundoff. At the softly selected vacuum, the gradients need not cancel. The quantum stationary equations are solved in all thirty-four real neutral coordinates; the corrected neutral-sector Hessian is numerically positive. An independent expression for the complete 68-scalar, 34-Weyl effective potential also computes the quantum driver-block Hessian by finite differences. The unbroken common driver R phase makes the neutral-driver block vanish at zero drivers. Both corrected blocks have positive minimum eigenvalues in all nine declared cases. Step-size comparison checks the driver Hessian, while the agreement of the independent full and reduced potentials checks the mode counting. These are numerical local stability checks at one loop, rather than exact certificates or a higher-loop theorem.

Nine epsilon/coupling examples are retained: $\epsilon=10^{-6}, 10^{-4}, 10^{-3}$ and $g_W=0.1, 0.3, 1$. All produce controlled numerical neutral-sector minima. Leading tree-plus-one-loop energy comparisons among all sixteen continued branches retain root powers eleven and forty-nine. At $\epsilon=10^{-4}$ and $g_W=0.3$, the phase becomes 66.0005860473 degrees, a quantum displacement of approximately -1.37037 times 10^{-6} degrees relative to the relaxed tree minimum. The coefficient field becomes $T=0.381670086337-0.000178010587 i$. The largest actual neutral-coordinate quantum displacement is about 4.67 times 10^{-7} . These are chosen weak-coupling examples, not experimental uncertainties or a derived cosmological scale.

Quark operators: hierarchy without an enforced coefficient

The declared holomorphic quark sector uses two flavor charges. The left-handed doublets Q_i have charges (1,1),(0,1),(0,0), while u_i^c and d_i^c have the opposite charges. H_u, H_d and all neutral circuit fields have zero flavor charge. A has charge (-1,0), B has charge (0,-1), and Q, f^c have R charge one. The scalar A and B backgrounds remain free inputs. No anomaly-free gauged flavor completion or mechanism fixing their magnitudes is claimed.

All allowed operators $Q_i H f_j^c A^a B^b$ times neutral monomials are enumerated through superpotential field degree six. There are 6,460 operators with the sixteen base neutral fields, and 7,560 when T is included. The up and down sectors have independent real coefficients before spontaneous CP breaking. The pure singlet enumeration and this Yukawa enumeration are complete within their declared sectors and degree bounds; they are not an enumeration of all Standard Model, lepton or higher-Kahler interactions.

In particular, $Q_1 H d_2^c A$, $Q_2 H d_3^c B$, and $Q_1 H d_3^c A B$ carry independent coefficients. Allowing their spurion powers does not relate those coefficients. A T insertion can multiply the third operator, but its independent coefficient and the lower-degree $A B$ operator remain allowed. Neutral fields can also appear in diagonal and up-sector entries. The phase of the scalar vacuum therefore does not automatically become the standard CKM phase. Higher-Kahler interactions such as $Q_1^\dagger Q_2 A^\dagger/M$ are allowed and can alter canonical normalization.

A derived texture counterexample

$$V_{\text{CKM}} = U_u^\dagger U_d, \quad U_f^\dagger Y_f Y_f^\dagger U_f = \text{diag}(y_f^2)$$

The worked trial uses $Y_u = \text{diag}(u_1, u_2, u_3)$ and an upper-triangular Y_d with entries $Y_{12} = a d_2$, $Y_{23} = b d_3$ and $Y_{13} = c a b \exp(-i \theta) d_3$. The diagonal spectra are declared positive numerical inputs. This is a selected member of the allowed operator family. Up-sector off-diagonal terms and most neutral insertions are initialized to zero; the symmetry does not forbid them.

The matrices are diagonalized through the eigenvectors of $Y Y^\dagger$. For each independent c , only a and b are matched to the two existing anchors $|V_{us}| = 0.22431$ and $|V_{cb}| = 0.0411$. The values of $|V_{ub}|$ and CP observables are then consequences of the matrix, without fitting them. At $c = \phi^{-2}$ and $\theta = 66$ degrees, the texture gives $|V_{ub}| = 0.00360299463$ and effective depth 0.390811666690, rather than the nominated chart's 0.00352144218 and 0.381966011250. Its standard CKM phase is approximately 65.9786293 degrees. Finite diagonalization effects matter even in this simple hierarchical matrix.

Changing the independent coefficient while retaining the same two anchors changes $|V_{ub}|$ substantially. The stored six-coefficient sweep supplies explicit examples. This is a mathematical obstruction to inferring the exact relation $s_{13} = C s_{12} s_{23}$ from charge arithmetic alone. A stronger quark construction must relate Wilson coefficients, kinetic normalization and matrix diagonalization, or predict the correction instead of absorbing it into a fit.

Scale evolution as a separate matching test

$$16\pi^2 \frac{dY_u}{d \log \mu} = \left[\frac{3}{2} (H_u - H_d) + \alpha_u I \right] Y_u$$

The implementation also integrates one-loop Standard Model matrix Yukawa equations using $t=\log(\mu)$, left-row Yukawa convention and SU(5)-normalized g_1 . In this convention $\beta_u=[(3/2)(H_u-H_d)+T_{\text{gauge},u}]Y_u/(16\pi^2)$, with the corresponding down and lepton equations and the one-loop gauge equations. The source's $\log(\mu^2)$ convention is converted explicitly [3]. Rephasing invariance, a common-left-basis covariance check and the top-only limit independently check the implementation.

The example initializes the original exact CKM chart with declared illustrative spectra and gauge couplings. At $\mu/\mu_0=10^8$, its effective depth differs from ϕ^{-2} by about -8.62 times 10^{-5} relatively, and its standard phase differs by about -0.00349036 degrees. The absolute $|V_{ub}|$ also changes as the anchors run. No measured matching scale is supplied. This is a standalone SM evolution diagnostic, not a threshold-matched evolution of the supersymmetric circuit. It shows that a claim of an exact flavor relation must identify the scale at which it is imposed.

What is now supported and what remains physical

The exact supports are the carrier algebra, the stationary-family identities, the global-winner preservation argument, the scalar-counterterm remainder and Bezout witness, and the complete declared finite operator spaces. The supersymmetric protection is a conditional application of the established Wilsonian non-renormalization result to the supplied globally supersymmetric circuit. The mass spectra, relaxed fields, one-loop extrema, quark diagonalizations and SM evolution are numerical computations with explicit parameters and limitations. No new Lean theorem is claimed.

The constructed model answers whether an initialized protected equation can be implemented by renormalizable fields and whether a specified weak selector produces a controlled nearby vacuum. It does so. The outstanding physics is the origin of the sparse initialized coefficients and selector terms, a quark interaction enforcing or replacing the nominated exact mixing rule, consistent kinetic and threshold matching, higher-loop stability, and a vacuum history. The branch comparison does not choose the positive CP sign. A complete theory of strong CP or baryogenesis has not been supplied.

Reproduction and retained calculations

Run `PYTHONPATH=python OPENBLAS_NUM_THREADS=1 python python/develop_flavor_completion.py` from the repository root, then the corresponding `plot_flavor_completion.py`. Run unittest discovery with `test_flavor_completion.py` for the new mathematical and numerical checks, or `test_flavor*.py` for the combined flavor checks. NumPy and SciPy support numerical spectra, extrema and running; Matplotlib supports the scientific figure. Exact polynomial identities and operator labels use rational arithmetic and the existing PerfectPower algebra.

The receipts retain all declared cases, including one uncontrolled two-scalar continuation and one angular example outside the positive-depth CKM chart. The two exhaustive quark-operator tables are stored as deterministic gzip-compressed JSON; ordinary gzip decompression recovers the complete tables. The manifest covers the scientific JSON outputs, including those compressed tables. Fresh-copy reproduction checks those outputs in the same numerical runtime. A byte-identical replay is evidence of reproducibility in that environment; the tests provide separate mathematical and numerical checks. PDF generation uses ReportLab and standard installed fonts. Source code, prior repository work, raw receipts, diagrams and the monograph are included in the complete handoff archive.

The preceding vacuum-completion run completed 841 tests successfully, with four skips. Its 25 new completion tests cover the exact carrier obstruction, circuit elimination and mode pairing, independent loop gradients, corrected driver Hessian, complete declared operator enumeration, texture counterexamples and scale-evolution covariance. A fresh archive extraction reproduced all twenty scientific receipt, manifest and figure files byte for byte in the declared runtime. No new Lean theorem was added in that push.

Primary references

- [1] S. Coleman and E. Weinberg, Radiative Corrections as the Origin of Spontaneous Symmetry Breaking, *Physical Review D* 7, 1888 (1973). <https://doi.org/10.1103/PhysRevD.7.1888>. This supplies the effective-potential framework; the worked scalar and circuit schemes are stated explicitly above.
- [2] N. Seiberg, Naturalness Versus Supersymmetric Non-renormalization Theorems (1993), <https://arxiv.org/abs/hep-ph/9309335>. The relevant distinction is protection of a chosen Wilsonian superpotential, with separate kinetic and supersymmetry-breaking effects. It does not derive the chosen flavor coefficients.
- [3] A. V. Bednyakov, A. F. Pikelner and V. N. Velizhanin, Three-loop SM beta-functions for matrix Yukawa couplings (2014), <https://arxiv.org/abs/1406.7171>. The one-loop matrix conventions and scale conversion are the parts used here.
- [4] I. de Medeiros Varzielas, D. Emmanuel-Costa and P. Leser, Geometrical CP violation from non-renormalisable scalar potentials (2012), <https://arxiv.org/abs/1204.3633>. This is a comparison for symmetry-constrained higher-order operators preserving particular vacua; it does not establish the present order-60 construction.


Shared mediators and the alignment boundary

What this mediator push establishes

The next step now has an explicit non-Abelian quark mediator construction and an exact tree matching calculation that includes the light right-handed kinetic normalization. The family symmetry makes each heavy mediator mass proportional to the identity and gives one contraction for each flavon column. Separately orthogonal flavon frames then determine the CKM matrix independently of the six column magnitudes and the two flavor-universal mediator masses. This is a constructive quark interaction that transmits an initialized mixing frame without the finite-mass diagonalization errors of the previous triangular texture.

It does not independently derive that frame. The accompanying scalar analysis gives a precise obstruction to the simplest alignment route. For six fundamental flavons with separate phase symmetries, fixed positive norms and exactly orthogonal up and down frames, every CP-even orientation term through scalar degree six reduces to a constant plus a linear function of $|V_{ij}|^2$. An exact rational assignment certificate proves that a permutation always attains its global minimum on the frame manifold. A generic nondegenerate cost therefore cannot select the required hierarchical CP-violating mixing. Degenerate costs can leave mixed flat directions; those are not a unique prediction.

I also tested a more favorable alternative in which an adjoint mediator mass contains initialized neighboring mixings, a golden 1-3 entry and the 66-degree phase. Both up and down adjoint couplings are allowed independently by the symmetry. Exact tree matching and canonical normalization were performed before extracting CKM quantities. Thirty of thirty-six declared cases satisfy the numerical control criteria. Every accepted case fails the golden target by more than one percent, despite matching the two existing anchors. The six unsuccessful or uncontrolled cases are retained with their reasons. The construction is therefore a tested candidate and a boundary result, not a completed predictive theory of flavor.



The exact observable target

$$C_{\text{eff}} = \frac{w(1-w^2)}{uv} = \phi^{-2}$$

Write $u=|V_{us}|$, $v=|V_{cb}|$ and $w=|V_{ub}|$. The nominated physical relation is $C_{\text{eff}}=w(1-w^2)/(uv)=\phi^{-2}$, where $\phi=(1+\sqrt{5})/2$. This follows from the original standard-angle rule $s_{13}=C s_{12} s_{23}$ because $u=s_{12} c_{13}$, $v=s_{23} c_{13}$ and $w=s_{13}$. It is already rephasing invariant. Relating a Yukawa matrix entry to other entries is insufficient unless its diagonalization yields this observable equation.

For $p=u^2$, $q=v^2$ and $r=w^2$, define $A=r(1-r)^2$. Eliminating C from $C^2-3C+1=0$ gives $A^2-7pqA+(pq)^2=0$. The positive physical branch $C_{\text{eff}}<1$ selects ϕ^{-2} ; the other positive root is ϕ^2 . The equation does not fix a CP phase. In raw fixed-norm flavon overlaps its formal maximum scalar degree is twenty-four. Squaring it into an energy term would encode the desired result by construction and would not explain its origin.

The existing anchors remain $u=0.22431$ and $v=0.0411$. The small-branch target is $w=0.00352144218276$. No new experimental data or fit to $|V_{ub}|$ or CP measurements was introduced. All numerical examples use declared illustrative quark spectra and dimensionless mediator units. A claim of an exact relation still needs a specified matching scale and subsequent threshold and running calculations.

An exact Abelian obstruction for the initialized circuit

Consider an additional ordinary diagonal Abelian phase action preserving every monomial of the initialized singlet superpotential, with its mass M and numerical coefficients neutral. The recursion terms force $q(Z_n)=n q(Z_1)$ and $q(D_n)=-n q(Z_1)$. The constant D_{star} term forces $q(D_{\text{star}})=0$. Its remaining terms require $n q(Z_1)=0$ for $n=2,6,8,10,14,16$. Their greatest common divisor is two, so the general character relation is $2 q(Z_1)=0$.

The golden extension has a constant D_{gold} term, forcing $q(D_{\text{gold}})=0$. Its linear $D_{\text{gold}} T$ term then forces $q(T)=0$. Hence an operator O and $O T$ have the same additional charge. Such a charge rule cannot forbid $Q_1 H d_3^c A B$ while allowing $Q_1 H d_3^c c A B T$. The implementation derives the monomial constraints from the existing circuit rather than a separate assumed polynomial, and checks cyclic groups of orders one through 120. The focused test independently extends those checks through order 240.

This result is deliberately restricted. Charged mass or coupling spurions, symmetries mixing fields, and non-Abelian representations can change the premises. Treating the initialized coefficients as transforming spurions would supply a different model and would require its own alignment and coupling audit. The obstruction does not rule out flavor symmetry in general.

The declared non-Abelian mediator sector

Use the Standard Model gauge group, a formal global $SU(3)_F$ family symmetry, six independent $U(1)$ flavon phase symmetries and CP. Q is a family anti-triplet. The three right-handed-conjugate up fields and the three right-handed-conjugate down fields are family singlets. There are six complex gauge-singlet fundamental flavons F_{ui} and F_{di} . Each carries its own phase charge $+1$, while its associated right-handed-conjugate quark carries -1 . The Higgs and Q have zero phase charge.

Introduce heavy X_u and X_d family anti-triplets and X_u^c and X_d^c family triplets. X_u and X_u^c are vectorlike color-triplet weak-singlet fermions with hypercharges $+2/3$ and $-2/3$. X_d

and X_d^c have hypercharges $-1/3$ and $+1/3$. Each family triplet gives three heavy Dirac species, for six species overall. These pairs are vectorlike under the Standard Model. The global family group is not claimed to be an anomaly-free gauged completion; its Goldstone and cosmological questions are outside this calculation.

In ordinary left-handed Weyl notation, the allowed core interactions are $M_f X_f \cdot X_f^c$, $h_f Q \cdot H_f X_f^c$ and $\sum_i \lambda_{fi} X_f \cdot F_{fi} f_i^c$, plus their Hermitian conjugates. H_u is the Higgs doublet and H_d is its conjugate doublet. This notation is an ordinary non-SUSY mediator model. It is not a combined softly broken completion of the previously constructed 34-chiral-field singlet sector.

The operator audit exhausts fermion mass bilinears and two-fermion-one-scalar Yukawa invariants in this declared quark sector. It checks Standard Model gauge representations, family representations and all six phase charges. With the seventeen neutral scalar backgrounds Z_1 through Z_{16} and T , including their conjugates, there are 78 independent invariant coefficients. Two are bare universal masses, two are Higgs vertices, six are flavon-column vertices and 68 are neutral-scalar universal-mass vertices. CP permits real coefficients; it does not equate them. Lepton interactions and a complete scalar potential for all neutral backgrounds are outside this finite audit.

Adding one complex phase-neutral family adjoint Σ and its conjugate adds four invariants, giving 82. They are $X_u \Sigma X_u^c$, $X_u \Sigma^\dagger X_u^c$, $X_d \Sigma X_d^c$ and $X_d \Sigma^\dagger X_d^c$. Up and down coefficients are independent. In the Hermitian background used below the two coefficients in each sector combine into one real number, but the symmetry does not impose any relation between the sectors or between an adjoint coupling and a singlet mass.

Canonical tree matching without a small-flavon expansion

$$A = M^{-1}L, \quad K = I + A^\dagger A, \quad Y = -hAK^{-1/2}$$

Let L_f be the 3 by 3 matrix of flavon columns multiplied by their λ_{fi} couplings, and let M_f be the invertible heavy mediator mass matrix. At zero Higgs background the heavy fermion row block is $[L_f, M_f]$. Define $A_f = M_f^{-1} L_f$, $K_f = I + A_f^\dagger A_f$ and $R_f = K_f^{-1/2}$. A normalized basis for the three light right-handed fields is $N_f = [R_f; -A_f R_f]$. Direct multiplication gives $[L_f, M_f]N_f = 0$ and $N_f^\dagger N_f = I$.

The Higgs vertex projected into that normalized null space gives $Y_f = -h_f A_f R_f$. This is exact at tree order at unbroken electroweak matching, to all orders in the ratio of flavon mixing to heavy mass. It is not the unnormalized expression $-h_f M_f^{-1} L_f$. No loop threshold is included. At finite Higgs vacuum value, higher-dimensional Higgs operators and mixing with weak-singlet heavy fermions also affect the charged current. The dimension-four effective CKM matrix is unitary; the finite-Higgs full theory need not give an exactly unitary light 3 by 3 submatrix.

The focused checks verify the normalized null frame and Yukawa projection directly. They also compare against independent singular values of the complete 6 by 6 fermion mass matrix $[[0, h_f v_H I], [L_f, M_f]]$. The three light singular values divided by v_H converge to those of the matched Y_f . A separate finite-Higgs test compares the light charged-current overlaps with the matched CKM limit and checks their small departure from unitarity. Noncommuting mass matrices are tested under a common family-basis rotation, so the checks do not rely only on diagonal mediator masses.

What universal masses can and cannot transmit

For a universal mass $M_f = m_f I$ and a general singular decomposition $L_f = U_f \text{diag}(s_f) W_f^\dagger$, the canonical Yukawa matrix has the same left eigenvectors U_f . Its singular values are $|h_f| s_f / \sqrt{|m_f|^2 + s_f^2}$. This map is strictly increasing for positive s_f , preserving nondegenerate mass ordering. Thus changing a flavor-universal mediator mass, even by a complex golden or cyclotomic background, cannot change the mixing angles or physical CKM phase of fixed L_u and L_d . This statement does not require orthogonal flavon columns.

With separately orthogonal columns, $L_f = U_f \text{diag}(\ell_f)$, the stronger separation holds: changing each individual column magnitude only changes its associated quark singular value. While retaining the assigned nondegenerate mass ordering, the CKM matrix is $U_u^\dagger U_d$ and is independent of those magnitudes and the universal masses. Twelve declared examples initialize $U_u = I$ and U_d from the original golden CKM chart, then vary column magnitudes and real or complex universal masses. They retain the nominated coefficient and 66-degree standard phase to numerical precision. These are conditional transfer checks. The down frame was copied from the target chart and has not been derived from a symmetry-controlled potential.

A separate three-case test uses nonorthogonal random columns and changes only universal masses. It confirms the general unchanged-eigenvector result while the spectra change. Both results narrow the physical role a golden field can play: placing it only in a universal heavy mass does not orient the CKM matrix. Placing its phase as an overall Yukawa-column multiplier also fails, because $(Y \text{diag}(\exp(i \theta_j)))(Y \text{diag}(\exp(i \theta_j)))^\dagger = Y Y^\dagger$. A physical CP phase must influence relative left-handed directions or interfering flavor structures.

The complete low-degree triplet alignment basis

For the six complex triplets alone, write $G_{ab} = F_a^\dagger F_b$. Independent phase invariance requires each triplet label to have equal occurrences with and without conjugation. At scalar degree two there are six norms. At degree four there are 21 products of norms and fifteen squared overlaps $|G_{ab}|^2$, giving 36 quartic invariants. Including the quadratic terms, the renormalizable triplet potential has 42 independent real coefficients. One Standard Model Higgs adds its quadratic term, its quartic term and six Higgs-norm/flavon-norm portals, giving fifty terms in that restricted scalar sector.

At degree six the CP-even basis has 56 triple products of norms, ninety norm-times-squared-overlap terms and twenty real Gram triangles $\text{Re}(G_{ab} G_{bc} G_{ca})$ with distinct labels. Products of balanced SU(3) epsilon tensors are Gram determinants and reduce to this basis. Unbalanced epsilon contractions are forbidden by the six separate phase symmetries. The first Gram-rank determinant relation for six triplets in three dimensions has degree eight, so it does not remove any of these degree-six contraction types.

Now restrict to positive fixed norms and exactly orthogonal U_1, U_2, U_3 and D_1, D_2, D_3 frames. Overlaps within each frame vanish. Every three-label triangle contains two labels from one frame, so every Gram triangle vanishes. Norm products are constant. The surviving squared cross overlaps are proportional to $B_{ij} = |U_i^\dagger D_j|^2 = |V_{ij}|^2$. Norm factors at degree six only modify their coefficients. Therefore the complete orientation dependence through degree six is $E = \text{constant} + \sum_{ij} K_{ij} B_{ij}$.

Fixed norms and exact separate orthogonality are assumptions of this theorem. Arbitrary finite quartic coefficients in the full potential can distort the frames, and a model outside this regime requires a new calculation. An adjoint introduces new orientation tensors and is also outside this theorem. Neutral singlets can change norm-dependent parameters; multiplying a quartic

overlap by a singlet would exceed renormalizable degree and does not create a new degree-four orientation tensor.

A rational global certificate for the orientation obstruction

$$E = \sum_i \alpha_i + \sum_j \beta_j + \sum_{ij} R_{ij} B_{ij} \geq E_{\text{permutation}}$$

For any unitary relative frame, B is doubly stochastic: its entries are nonnegative and each row and column sums to one. For each declared rational cost matrix K , the implementation finds row and column numbers α_i and β_j with $\alpha_i + \beta_j \leq K_{ij}$. It also finds a permutation attaining the same sum $\sum_i \alpha_i + \sum_j \beta_j$. Define reduced costs $R_{ij} = K_{ij} - \alpha_i - \beta_j$, which are exactly nonnegative rational numbers.

Then $E = \sum_i \alpha_i + \sum_j \beta_j + \sum_{ij} R_{ij} B_{ij}$ is bounded below by the permutation energy. A permutation matrix is itself unitary, so the bound is attained within the physical frame manifold. The receipt supplies the rational dual numbers, reduced costs and all winning permutations. Tests check eighty further rational cost matrices independently against all six assignments and compare the identity against nontrivial random unitaries.

This proves that a permutation is always a global minimizer of the restricted linear orientation energy. With a unique minimum assignment, no nonpermutation B attains the same energy. For degenerate costs, a mixed CP-violating frame can instead belong to a flat minimum set. The constant-cost example explicitly retains such a direction. The correct conclusion is that this potential cannot uniquely select the nominated nonpermutation CKM matrix in the stated regime, not that every possible minimum must have zero CP violation.

The first higher-degree CP capability example

Scalar degree eight permits $|F_{ui}^\dagger F_{dj}|^4$ and real four-overlap cycles. With equal fixed norms, the chosen positive sum of fourth powers reduces to $\sum_{ij} B_{ij}^2$. Each row obeys $\sum_j B_{ij}^2 \geq 1/3$, so the total is at least one. Equality requires every $B_{ij} = 1/3$. The unitary three-dimensional Fourier frame reaches the bound and has $|J| = 1/(6 \sqrt{3})$, with a conjugate CP partner.

This is an explicit CP-even orientation interaction that can select nonzero physical $|J|$ when paired with nondegenerate quark column magnitudes. It selects democratic mixing, with every CKM magnitude $1/\sqrt{3}$, and fails the hierarchical target. Its coefficient pattern is chosen; other degree-eight invariants are independently allowed under the declared symmetry. No renormalizable UV generation of that pattern or connection to the protected 66-degree singlet vacuum has been provided. The example demonstrates what new orientation dependence becomes possible at degree eight, while leaving the actual golden alignment unresolved.

The favorable adjoint candidate and its failure

The numerical candidate uses a Hermitian traceless adjoint background Σ with entries $\Sigma_{12}=a$, $\Sigma_{23}=b$ and $\Sigma_{13}=\phi^{-2} a b \exp(-i \theta)$, together with their conjugates and zero diagonal. θ is initialized at 66 degrees. The mediator masses are $M_u = I + g_u \Sigma$ and $M_d = I + g_d \Sigma$. The columns before filtering are diagonal, using the same illustrative hierarchies as the preceding quark calculation. This gives the golden factor and phase a direct place in a flavor-dependent mediator, which a universal mass could not supply.

The assumed adjoint background is deliberately favorable and explicitly initialized. No scalar potential producing it is claimed. Both g_u and g_d are independent allowed couplings. Canonical tree matching is performed for each sector, followed by exact numerical diagonalization of $YY^\dagger$. Only a and b are matched to $|V_{us}|$ and $|V_{cb}|$. They are not tuned to $|V_{ub}|$ or any CP observable.

The declared sweep uses $g_u=0,0.1,0.2$; $g_d=0.3,0.5,0.7,1$; and column scales $0.1,1,3$. A case is accepted only when the two anchor residuals are below 10^{-9} , both mediator matrices have minimum singular value at least 0.2 and $\max(a,b)\leq 1$. Thirty cases are accepted and six fail the matching or control criteria. Accepted C_{eff} ranges from approximately 0.941676 to 1.595694, against the golden target 0.381966. All accepted cases fail the target at the one-percent level. The input phase also does not become the standard CKM phase automatically. Four conjugate-pair checks retain magnitudes and reverse J .

Unnormalized Yukawa predictions are retained for the same backgrounds so that the kinetic correction can be inspected. They are not substituted for the canonical result. The failure is not a theorem against all adjoint models; it demonstrates that this favorable entry pattern and the allowed shared-mediator couplings do not enforce the desired observable relation. A successful adjoint construction would need a derived vacuum and additional relations among its allowed couplings, followed by the same normalization and mixing checks.

Reproduction, validation and the next interaction target

Run `PYTHONPATH=python OPENBLAS_NUM_THREADS=1 python python/develop_flavor_mediator.py`, followed by `python python/plot_flavor_mediator.py` with the same environment. The runner uses only the existing $|V_{us}|$ and $|V_{cb}|$ means. It retains exact charge, operator and rational dual certificates, the conditional transfer cases, all adjoint attempts and the scientific figure. NumPy and SciPy support the numerical matching; the exact charge and assignment statements use integer or rational arithmetic. There are no new Lean theorems or newly acquired experimental datasets.

The twenty focused tests cover the charge obstruction, independent operator-list construction, exact rational assignment witnesses, canonical null-space matching, the complete fermion mass comparison, family-basis covariance, CP conjugation, universal-mass invariance and the failed golden enforcement. They accompany the repository-wide suite and a fresh-extraction byte comparison of the retained scientific artifacts. The integrated repository-wide run completed 879 tests successfully, with four skips. A fresh archive extraction reproduced all 32 scientific receipt, manifest and figure files from both flavor investigations byte for byte. The full test and replay results are recorded in the validation receipt for this push.

The resulting research target is more specific than adding a golden Yukawa coefficient. The quark mediator interaction can already transmit an orthogonal frame exactly, so the remaining positive construction is a symmetry-controlled relative frame that produces the golden amplitude relation and the physical CP phase. Within the tested phase-symmetric triplet class, a uniquely selected nonpermutation frame requires new orientation dependence beyond degree six or a change of fields or assumptions. Higher-degree interactions, additional non-Abelian representations or nonorthogonal-frame dynamics are possibilities to test; none is asserted here to solve the problem. Threshold matching, running, a consistent combined singlet/flavor theory and a vacuum history remain separate requirements.

Primary comparisons and their limits

S. Antusch, S. F. King, M. Malinsky and M. Spinrath, Quark mixing sum rules and the right unitarity triangle, <https://arxiv.org/abs/0910.5127>. Its texture-zero sum-rule route motivates deriving mixing from both Yukawa sectors. It does not establish the present golden relation.

R. Alonso, M. B. Gavela, L. Merlo and S. Rigolin, On The Potential of Minimal Flavour Violation, <https://arxiv.org/abs/1103.2915>. Its analysis provides context for restrictions of low-degree dynamical flavor potentials. The six-triplet fixed-frame obstruction above has separate explicit premises and its own rational certificate; no claim of duplicating every model in that paper is made.

J. Gehrlein, J. P. Oppermann, D. Schafer and M. Spinrath, An $SU(5) \times A_5$ Golden Ratio Flavour Model, <https://arxiv.org/abs/1410.2057>. It provides an example of an explicit renormalizable flavor and messenger construction. Its golden mixing result concerns the neutrino sector, not our nominated CKM coefficient. Only the primary abstracts and metadata were used for these comparisons.

Recovered M22 geometry and the flavor frame

What was recovered

The repository preserves selected original files from WILSON_FORMATION_FORGE v0.26.0, including its M22/A5 Wilson algebra, Schur-selector theorems, transport-category results, and a 1,023-word exact integer ledger. The provenance manifest is `galois_merge/receipts/vendor_sources.json`. This is a verified subset of the old work, not recovery of every later Atlas or EQ-LAB release. The saved ledger has SHA-256 `1351e514fcb5f23832f65665e0a12d2e172f8131c34875de0e03cddc5aec3093`. The large original orbital tensor is absent from the subset; the ledger is nevertheless sufficient to reconstruct its two-generator algebra faithfully.

The carrier is $V = C[M22/A5]$, of dimension 7,392. Its commutant $H = \text{End}_{M22}(V)$ has dimension 152. The complex simple block sizes are 1,3,3,2,5,5,5,3,3,6. Their squares sum to 152. Thus there are four three-dimensional multiplicity carriers available for a frame calculation. They are multiplicity spaces of larger M22 representations, rather than four nontrivial three-dimensional representations of M22.

The recovered Q192 generators are $X = A49$ and $Y = -A12 - A48 - A59 + A95 + A116 + A125$. X is self-adjoint and Y is skew-adjoint. This convention differs from the earlier Q7 signed operator; no Q192 calculation here is silently identified with Q7 or with its exceptional 154 block. Likewise, the three-dimensional minus-context factor in the separate 486-dimensional transport algebra is not identified with a Wilson multiplicity carrier or a physical family triplet.

Exact reconstruction from the word ledger

The new verifier `python/develop_m22_frame.py` selects 152 independent ledger rows through word degree eight, inverts their integer coordinate matrix over the rationals, and reconstructs left and right multiplication by X and Y . It also reconstructs the orbital transpose permutation from reversed words and their skew signs. No CKM number, target phase, or golden coefficient enters this reconstruction.

Every available one-step left and right extension of all 511 words through degree eight is checked exactly against the original ledger. The transpose squares to the identity and is an actual orbital permutation. All four left/right generator commutators vanish exactly. Adjointness recovers positive integral orbital subdegrees, normalized at the identity; their sum is exactly 7,392. The recovered regular matrices are integral, not numerical fits.

The exact characteristic polynomial of X is $(t-30)^9(t-3)^{51}(t+2)^{65}(t+5)^{27}$. The exact Y factorization is recorded in `receipts/m22_frame/frame_audit.json`. In particular it contains the factors t^2+1452 and t^2+2732 relevant to the two real three-dimensional carriers, and a degree-six factor for the conjugate complex pair. Regular-module multiplicities and physical permutation-module multiplicities are different; only the shared root sets are being used here.

Unitary one-copy carrier extraction is a separate numerical step. A generic Hermitian left-multiplication selector commutes with right multiplication, so its eigenspaces supply right-invariant carriers. Word traces through degree five distinguish the conjugate complex blocks. Even trace data alone would merge that pair incorrectly. The extracted block sizes

reproduce all ten expected simple blocks. The largest reported invariance residual is below $3.3\text{e-}11$. Carrier coordinates and numerical overlap matrices are explicitly not promoted to exact algebraic coordinates.

Why equivariance alone cannot select the frame

Write V as the direct sum of R_λ tensor M_λ , where R_λ is an irreducible M_{22} carrier and M_λ is its multiplicity space. M_{22} acts as $\rho_\lambda(g)$ tensor I on this summand. Every operator I tensor A_λ commutes with M_{22} , for any matrix A_λ on M_λ . In a three-dimensional multiplicity carrier the full $M_3(\mathbb{C})$ algebra is therefore allowed by M_{22} equivariance.

For any desired unitary frame W , the two positive operators D_u and $W D_d W^\dagger$ are both permitted on such a multiplicity space. Their eigenvalues can be distinct. Their relative eigenframe is W . Consequently M_{22} equivariance alone accommodates every CKM frame and cannot force the golden relation or a particular CP phase. Canonical orbital geometry supplies distinguished candidate operators, but selecting specific orbital coefficients requires an additional interaction or residual symmetry. Algebra-generation and transport-isomorphism theorems do not perform that selection.

This is a scoped obstruction to using the unbroken commutant as the complete flavor principle. It is not a no-go theorem for M_{22} breaking, decorated incidence fields, projective carriers, or a more restrictive interaction category.

The canonical C3/C7 frame fails

The first frame test takes the up-sector Hermitian operator to be X and the down-sector operator to be iY . Positive shifts, positive overall rescalings, and scalar functions that preserve the nondegenerate eigenprojectors do not change this frame test. Both ordinary real three-dimensional carriers have simple spectra for X and iY .

On the first real carrier the X eigenvalues are $-5, 3, 30$ and the iY eigenvalues are $-\sqrt{1452}, 0, \sqrt{1452}$. Its squared overlap matrix, in increasing spectral order, has rows approximately $(0.422078, 0.155844, 0.422078)$, $(0.196970, 0.606061, 0.196970)$, and $(0.380952, 0.238095, 0.380952)$. On the second real carrier X has eigenvalues $-5, -2, 3$ and iY has eigenvalues $-\sqrt{2732}, 0, \sqrt{2732}$. Its rows are approximately $(0.00512445, 0.98975110, 0.00512445)$, $(0.49897511, 0.00204978, 0.49897511)$, and $(0.49590044, 0.00819912, 0.49590044)$.

The equality of the outer probability columns is structural. In a real carrier X has a real eigenbasis, while a real skew Y has conjugate eigenvectors for the opposite eigenvalues of iY . Their squared overlaps with any real X eigenvector are equal. The numerical column differences are below $6\text{e-}16$. Row and column permutations retain the identical-column obstruction. Two identical probability columns cannot each concentrate near one in different rows, so neither frame has the CKM hierarchy. This statement does not require interpreting a convention-dependent eigenvector phase as the CKM phase.

The other two three-dimensional carriers are a conjugate complex pair, so the preceding real-carrier argument must not be applied to them individually. Both have X spectrum $-2, -2, 3$. The repeated eigenvalue leaves a $U(2)$ freedom and prevents X from selecting a unique family frame. The unique X eigendirection has overlaps approximately $0.0381554, 0.5982882, 0.3635564$ with the iY eigenlines, in reverse order on the conjugate carrier. These overlaps are unchanged by rotations in the repeated eigenspace. Their maximum

is below 0.599, so even that fixed direction cannot be a nearly unmixed CKM row under any permutation. The receipt omits a full mixing matrix for these degenerate cases because it would depend on arbitrary eigenbasis choices.

Thus all four available three-dimensional blocks fail for this canonical pair. This does not exclude pairs built from additional unsigned channels, higher words, decorated C5 incidence, or another vacuum-selected Hermitian operator.

The twelvefold cover supplies a phase carrier

The primary ATLAS data gives M22 Schur multiplier of order twelve and lists its twelvefold cover. The recovered Formation work already tracks separate 2-primary and 3-primary cover restrictions. The full universal cover has central C12. An order-five element of M22 can be lifted to an order-five element of the cover: its fifth power lies in C12, and the fifth-power map is invertible on C12, so a central correction cancels that power. This lift commutes with the center. Their subgroup is C12 times C5, hence C60.

This is a group-theoretic inference from the cover data. It avoids the ordinary-M22 restriction that a single group element has no order sixty. It is not a claim that the old untwisted 7,392-state permutation module sees the central character; that module factors through M22 and its center acts trivially.

A character of this C60 subgroup has phase residue $n=5a+12b$ modulo 60, with a modulo 12 and b modulo 5. All sixty residues occur. The primitive positive phases are 6,42,66,78,102,114,138,174 degrees. The residue eleven, corresponding to 66 degrees, can be written with $a=7$ and $b=3$. These are possible character labels, not dynamically selected quantum numbers.

If one retains the previously nominated readout $C=1-2 \cos(12 \theta)$, the primitive residues n congruent to plus or minus one modulo five give $C=(3-\sqrt{5})/2=\phi^{-2}$. Its exact polynomial is C^2-3C+1 . The positive angles on this branch are 6,66,114,174 degrees. Thus the cover supplies both the relevant cyclotomic arithmetic and several competing branches. It neither selects 66 degrees nor derives the equality of this singlet readout with the observable CKM coefficient. The other primitive residues give the larger golden root.

Central phases do not select a residual eigenframe. Multiplying a residual group matrix by a central scalar changes its eigenvalues but leaves all its spectral projectors unchanged. Such scalar factors therefore cancel from projector overlaps. Similarly an overall Yukawa or Yukawa-column phase cancels from YY-dagger. A primitive eigenphase of 66 degrees is not evidence for a physical CKM phase of 66 degrees. Multiple charge-compatible interfering contractions and a nonzero rephasing invariant must be demonstrated explicitly.

The next interaction calculation

The useful continuation is a decorated order-five incidence construction coupled to the recovered signed C7 channels. First derive the allowed cover characters and local contractions from the actual Schur restrictions. A faithful central character cannot simply be attached to an ordinary A5 triplet without checking its extension and charge constraints. Build the resulting twisted or decorated carrier and identify two Hermitian operators with simple three-dimensional eigenframes. Their coefficients must follow from the permitted contractions or a solved vacuum, rather than from a fitted CKM matrix.

Then compute the relative spectral-projector overlaps and a CP-sensitive closed path invariant. Check whether any central phase survives these neutral observables. Fit only the existing two anchor magnitudes, if the construction admits two independently justified scales; evaluate the remaining CKM magnitude, effective golden coefficient, and physical phase as predictions. Enumerate every allowed orientation and character branch, including CP conjugates and the four golden-angle candidates. Finally enumerate symmetry-allowed counterterms to test whether the frame survives coefficient changes.

The recovered algebra has enough structure to make this calculation concrete. What it currently establishes is an exact operator reconstruction, a failure of the simplest canonical frame, and an order-60 cover route with an explicit branch and rephasing obstruction. A symmetry-protected quark frame has not been derived.

Reproduction and sources

Run `python python/develop_m22_frame.py` from the repository root with NumPy, SciPy, SymPy and python-flint installed. The output operator matrices and transpose/subdegree data are in `receipts/m22_frame/recovered_operators.json`. The diagnostic, exact polynomial factors and phase enumeration are in `frame_audit.json`. The trusted original object-array archive is the same one used by the byte-preserved Forge verifier; do not replace it with an untrusted NumPy pickle.

Recovered source theorems include `RAMANUJAN_WILSON_FAMILY.md`, `WEDDERBURN_35_RAMANUJAN_REDUCTION.md`, `RAW_FIVE_BLOCK_STAR_TRANSPORT_v11.md`, `CHAR0_TRANSPORT_GLUE_M5_v13.md`, `M22_PRIMARY_SCHUR_SIGNATURES_v11.md` and `L34_MATCHED_SELECTOR_V24.md` under `galois_merge/vendor/forge/THEOREMS`. The primary external cover reference is the ATLAS M22 page, <https://brauer.maths.qmul.ac.uk/Atlas/spor/M22/>. Its multiplier, cover and conjugacy-class data were checked on 5 October 2026. The CKM target and the mediator matching remain those of the existing protected-flavor chapters.

Triplet transport and exact residual obstructions

Research manuscript and continuation of the protected-flavor monograph. Prepared 5 October 2026. Status: computationally certified mathematical results with proofs; publication novelty and the proposed particle-physics interpretation require independent review. No golden CKM coefficient, CKM phase, or experimental mixing anchor enters the group, connection, or overlap enumeration.

Abstract and the claim being tested

We reconstruct a canonical decorated connection on a 7,392-cap M22/A5 geometry recovered from the Wilson work. Its order-five scalar adjacency B satisfies $B^2 = X + 6I$ in the preserved 152-dimensional commutant, so it cannot supply eigenframes independent of X . The cap graph consists of 176 distance-regular components with 42 vertices, with intersection array $\{6,5,1;1,1,6\}$. Each cap has five A4 subgroups. Matching them across a common D5 by their common involutions defines a parameter-free five-letter edge permutation. The resulting holonomy is S5. On the exterior square of the standard four-dimensional hyperplane representation, every edge exchanges the two local golden A5 triplets. There is no parallel rank-three projector on the base graph. Its connected 84-vertex orientation cover has A5 holonomy and admits exactly two such projectors; a positive interaction selects them. Independently, an exact census of the genuine complex triplet of 3.A6 gives 151 complete Abelian residual frames and 111 squared entry values. Every nonzero entry has magnitude at least 0.0759431758699. Two involution residuals also have an exact fixed-entry obstruction. These results identify concrete failures and a precise alignment repair, rather than deriving the nominated golden CKM relation.

The central question is whether the recovered symmetry can enforce $C_{\text{eff}} = |\text{Vub}|(1-|\text{Vub}|^2)/(|\text{Vus}||\text{Vcb}|) = \phi^{(-2)}$ and a physical CKM phase of 66 degrees. The results below do not establish either equality. In particular, geometric transport, a representation's character field, a central scalar phase, and the physical CKM invariant are different objects.

Exact scalar identity and the 42-cap component

$$B^2 = X + 6I, \quad B^3 = 4X + 11B + 6D$$

The trusted source is `galois_merge/vendor/forgedATA/RELATION/TASK_SELECTED_WORD_PROGRAMS_V23/exact_prefix_1023.npz`, SHA256

1351e514fcb5f23832f65665e0a12d2e172f8131c34875de0e03cddc5aec3093. It records the original integral Hecke coordinates of 1, X, Y and their binary words. The original X is orbital A49, and the original Y is -A12-A48-A59+A95+A116+A125. The sole valency-six orbital is B = A134; the sole valency-five orbital is D = A139. Solving against 152 independent preserved words and checking all 511 available extensions reconstructs multiplication by B and D without CKM input.

Theorem 1. In the full reconstructed regular commutant representation, $B^2 = X + 6I$ and $B^3 = 4X + 11B + 6D$. Consequently $B^4 - 4B^3 - 17B^2 + 24B + 36I = 0$. Its regular characteristic polynomial is $(t-6)^9(t-2)^{65}(t+1)^{27}(t+3)^{51}$. These are exact integral matrix identities, not rounded

spectral matches.

Proof. Express the orbitals B and D in the selected exact word basis using rational matrix inversion. Their recovered multiplication matrices have integral entries. Direct integer matrix products give the stated identities. The characteristic polynomial is computed independently from the integral matrix. The complete matrices and replay script are supplied with the receipts. The four distinct roots occur, so the displayed quartic is the minimal polynomial.

An independent degree-22 permutation construction uses the ATLAS standard M22 generators. A subgroup generated by an involution and an element of order three with product of order five has order 60. The selected cap class has subdegree counts $1^1, 5^1, 6^1, 10^1, 15^2, 20^4, 30^{42}, 60^{100}$, matching the recovered action. Enumerating its cosets accounts for all 443,520 group elements and 7,392 caps. Each cap contains six C5 subgroups; a C5 lies in exactly two caps in this class. The corresponding undirected cap graph has valency six. The implementation finds a 42-vertex component with distance layers 1,6,30,5 and the stated intersection array; transitivity gives 176 components.

The component graph is a known Hoffman-Singleton second-subconstituent graph. Its spectrum is $6^1, 2^{21}, (-1)^6, (-3)^{14}$. Neither the graph nor its intersection array is claimed as a new discovery. The five distance-three neighbors together with a vertex form an antipodal six-cap block; the component has seven such blocks. The exact component stabilizer acts faithfully as all 2,520 even permutations of the seven blocks, identifying it as A7. A six-cap block stabilizer acts faithfully as all 360 even permutations of its six caps, identifying an ordinary A6 containing the cap A5 with index six.

The identity $B^2 = X + 6I$ means every X eigenspace is B-invariant and conversely B selects only branches already lying in these spaces. Since the recovered X eigenvalues correspond to four distinct B^2 values, B and X have the same eigenspace decomposition. The order-five scalar operator therefore does not repair the earlier failure to obtain a new three-family mixing frame.

A canonical decorated transport

$$\Omega = \frac{EG_2}{\sqrt{5}}, \quad \Omega^2 = I, \quad \Omega T = -T\Omega$$

Each A5 cap has five A4 subgroups, equivalently the normalizers of its five Klein four subgroups. Adjacent caps intersect in D5. Each local A4 meets this D5 in a single order-two subgroup, and the five A4 subgroups exhaust its five involutions. Match the source and target A4 subgroups with the same actual involution. This defines a unique bijection of the five local letters, hence a permutation edge transport. The construction is covariant under M22 conjugation and has no continuous coupling or chosen CKM angle.

Theorem 2. For this incidence-matching connection, every edge permutation is odd and the holonomy of a 42-cap component is S5. On $W = \text{exterior-square } R$, where R is the four-dimensional standard hyperplane representation of the local five letters, transport exchanges the self-dual and anti-self-dual three-dimensional subspaces.

Proof. The supplied cap permutations identify all local subgroups and all 252 directed edge transports by exact set intersections. Every edge is odd. Products around loops, after a spanning-tree trivialization, generate all 120 permutations of five letters. An independently written verifier recomputes every transport from conjugated A4 and cap intersections, uses a depth-first rather than breadth-first tree, and again obtains the entire S5. One explicit five-cycle is cap 0, cap 1083, cap 2333, cap 3074, cap 5702, cap 0. Its loop permutation is (2,0,4,3,1) in zero-based image notation, an odd permutation. Vertex labels refer to the supplied deterministic

coset enumeration.

For the representation-theoretic statement, use the hyperplane basis $e_i - e_5$, $i=1,\dots,4$, with metric $G = I + 11^T$, of determinant five. Let G_2 be its induced exterior-square metric and E the matrix of wedge pairing in the six ordered bivector coordinates. The Hodge operator is $\Omega = E G_2 / \sqrt{5}$, and $\Omega^2 = I$. Its two projectors $(I \pm \Omega)/2$ have rank three. For any five-letter permutation p , its induced bivector transport T satisfies $\Omega T = \text{sign}(p) T \Omega$. Every edge is odd, hence maps one Hodge branch to the other. The local A_5 preserves the branches. Their order-five characters are $(1+\sqrt{5})/2$ and $(1-\sqrt{5})/2$ in opposite classes, identifying the familiar two real golden triplets. This is a real representation-theoretic branch exchange, not a CP phase calculation.

For an independent irreducibility check, the bivector character is $\chi(p) = ((\text{fix}(p)-1)^2 - (\text{fix}(p^2)-1))/2$. Its squared character sum is 120 over S_5 , giving inner product one, and is 120 over A_5 , giving inner product two. The full S_5 commutant is therefore scalar, while the A_5 commutant is spanned by I and Ω . Exact commutator-rank calculations reproduce dimensions one and two.

Corollary. There is no parallel Hermitian rank-three projector on this connected base connection. Any such projector would commute with every loop at the root. The irreducible S_5 action forces it to be a scalar idempotent, hence either zero or identity, of rank zero or six. The theorem excludes parallel projectors for the specified connection; it does not prohibit homogeneous A_5 triplet bundles or all possible M_{22} models.

What edge symmetry actually enforces

$$\dim \text{Hom}_{D_5}(3, 3^t) = 1, \quad \dim \text{Hom}_{D_5}(3, 3^{\bar{t}}) = 2$$

The D_5 intersection reveals an additional coefficient issue. The restriction of one golden triplet to D_5 is a one-dimensional sign axis plus a two-dimensional plane carrying C_5 weights ± 1 ; its Galois partner has the same axis and plane weights ± 2 . Transport between adjacent caps exchanges the two five-cycle character classes. Taking the exact character inner products over the ten common elements gives dimension one for a same-branch edge intertwiner and dimension two for an opposite-branch edge intertwiner.

Thus a same-branch covariant map has rank at most one. An opposite-branch map can be invertible, but its axis and plane couplings are independent. The particular incidence matching above picks a unitary map on both channels; D_5 symmetry alone does not force their relative strengths. It would be incorrect to turn this particular geometric isometry into a general coupling-protection theorem. A particle interaction must explain why it uses the incidence matching rather than the full symmetry-allowed two-channel family.

Multiplying edges by central scalar phases, including phases from a central cover, leaves $\Omega T = -T \Omega$ unchanged. A phase factor also cancels in $T P T^{-1}$, so it cannot repair the parallel-projector obstruction. A noncentral transport or different fiber content could do so; neither is excluded here. No identification of a cover eigenphase with the CKM phase is made.

An explicit protected alignment interaction

$$J_{V,s} = (-1)^s \Omega_V, \quad P_{\pm} = \frac{I \pm J}{2}$$

The orientation cover has vertices (v,s) , with $s=0,1$, and lifts an edge $v-w$ to $(v,s)-(w,1-s)$. It is connected because the base has odd loops. Each component has 84 vertices. The lifted

holonomy consists exactly of the 60 even permutations, A_5 . The field $J_{(v,s)} = (-1)^s \Omega_v$ is parallel because edge transport anticommutes with Ω . The deck exchange changes the sign of J .

Theorem 3. On this cover the only parallel Hermitian rank-three projectors are $P_+ = (I+J)/2$ and $P_- = (I-J)/2$. For dimensionless Hermitian endomorphism order parameters P_a , an explicit potential selecting exactly these configurations is the sum of three nonnegative terms: a positive coefficient λ times the sum of squared norms of $P_a^2 - P_a$, a positive coefficient μ times the sum of $(\text{Tr } P_a - 3)^2$, and a positive coefficient κ times the sum over edges of squared norms of $P_b - T_{ba} P_a T_{ba}^{-1}$. The norms use the positive fiber metric.

Proof. Zero energy enforces a rank-three projector and parallel transport. At a root the A_5 commutant is $\text{span}\{I, J\}$. A Hermitian idempotent in this commutant has eigenvalues $a+b$ and $a-b$ on the two inequivalent triplets. Rank three forces these to be 1 and 0 in either order. This gives exactly P_+ and P_- . Conversely, their parallelism and rank make all terms vanish. On the base the identical constraints have no zero-energy configuration, by the scalar S_5 commutant. The cover is therefore a concrete repair of this alignment problem, rather than a choice of a fitted frame.

The selected projectors remain exact for arbitrary positive λ , μ and κ within this stated interaction. This is a classical robustness result for the declared potential family. Symmetry allows additional interactions; no general protection against counterterms or radiative corrections is claimed.

This establishes a local rank-three subspace. It does not by itself yield exactly three light global quark modes, nondegenerate quark masses, a relative up/down frame, a UV origin for the interaction, or a cosmological selection of its two branches. In particular, the two real Hodge sectors are not the positive and negative physical CP vacua. Additional physical fields and a light-mode analysis are necessary before treating the construction as a flavor model.

Exact complex-triplet residual gap

$$|V_{ij}| \in \{0\} \cup [0.0759431758\dots, 0.9566183482\dots] \cup \{1\}$$

The identified A_6 block supplies a concrete place to investigate a genuine complex triplet. Ordinary A_6 has no faithful ordinary triplet; its triple cover $3.A_6$, the Valentiner group $\text{Sigma}(360 \times 3)$, does. We use the published generators of Hagedorn, Meroni and Vitale, equations (7) and (33), and close them exactly in $Q(\sqrt{5}, \omega)$, with $\omega^2 + \omega + 1 = 0$. The closure has 1,080 elements and a center of order three, and its real A_5 subgroup has order 60. A consistent embedding of this triple cover into a particular central M_{22} cover is a separate requirement, not an assumption silently imposed by this calculation.

The complete minimal Abelian residual frames comprise 40 order-three frames, 45 order-four frames, 36 order-five frames, and 30 joint Klein frames, totaling 151. The order-three frame count is smaller than the number of cyclic order-three subgroups because central multiplication gives the same projectors. Exact spectral-projector enumeration checks coverage of all 474 noncentral elements of orders three, four and five. Order-twelve and order-fifteen lifts have the same frames as their cubes; order-two and order-six residuals leave a two-dimensional degeneracy. Higher Abelian residuals add the center or already contain one of the complete frames. This classification agrees with the published subgroup analysis; it is not presented as a newly discovered flavor group.

Theorem 4. Every entry probability between the 151 complete frames belongs to the supplied exact set of 111 values. The smallest positive value is $p_{\min} = 0.005767365961210174\dots$, and the largest value strictly below one is $p_{\max} = 0.9151186641666433\dots$. Consequently every nonzero magnitude is at least $0.07594317586992\dots$, and every subunit magnitude is at most $0.956618348228092\dots$. Rephasing and all row and column mass permutations preserve these bounds.

Proof. Construct cyclic projectors as the finite Fourier sums of exact group powers, retaining the three projectors of trace one. Construct Klein projectors by their four characters, retaining the three projectors of trace one. For all 11,476 unordered frame pairs with repetition, evaluate all nine traces $\text{Tr}(P_i Q_j)$, giving 103,284 exact comparisons. Computation is in $\mathbb{Q}(\zeta_{60})$, reduced modulo Φ_{60} . All probabilities are exactly fixed by complex conjugation. Convert them to degree-seven rational polynomials in $t = \zeta_{60} + \zeta_{60}^{-1}$. Its minimal polynomial is $t^8 - 7t^6 + 14t^4 - 8t^2 + 1$, and the physical embedding is its largest real root, $2 \cos(\pi/30)$. Rational root isolation and interval Horner evaluation give strictly ordered, disjoint rational enclosures for all 111 values. This proves both extrema and their ordering without using floating point to choose the winner. The 60th root is a common arithmetic field for orders 3,4,5; its appearance is not a prediction of 66 degrees.

Explicitly, $p_{\min} = -t^{7/10} - t^{6/20} + 11t^{5/20} + 3t^{4/10} - 3t^{3/5} - 9t^2/20 - t/5 + 1/2$. The upper endpoint below one is $p_{\max} = 2t^{6/9} - 4t^{4/3} + 2t^2 + 1/9$. The full rational polynomial table and certified intervals are in `receipts/m22_interactions/valentiner_exact_gap.json`. A separate unitary numerical census of all 22,801 ordered frame pairs independently reproduces the extrema; 36 mass label permutations per pair correspond to 820,836 labeled pairs, and no labeling changes the entry spectrum.

The nominated values $|V_{cb}|=0.0411$ and $|V_{ub}|=0.00352144218276$ fall inside the forbidden nonzero gap. Thus complete Abelian residual symmetries cannot give that CKM hierarchy in this triplet without symmetry breaking. These numbers are used only after the exact spectrum is constructed, as a comparison to the existing nominated model, not as a current experimental global fit. The result excludes the specified exact residual construction; it does not exclude 3.A6 as a broken flavor symmetry.

Partial residuals and the fixed-line obstruction

$$|V_{ij}|^2 \in \left\{ 0, \frac{3-\sqrt{5}}{8}, \frac{1}{4}, \frac{1}{2}, \frac{3+\sqrt{5}}{8}, 1 \right\}$$

Leaving a two-dimensional degenerate sector in each quark mass matrix permits continuous rotations, so the preceding complete-frame gap does not apply directly. Nevertheless, retaining an involution in both sectors fixes one line in each. There are 45 order-two elements. Their rank-one projectors are $(I+g)/2$. Evaluating all 2,025 ordered projector overlaps exactly gives squared fixed entries $0, (3-\sqrt{5})/8, 1/4, 1/2, (3+\sqrt{5})/8$, and 1 . Their magnitudes are $0, 0.309016994\dots, 0.5, 0.707106781\dots, 0.809016994\dots$, and 1 .

Theorem 5. If both nondegenerate quark Hermitian mass matrices commute with nontrivial involutions in the stated triplet, at least one CKM entry is in this fixed set, irrespective of the two otherwise arbitrary $U(2)$ rotations and irrespective of any added compatible CP constraint. Distinct fixed lines cannot give the nominated hierarchical CKM matrix. Identical fixed lines give an exactly decoupled family and hence a two-family mixing block. A zero fixed overlap also supplies an exact zero entry. All these cases require symmetry breaking to realize the nominated matrix with nine nonzero entries.

Proof. Commutation with an involution preserves its one-dimensional $+1$ eigenspace and its two-dimensional -1 eigenspace. Nondegenerate mass eigenstates therefore include the fixed line in each sector. Their overlap is a CKM entry. Rotations confined to the two-dimensional complements do not change this overlap. The exact projector calculation gives the stated list. The nominated matrix's off-diagonal magnitudes are below 0.3090 and its diagonal magnitudes exceed 0.8091; none is zero or one. Adding CP cannot change the representation-fixed line overlap. This argument does not assume a specific CP automorphism or a chosen mixing phase.

Publication scope and the next physical problem

The result suitable for scrutiny is a conjunction of exact transport obstruction, explicit cover alignment mechanism, and exact quark residual gaps. It is not a derivation of the golden CKM relation. Known ingredients include the graph, the two A5 triplets, the Valentiner generators and residual-subgroup classification, and the Hodge transformation law. The documented contribution is their explicit reconstruction on the recovered cap system, the canonical involution-matching connection with certified holonomy, the scoped interaction repair, and exact gap certificates applied to the nominated quark mechanism. A literature search did not establish novelty of this combination; independent specialist review remains necessary before submitting a novelty claim.

The remaining viable problem is now narrower. A proposed theory must specify a noncentral, charge-compatible interaction that produces three light quark modes, breaks the exact residual obstruction in a controlled way, and correlates the up/down breaking operators after canonical normalization. It must then derive the coefficient $\phi^{(-2)}$ and a neutral CP observable corresponding to 66 degrees without inserting either target. Central phases alone, scalar order-five adjacency alone, and exact complete or two-involution residual frames cannot supply that result. Small free corrections that merely fit the missing entries would not resolve the original coupling-protection problem.

Reproduction and sources

Install `python/m22_frame_requirements.txt`. From the repository root run `python python/certify_m22_c5.py`, `python python/develop_m22_transport.py`, `python python/verify_m22_transport.py`, `python python/develop_valentiner_frames.py`, and `python python/certify_valentiner_gap.py`. The first three scripts distinguish the preserved Wilson regular algebra from the independently enumerated cap geometry. The independent verifier reconstructs subgroup intersections directly and does not reuse the connection's cocycle code or centralizer solver. The exact residual-gap script checks its entire field and real embedding using rational intervals. Logs and JSON receipts are included. A cold replay compares JSON outputs byte for byte; the compressed permutation file is validated by its array contents, not by ZIP container metadata. No CKM input enters any of these five computations.

The standard degree-22 generators are from the ATLAS of Finite Group Representations, <https://brauer.maths.qmul.ac.uk/Atlas/spor/M22/>, generator files `gap/M22G1-p22B0.g1` and `gap/M22G1-p22B0.g2`. Thomas Connor and Dimitri Leemans, The subgroup lattice of M22 (2012), <https://leemans.dimitri.web.ulb.be/atlaslat/m22.pdf>, independently documents the relevant A5, A6 and A7 subgroup classes; the selected class is recorded by explicit permutations rather than inferred only from a class name. Claudia Hagedorn, Aurora Meroni and Lorenzo Vitale, Mixing Patterns from the Groups $\Sigma(n, \phi)$, arXiv:1307.5308, <https://arxiv.org/abs/1307.5308>, supplies the Valentiner generators and prior residual analysis. The known 42-vertex graph is described in Wei Jin and Cheryl E. Praeger, Normal quotients of

diameter at most two of finite three-geodesic-transitive graphs, *Journal of Combinatorial Theory A* 178 (2021), 105349, <https://doi.org/10.1016/j.jcta.2020.105349>. These sources establish antecedents; they do not independently verify the new receipts or confer a publication claim.

Valentiner interactions and the protection boundary

Result and scope

The recovered cap geometry has a supported central-cover route to a genuine complex triplet. The triplet's first new symmetric invariant is a unique sextic, derived here directly from its exact generators. A minimal vectorlike mediator model has exactly three light modes per charge sector before electroweak breaking, and its canonical matching transmits the flavon frame. Its complete scalar operator space nevertheless contains independent cross-sector quartics. A target-free numerical local-vacuum experiment shows one such operator changing the observable CKM coefficient. Neither the golden coefficient nor the 66-degree physical phase is derived.

These results replace an unspecified symmetry search with a concrete representation, a fixed invariant tensor, specified field content and a counterterm audit. The sextic itself is the classical Valentiner/Wiman invariant, not a newly discovered polynomial. Publication novelty of the combined construction and obstruction requires comparison with the existing invariant-theory and flavor literature.

The central-cover connection

The earlier cap computation established a component stabilizer A_7 and a six-cap block stabilizer A_6 of index seven in that A_7 . Both A_7 conjugacy classes in M_{22} lift to $3.A_7$ in $3.M_{22}$. The GAP Character Table Library lists the index-seven maximal subgroup of $3.A_7$ as $3.A_6$, of order 1080. Thus the full preimage of the cap-block A_6 is the nonsplit Valentiner cover, rather than the split product $C_3 \times A_6$. This establishes the abstract central-cover chain $3.A_6 < 3.A_7 < 3.M_{22}$ for the recovered stabilizer geometry.

This conclusion uses the independently computed ordinary stabilizers together with published subgroup classifications. We have not constructed an explicit matrix representation of $3.M_{22}$ or lifted every recovered permutation generator. The local triplet does not extend to a three-dimensional representation of the whole cover by this argument. A global bundle, a global mass operator on 7392 caps, and decoupling of its other modes remain separate obligations.

Primary sources:

<https://www.math.rwth-aachen.de/~Thomas.Breuer/ctbllib/ctbltoc/data/3.M22.html> ;

<https://www.math.rwth-aachen.de/~Thomas.Breuer/ctbllib/ctbltoc/data/3.A7.html> ;

<https://www.math.rwth-aachen.de/homes/sam/ctbllib/ctbltoc/data/3.A6.html> . The exact local $SU(3)$ generators are those of Hagedorn, Meroni and Vitale, <https://arxiv.org/abs/1307.5308> , equations (7,33).

An exact tensor boundary

Let R be the faithful complex triplet and $G=3.A_6$. We closed its four generators exactly in $Q(\sqrt{5}, \omega)$, with ω squared plus ω plus one equal to zero, obtaining all 1080 elements. For every pair p, q with total degree at most eight, we evaluated the character average $\dim \text{Inv}(R \text{ tensor } p \text{ times } \bar{R} \text{ tensor } q) = \sum_g \text{tr}(R(g))^p \text{conjugate}(\text{tr}(R(g)))^q / 1080$. We independently computed the corresponding $SU(3)$ tensor multiplicities from partitions and hook lengths.

Theorem. At total degree at most five the G -invariant tensor space equals the $SU(3)$ -invariant tensor space, for arbitrarily labeled triplet fields. At degree six, the $(3,3)$ space still has dimension six for both groups. The $(6,0)$ space has dimension six for G and five for $SU(3)$, leaving exactly one new tensor. The conjugate statement holds for $(0,6)$; the other degree-six bidegrees add none.

Proof. $SU(3)$ invariants form a subspace of G invariants. The exact character averages and independent hook-length counts coincide in the stated bidegrees. Equality of dimensions therefore proves equality of spaces. In degree six, the symmetric-power average gives $\dim \text{Inv}(\text{Sym}^6 R) = 1$. Its tensor cannot be an $SU(3)$ symmetric invariant, because there is no positive-degree $SU(3)$ -invariant polynomial on one fundamental triplet. It exhausts the one-dimensional quotient. All character coefficients cancel exactly in the four-dimensional number field. The full counts are in `receipts/m22_interactions/valentiner_invariants.json`.

For one commuting triplet, through degree six the only nonconstant invariant bidegrees are $(1,1)$, $(2,2)$, $(3,3)$, $(6,0)$ and $(0,6)$, each with dimension one. The first three are powers of the norm. Consequently every renormalizable single-triplet potential has an accidental continuous $SU(3)$ orientation symmetry. A sextic is the first possible finite-group anisotropy. This is an effective interaction in four spacetime dimensions, requiring a suppression scale or a specified mediator completion.

The unique sextic in the recovered basis

$$F_6 = x^6 + y^6 + z^6 + A(x^4 y^2 + y^4 z^2 + z^4 x^2) + B(x^2 y^4 + y^2 z^4 + z^2 x^4) + D x^2 y^2 z^2$$

Writing the triplet as (x, y, z) , normalize the sextic by its x^6 coefficient. It is $F_6 = x^6 + y^6 + z^6 + A(x^4 y^2 + y^4 z^2 + z^4 x^2) + B(x^2 y^4 + y^2 z^4 + z^2 x^4) + D x^2 y^2 z^2$, with $A = [15 - 9\sqrt{5} + i(15\sqrt{3} + 3\sqrt{15})]/8$, $B = [15 + 9\sqrt{5} + i(-15\sqrt{3} + 3\sqrt{15})]/8$ and $D = 15 - 3i\sqrt{15}$.

The coefficients arise from solving invariant linear equations, without a selected angle or golden CKM coefficient. Sparse monomial generators first reduce the degree-six space; the remaining generator leaves a single invariant. Each of the four polynomial identities $F_6(g \phi) = F_6(\phi)$ is then checked exactly in the number field. The receipt records every monomial coefficient.

A potential contains a free complex overall coefficient multiplying F_6 plus its conjugate. Uniqueness fixes internal component ratios; it does not fix the coefficient relative to other invariants. Its complex coefficients also do not establish physical CP violation. A consistent generalized CP action on all fields and its allowed couplings must still be constructed. Ordinary componentwise conjugation is not assumed to be a symmetry in this basis.

Specified quarks, flavons and mediator interactions

Take Q_L in R , the Higgs in the trivial family representation, three right-handed up quarks and three right-handed down quarks in family singlets, and one vectorlike quark triplet F_u and one F_d in R . These vectorlike fields are Standard Model weak singlets with the respective up or down electric charge. Introduce six Standard Model singlet flavons ϕ_i in R . This is a local four-dimensional effective model after selecting the cap stabilizer; it is not a derived global cap spectrum.

Add a specified shaping group C_6 to the sixth power. Each right-handed singlet f_R has charge +1 under its own factor, its flavon has charge -1, and all other listed fields are neutral under that factor. The family and shaping symmetries permit, per sector, $M_f F_{\bar{L}} F_R$, $h_f Q_{\bar{L}} H_f F_R$, and $\sum_i \lambda_i F_{\bar{L}} \phi_i f_R$, plus Hermitian conjugates. Here H_u is the conjugate Higgs doublet and H_d the ordinary one. Each family contraction is unique, by $\dim \text{Hom}_G(R, R) = 1$. The mediator mass is therefore flavor-universal. Cross-column flavon couplings are forbidden by the specified shaping charges. A bare $Q_{\bar{L}} H_f f_R$ coupling has no family singlet. A mediator-triplet bilinear with one triplet flavon is forbidden by the nontrivial central charge. These exhaust renormalizable mass and Yukawa structures for this listed field content.

The symmetry allows separate M_f , h_f and three λ_i in each sector. It does not correlate the up and down parameters. The shaping group is an explicit model assumption, not something derived from M_{22} . Its gauging, anomalies, a scalar messenger completion and quantum thresholds are not claimed.

Three light modes and canonical projectors

$$K_f = I + A_f^\dagger A_f, \quad Y_f = -h_f A_f K_f^{-1/2}, \quad |V_{ij}|^2 = \text{Tr}(P_i^u P_j^d)$$

At $H=0$ define C_f with columns λ_i times the flavon expectation values. The heavy row block is $[C_f, M_f I]$, with three left-handed rows and six right-handed columns. For nonzero M_f it has rank exactly three, regardless of C_f . Its right kernel therefore has dimension three; the three Q_L components are the corresponding left chiral light modes. The three heavy singular values are $\sqrt{|M_f|^2 + \sum_i \lambda_i^2}$. There are exactly three light families per charge sector in this specified local model, without interpreting a rank-three fiber as a global family count.

Set $A_f = C_f / M_f$, $K_f = I + A_f^\dagger A_f$. The normalized light right frame is $[I; -A_f] K_f$ to the power $-1/2$, and the matched Yukawa matrix is $Y_f = -h_f A_f K_f$ to the power $-1/2$. These are exact tree identities at $H=0$, with no small-flavon expansion. They imply $H_f = Y_f Y_f^\dagger = |h_f|^2 S_f (|M_f|^2 I + S_f)$ inverse, where $S_f = C_f C_f^\dagger$.

The positive spectral function is strictly increasing. Thus the left spectral projectors of H_f and S_f agree, including the mass ordering for a nondegenerate spectrum. Quark mixing is computed as $|V_{ij}|^2 = \text{Tr}(P_i^u P_j^d)$, and $J = \text{Im Tr}(P_1^u P_1^d P_2^u P_2^d)$. Arbitrary flavon frames remain arbitrary physical quark frames; canonical normalization neither creates nor protects the golden relation. At finite Higgs expectation value, heavy-light mixing supplies higher-dimensional EFT corrections and potential charged-current nonunitarity. Those corrections are not included in this matching statement.

The complete scalar space through degree six

For the six specified flavons and C6 shaping factors, a degree smaller than six has equal numbers of each flavon and its conjugate. Consequently the scalar operators coincide with balanced Gram contractions: six quadratic norms; 36 quartics consisting of 21 norm products and 15 absolute overlap squares. With one Higgs, add its quadratic norm, its quartic norm and six Higgs-norm/flavon-norm products, for 50 real renormalizable scalar couplings in total.

At degree six the general Hermitian flavon space has 198 real parameters: 56 cubic norm products, 90 norm-times-overlap squares, the real and imaginary parts of 20 Gram triangles, and the real and imaginary parts of six individual $F6(\phi_i)$. Balanced epsilon contractions are Gram determinants and add no independent operators. Gram rank relations first occur at degree eight. The shaping charges forbid mixed polarizations of the new sextic at degree six: an unbalanced field must occur six times by itself. Under a specified CP action the count can decrease, but no CP classification has been assumed here. Every absolute overlap square remains CP-even under a common unitary anti-linear CP action on the triplets.

In particular, $\epsilon |\phi_{u1}^\dagger \phi_{d1}|^2$ squared is an independent allowed quartic. Neither the group nor the shaping charges constrain its coefficient to the chosen sextic coefficient. This is a concrete obstruction to claiming protected alignment merely from uniqueness of F6. It is the interaction-level counterpart of the earlier allowed linear angular deformation, although no equality between those two coordinate descriptions is assumed.

A target-free local-vacuum deformation experiment

We minimized the phase of each sextic analytically, fixed each flavon norm to one, and considered the reduced potential $\sum_i -|F6(\phi_i)| + \epsilon |\phi_{u1}^\dagger \phi_{d1}|^2$ squared. This is the fixed-norm reduction of polynomial sextic terms plus their conjugates, with independent phases minimized. Radial stability requires separate radial terms; no global minimum or cosmological vacuum selection is certified.

Six independently generated sextic stationary directions were selected from the numerical group orbit, and their column strengths set to 0.07, 0.31 and 0.9 in each sector. None is a measured CKM anchor or the golden target. Analytic gradients are checked against independent central finite differences, and each numerical branch must have gradient norm below $1e-7$. The complete matrices, CP invariant and effective golden coefficient are recorded in `receipts/m22_interactions/valentiner_alignment_response.json`. For $\epsilon=0$, C_{eff} is 0.4830422217 and J is 0.01663167860. At $\epsilon=0.01$ they become 0.4831824152 and 0.01663578380. The intermediate positive and negative 0.001 cases also give distinct values. The $\epsilon=0$ and $\epsilon=0.01$ branches have different C_{eff} ; this change survives the exact canonical mediator matching theorem. The mixing is not phenomenologically realistic, which is appropriate for an allowed-deformation counterexample rather than a fitted proposal.

The found single-flavon extrema form a numerical 60-line orbit, and sixteen independent searches reach modulus approximately 1.0886621079. These are numerical observations, not an exact orbit classification or a proof of the global modulus bound. Optimizer success flags and final gradients are retained, including precision-loss flags when a sufficiently small verified gradient remains.

What the next derivation must add

The sextic supplies a real representation-derived coefficient pattern, and the mediator calculation supplies a concrete light-family and normalization result. Their combination does not protect the observable golden relation. The next justified search must introduce interactions that correlate the sectors while controlling all 15 independent quartic overlaps, permit useful mixed sextic contractions without opening arbitrary singlet-copy matrices, and define a consistent generalized CP action. A proposed mechanism must then demonstrate its chosen spectrum and vacuum, test all remaining free parameters, and match the observable projectors. No choice of the desired CKM angle enters the invariant derivation above.

Reproduction

Run `python python/develop_valentiner_invariants.py`, then `python python/develop_valentiner_alignment.py` from the repository. The first command produces an exact invariant receipt and 24 canonical-matching scenarios. The second produces a numerical local-branch receipt with an independent gradient check. The exact group code and dependencies are shared with `python/develop_valentiner_frames.py` and `python/m22_frame_requirements.txt`. The monograph renderer includes this complete chapter. No new Lean theorem or experimental measurement is asserted.

Product-family CP and the global spectrum

The subsequent [input audit](FLAVOR_SEARCH_INPUT_AUDIT.md) distinguishes these separate constructions and checks their missing matching steps. In particular, the supersymmetric messenger superpotential is not the supplied positive scalar alignment potential, and the nonsupersymmetric quark example's conjugate-link reverse coupling cannot be copied unchanged into a superpotential with just one chiral link. The audit also retains updated experimental comparisons without changing the historical inputs.

Concrete advance and remaining boundary

This push constrains the problematic cross-sector interaction by changing the specified family group to $G_u \times G_d$, with each $G_f = 3.A6$, and joining the sectors with a bifundamental link. The bare absolute overlap quartic is forbidden. The first dressed overlap is degree six. The link has a unique new holomorphic sextic tensor, whose component ratios come from the previously derived Valentiner invariant. A renormalizable supersymmetric messenger chain generates this contraction at leading tree order, with its complete allowed holomorphic operator audit retained. A positive alignment interaction has exact zero-energy states at every group matrix, without inserting a CKM angle.

We also construct an exact anti-linear involution on the specified triplet and singlet fields. It leaves the sextic conjugate as required, and a canonical quark calculation produces a pair with opposite nonzero Jarlskog invariants. This establishes physical CP violation in a conditional matched example; it does not derive a unique phase or select the physical vacuum. Finally, the canonical cap connection is certified to have no zero modes. A fully specified modified global mass-squared operator has exactly three zero modes among 44,352 modes, with a gap of one half. Its required symmetry breaking and its chiral quark completion remain independent assumptions.

Exact CP on the elementary field content

$$X\overline{R(g)}X^{-1} = R(u(g)), \quad X\overline{X} = I$$

Use the published outer automorphism that sends the faithful triplet to its conjugate: a maps to a^2 , h to q , f to $f a^2 h f$, h to q , and q to a . We evaluate these words on the exact generators, propagate the map through all 1080 elements, and check that every repeated group word has the same image. We solve $X \text{conjugate}(R(g)) = R(u(g)) X$ as exact linear equations over $Q(\sqrt{5}, \sqrt{-3})$. The published representative can square to an inner action. An explicitly recorded inner adjustment yields a symmetric unitary X with $X \text{conjugate}(X) = I$.

The receipt gives the matrix entries, its normalization and the adjustment word. Before normalization, $X_0^\dagger X_0 = (6-2\sqrt{5}) I$. The sextic identity is checked coefficient by coefficient: $F_6(X_0 \phi) = (6-2\sqrt{5})^3 \text{conjugate}(F_6)(\phi)$. Equivalently $F_6(X \text{conjugate}(\phi)) = \text{conjugate}(F_6(\phi))$. Thus real coefficients of F_6 plus its conjugate define an invariant interaction on this specified field content. Ordinary componentwise conjugation in the original basis would not establish this result.

There is an important scope restriction. We enumerate all 17 conjugacy classes and the four outer-class representatives supplied by the known Klein outer automorphism group. Every representative fails to send at least one class to its inverse class. For the representative interchanging this triplet with its conjugate, two classes fail. Inner adjustments do not change these class permutations. There is therefore no class-inverting automorphism that could define the same physical CP operation for arbitrary irreducible field content. The anti-linear operation constructed here is restricted to the elementary triplet, singlet and bifundamental fields that pass the explicit consistency and Lagrangian checks. Adding a different elementary representation requires a fresh audit.

Primary sources: Hagedorn, Meroni and Vitale, <https://arxiv.org/abs/1307.5308> , equation (34), for the outer maps and their classification; Chen et al., <https://arxiv.org/abs/1402.0507> , for the class-inversion requirement in generic finite-group CP; <https://doi.org/10.1140/epjc/s10052-025-15050-8> , for the representation-content dependence of conjugation in Sigma(1080). Our exact class enumeration and intertwiner calculation are independent computations using the recovered generators.

Cross-sector sequestering and its exact operator boundary

$$G_u \times G_d, \quad |\phi_{ui}^\dagger \wedge \phi_{dj}|^2 \quad (\text{degree } 6)$$

Take ϕ_{ui} in (3,1), ϕ_{dj} in (1,3), and Λ in (3,3 $\bar$). Retain one independent C6 shaping factor for each right-handed singlet and its flavon. Before a link is inserted, $|\phi_{ui}^\dagger \phi_{dj}|^2$ squared is not an invariant of the product family group. There is no invariant identification of its independent family indices. Its proposed replacement $|\phi_{ui}^\dagger \Lambda \phi_{dj}|^2$ squared is invariant and has field degree six. Its nine coefficients remain independent. Arbitrary additional phase charges could not have forbidden the original absolute overlap square in the diagonal-group model: its charges cancel identically.

The bifundamental Molien calculation averages over the 1080 squared product group using exact power-trace class signatures and Newton identities. Through total degree six it has nonzero spaces (1,1) of dimension one, (3,0) and (0,3) of dimension one, (2,2) of dimension two, (4,1) and (1,4) of dimension one, (3,3) of dimension three, and (6,0) and (0,6) of dimension two. This is the complete self-operator count through that degree, not only a selected contraction.

The degree-three invariant is $\det \Lambda$. The two holomorphic degree-six invariants are its square and $I_6(\Lambda)$, obtained by contracting the unique symmetric sextic tensor with six link matrices and its conjugate tensor. Consequently there is only one genuinely new holomorphic orientation tensor at this degree. Balanced link-only sextics are the three symmetric polynomials of the singular values of Λ . At degree five the invariant is $\det \Lambda \times \text{Tr}(\Lambda \dagger \Lambda)$. Real parts are selected by the field-content CP involution; none of these counts supplies a relation between independent coupling constants.

The complete renormalizable scalar count for the listed six flavons, one link and one Higgs is 58 real parameters with this CP restriction. There are eight quadratic norms, one real cubic determinant coefficient, and 49 quartics. The quartics consist of 21 flavon norm products, six within-sector absolute overlaps, two link self-contractions, six link-norm/flavon-norm products, six $\phi \dagger \Lambda \Lambda \dagger \phi$ or conjugate-side contractions, and eight Higgs quartics and norm portals. The nine direct up/down overlap quartics of the earlier diagonal-group model are absent. Without CP the determinant coefficient has two real components, giving 59 parameters. The counts refer to this explicitly assumed product family and shaping content, not to every possible UV theory.

On a unitary link, $\Lambda^\dagger \Lambda$ is proportional to I . The six link/flavon orientation quartics then become norm terms and cannot rotate a source frame. Thus this construction removes the previous quartic angular perturbation in that regime. At degree six the dressed overlaps return, and a general Wilsonian effective theory permits them even if a chosen tree potential sets their coefficients to zero. They must be controlled by an additional mechanism before an exact CKM prediction could be protected.

A representation-derived positive link interaction

$$I_6(U) = \langle T, T \circ U \rangle, \quad \text{Re } I_6(U) \leq \|T\|^2 = 16$$

Let T be the symmetric coefficient tensor defined by $F_6(\phi) = T$ contracted with six copies of ϕ . Tensor norms require multinomial weights; the exact polynomial coefficients give $\text{norm}(T)^2 = 16$. Define $I_6(\Lambda) = \text{inner product of } T \text{ with } T \text{ composed with } \Lambda$. For unitary U , $\text{norm}(T \text{ composed with } U) = \text{norm}(T)$, so $\text{Re } I_6(U)$ is at most 16 by Cauchy-Schwarz. Equality is equivalent to $T \text{ composed with } U = T$. All 1080 exact group matrices satisfy that tensor identity, independently of the chosen quark model or CKM target.

A globally nonnegative polynomial alignment energy is $V_L = a \text{Tr}[(\Lambda^\dagger \Lambda - \text{ell } I)^2] + b |\det \Lambda - \text{ell } I|^2 + c \text{norm}[T \text{ composed with } \Lambda - \text{ell } I \text{ to the sixth } T]^2$, for positive a, b, c and ell . Every $\Lambda = \text{ell } R(g)$ is an exact zero-energy state. On the unitary, determinant-fixed manifold its last term is proportional to $16 - \text{Re } I_6(U)$. The last norm square contains degree-six and degree-twelve terms; the interaction is not a wholly renormalizable potential. Its coefficients and its higher-degree completion are explicit choices. We establish these zero-energy states and their positivity, without claiming an independently derived UV origin or a complete classification of all zero-energy matrices.

The link transforms under the elementary CP operation as Λ maps to X conjugate(Λ) $X^\dagger$. I_6 and $\det \Lambda$ then map to their conjugates. The positive interaction is therefore invariant. It correlates the sectors through a fixed representation tensor rather than through an arbitrary fitted matrix. Its degenerate group-related states do not select a physical CKM phase by themselves.

A renormalizable messenger origin for the sextic

The sextic also has an explicit interaction-level origin. Symmetric squared of the faithful triplet is an irreducible complex six, and symmetric cubed is an irreducible real ten. The latter has a unique invariant symmetric bilinear Q_{10} . Polarizing the sextic across two groups of three indices gives this bilinear, with $F_6(\phi) = \text{Sym}^3(\phi)$ transpose $Q_{10} \text{Sym}^3(\phi)$. In an orthonormal symmetric-cube basis its exact identity is $Q_{10}^\dagger Q_{10} = (8/5) I$. We certify all 100 entries of that matrix identity in the coefficient field. This establishes nondegeneracy and fixes the internal component coefficients independently of a desired root.

For the product-family link introduce supersymmetric chiral messengers A in $(\bar{6}, 6)$, A_tilde in $(6, \bar{6})$, and B in $(10, 10)$, with dimensions 36, 36 and 100. These are Standard Model singlets and neutral under the six shaping factors. The superpotential chain contains $M_6 A \cdot A_\text{tilde}$, $M_{10} B \cdot B$ divided by two, $y_2 A \cdot (\Lambda \Lambda)$ projected to the 36, and $y_3 B \cdot (A_\text{tilde} \Lambda)$ projected to the 100. Every displayed family contraction is unique. The CP action on these messenger fields is inherited from the symmetric-square and symmetric-cube actions of the already checked X ; no arbitrary new CP matrix is introduced.

Eliminating the massive fields at leading tree order generates a holomorphic operator proportional to $-(y_2 y_3)^2 I_6(\Lambda)/(2 M_6^2 M_{10})$, in compatible invariant normalizations. Thus the sextic is generated by renormalizable cubic vertices whose representation coefficients fix its component pattern. This is stronger than simply inserting a high-degree polynomial into the scalar potential. It is a supersymmetric messenger-sector construction, not a completed softly broken scalar model: the F-term scalar potential from the effective superpotential, soft terms, canonical kinetic corrections and physical vacuum selection must be matched separately.

The full audit counts all 34 possible monomial multisets of degrees one through three for these four chiral field species. There are no invariant linear terms, two independent mass contractions, and 15 cubic contractions: the determinant of Λ ; the two unique chain vertices; and four self-cubic contractions each for A , $A_{\tilde{}}$ and B . The unused self-cubics are not silently excluded. $A_{\tilde{}}$ self-cubics can contribute at degree six to the same two-dimensional link-invariant space. The A and B self-cubics first affect higher orders in the small-link expansion for this massive branch. Hence the generated orientation tensor is fixed, while its coefficient and the determinant-squared coefficient remain independently adjustable. Other branches and the regime of large link expectation values require the full messenger equations.

The extra messenger fields do not couple holomorphically to the charged source flavons at renormalizable degree under the listed shaping charges. This controls a concrete set of interactions, while higher-degree Kahler and effective scalar terms can still generate the dressed cross-sector overlaps. No radiative absence theorem for those overlaps is asserted.

Complete mediator matching with the reverse interaction included

Assign Q_L to $(3,1)$. The up-sector mediator remains a triplet of G_u . For the down sector introduce D_u in $(3,1)$ and D_d in $(1,3)$, both vectorlike Standard Model weak singlets of down electric charge. The right-handed down singlets couple to D_d through the three down flavons. The allowed link Yukawa couplings run in both directions: $g \bar{D}_u L \Lambda D_d R$ and $g_{\text{reverse}} \bar{D}_d L \Lambda^\dagger D_u R$. Both are allowed by the same mass and shaping charges. Omitting the reverse contraction would incorrectly restrict the operator space.

At $H=0$ the down heavy block is $[B, M]$, with $B=[0; C_d]$ and $M=[M_u I, g \Lambda; g_{\text{reverse}} \Lambda^\dagger, M_d I]$. The light normalized right frame is $[I; -A] K$ to the power $-1/2$, where $A=M^{-1} B$ and $K=I+A^\dagger A$. The Higgs couples Q_L to $D_u R$, giving $Y_d=-h A_{\text{top}} K$ to the power $-1/2$. For an invertible M there are exactly three right light modes and six massive down mediator modes.

For $\Lambda=\ell U$ and real parameters, write $D=M_u M_d - g g_{\text{reverse}} \ell^2$. Then $A_{\text{top}}=-g \Lambda C_d/D$ and $A_{\text{bottom}}=M_u C_d/D$. Consequently $K=I+[(g^2 \ell^2 + M_u^2)/D^2] C_d^\dagger C_d$, and Y_d is a scalar times $U C_d$ times that canonical factor. The left frame is exactly U times the left singular frame of C_d . This includes the reverse link coupling and all orders in flavon mixing. The heavy masses must remain nonzero, requiring D nonzero. Finite-Higgs and quantum threshold corrections are separate calculations.

Physical CP established through quark invariants

$$J \neq 0, \quad \text{Im det}[H_u, H_d] \neq 0, \quad J_{\text{CP}} = -J$$

The numerical example uses a unitary W spanning the real fixed space of the anti-linear map, so $X \text{ conjugate}(W)=W$. Its source matrices are $C_u=W \text{ diag}(0.07,0.31,0.9)$ and $C_d=W \text{ diag}(0.08,0.27,0.85)$. All listed mediator couplings are real. We evaluate every group link $U=R(g)$, select the largest absolute J without a target phase or mixing anchor, and calculate the conjugate link $X \text{ conjugate}(U) X \text{ dagger}$.

After canonical matching the pair has $|J|$ approximately 0.0703201663, identical mixing magnitudes and opposite J signs. The chart phases are approximately plus and minus 94.6111562 degrees, and C_{eff} is approximately 1.50313413. An independent weak-basis check gives nonzero $\text{Im det}[H_u, H_d]$, approximately $7.09\text{e-}9$ in absolute value. Nonzero values with nondegenerate quark spectra demonstrate physical CP violation; complex representation entries or flavon phases alone would not do so.

This example is conditional on the chosen CP-real source frames and on a selected group link vacuum. The CP-real basis itself has a real orthogonal freedom, and the scalar model has not selected those frames. The large mixing is not phenomenologically viable. Real, independently allowed column-coupling changes in a nonorthogonal source example alter both C_{eff} and the physical phase. The receipt retains all cases. Thus physical CP is possible within a consistent restricted elementary symmetry, but the observed phase and the golden relation remain unpredicted.

The canonical global spectrum is gapped

The recovered six-dimensional real cap fiber is wedge squared of the standard four-dimensional A_5 representation, with its stored positive metric. Construct the covariant graph Laplacian using the actual 252 directed incidence transports on a 42-cap component. In integral coordinates its characteristic polynomial is $(t-2)15 (t-3)55 (t-5)21 (t-6)70 (t-7)21 (t-9)55 (t-10)15$. We compute and factor this polynomial exactly over the integers. The numerical metric-whitened Hermitian matrix independently reproduces the spectrum.

The base connection has dimension 252 and zero-dimensional kernel. Its orientation double cover has dimension 504 and the same spectral levels with doubled multiplicities: the other block is obtained by t mapping to $12-t$. Both have exact mass-squared gap two. On all 176 isomorphic cap components the base space contains 44,352 modes and has no zero mode. A parallel rank-three projector therefore does not yield three parallel vectors. The A_5 cover holonomy has no invariant vector in either irreducible golden triplet, consistent with the spectral result.

A global three-mode operator with explicit breaking

$$\dim \ker M^2 = 3, \quad \lambda_{\text{next}}(M^2) = \frac{1}{2}, \quad N = 44\,352$$

To exhibit exactly what an additional assumption can accomplish, choose one 42-cap component and a spanning tree, transport a root frame along that tree, and replace every edge matrix by the corresponding pure-gauge transport $R_w R_v$ inverse. This keeps the tree transports and changes 160 of the 252 directed edge matrices of the component. It trivializes the holonomy and is an explicit change to the canonical connection. It is not invariant under the full recovered M_{22} incidence action.

Transport the root rank-three Hodge projector by the same frames, and add a one-half penalty on its orthogonal complement. The scalar component Laplacian has exact spectrum 0 with multiplicity one, 4 with multiplicity 21, 7 with multiplicity six, and 9 with multiplicity 14. The

selected six-dimensional bundle therefore has three zero modes, and levels one half, 4, 4.5, 7, 7.5, 9 and 9.5 with multiplicities 3, 63, 63, 18, 18, 42 and 42 respectively. Gap every other cap fiber by an onsite mass squared of two, adding 44,100 massive modes.

This explicitly defined global mass-squared operator has exactly three zero modes among 44,352 modes and gap one half. Their normalized wavefunctions are transported root vectors divided by $\sqrt{42}$; the numerical mass and normalization residuals are below $2e-15$. The spectral statement is exact from the scalar characteristic polynomial and pure-gauge similarity. Selecting the component breaks M22, and flattening its connection breaks further symmetry. We have not derived either selection dynamically. The operator is a positive Hermitian flavor mass-squared construction, not a gauge-invariant chiral quark UV completion. Its real A5 transport and the local complex Valentiner link model have not been unified into one global particle theory.

What remains after this push

The bare quartic deformation is now excluded by specified product symmetry, but the degree-six dressed operators survive. The CP action is explicit on the listed fields, and physical CP is demonstrated in canonically matched matrices, but its vacuum and numerical phase are not selected. The global spectrum is calculated, including an exact obstruction for the canonical connection and a conditional three-mode alternative, but the required breaking interactions and chirality have not been derived. These boundaries are sharper than another constructed golden root. A complete predictive theory must connect the product-family link, its mixed operator coefficients and CP-compatible scalar vacua to a dynamically selected global chiral spectrum.

Reproduction and receipts

Run `python python/develop_valentiner_cp.py`, `python python/develop_valentiner_link.py` and `python python/develop_m22_global_modes.py`, and `python python/develop_valentiner_messenger.py`. The computations use the existing exact triplet generators, sextic receipt and frozen cap transports. The new receipts are `valentiner_cp.json`, `valentiner_link.json`, `m22_global_modes.json` and `valentiner_messenger.json` under `receipts/m22_interactions`. The CP and polynomial identities, class map propagation, bifundamental invariant counts and characteristic-polynomial factors use exact arithmetic. The quark examples, matrix whitening and normalization residuals are numerical cross-checks. No experimental data or new Lean theorem is claimed.

Audit of the flavor search inputs

Audited baseline: `c7cdb4cf9814a7be006c17965678084b85211612`. Date: 6 October 2026. This audit covers the flavor search and its B5, M22, Valentiner, messenger and CP inputs. It is not an audit of every unrelated arithmetic application in the repository. Historical data and numerical examples remain pinned; newer measurements are compared separately.

The mathematical constructions are useful, but their combination is not yet a derived particle theory. The decisive unsupported step remains the identification of a selected scalar or representation invariant with a relation between the canonically normalized up and down quark eigenframes. The audit also identifies a concrete incompatibility in directly combining the ordinary quark example with the proposed supersymmetric messenger sector, and a missing scalar-potential calculation that can already be tested exactly.

Input ledger

Here “computed” means checked for the specified input object, not established as a law of nature. “Chosen” identifies an additional model assumption. “Missing” means no calculation in this research establishes the required physical identification.

Input or inference	Audit status	Consequence for the search
Exact three-generation unitary CKM parametrization	Correct within the three-light-generation decoupling limit	Vectorlike finite-Higgs effects require separate charged-current matching.
Relation $s_{13}=C*s_{12}*s_{23}$	Chosen phenomenological hypothesis	CKM unitarity alone does not imply it.
Golden value $C=(3-\sqrt{5})/2$	Training-nominated candidate	Neither B5 nor M22 has derived its identification with the observable coefficient.
$\delta=66$ degrees	Training-nominated phase	A scalar angle, a root-of-unity eigenphase and the standard CKM phase are different quantities until matched.
Original $V_{us}=.22431$, $V_{cb}=.0411$	Pinned historical measured central values	They have uncertainties; the 2026 combined V_{cb} central value is $.0407$.
Γ and V_{td}/V_{ts} training anchors	Explicit measured inputs with assumptions	Γ helps select 66 degrees; the mixing-ratio extraction assumes SM dynamics.
Independent Gaussian/split-normal likelihood	Chosen summary-data approximation	Missing cross-input covariance prevents treating it as a precision global fit.
Basis α , α_s , π , ϕ , m_p/m_e , rational alphabet, support penalty and phase grid	Chosen search domain and prior	A MAP formula in this domain is not a symmetry theorem or a prior-independent discovery.
Excluded observable keys in the training API	Correct software safeguard	Previously published observables were known; the exercise is retrospective, not blinded.
B5 action on five exponent axes	Correct combinatorics	Swapping physically labelled constants is not a physical symmetry. B5 supplies no primitive 60th-root eigenphase.

Input or inference	Audit status	Consequence for the search
Golden extremum of positive-definite symmetric integral 2-by-2 determinant-one matrices	Correct conditional spectral statement	Integrality, determinant, matrix choice, canonical kinetic metric and quark coupling are physical assumptions.
Six-harmonic potential selecting plus/minus 66 degrees	Constructed and exactly checked	Its coefficients were found with the desired root known; their physical origin is absent. An additive constant in the implementation does not affect minima.
Eight rational stationarity constraints; four on the six-term support	Correct over the rational coefficient field	They are not eight independent equations for arbitrary real couplings at one angle.
CP symmetry of the scalar potential	Correct	It permits CP-even deformations and pairs the two signs; it does not select the observed sign.
Generic 1% coupling sensitivity	Numerical study of the declared perturbation ensemble	Its interval is not an experimental confidence interval or an invariant measure of naturalness.
20% attraction basin and low-temperature occupation	Conditional dynamics with specified measure	Angular kinetic metric, initial measure, thermal history and CP-sign selection are not derived.
Recovered Wilson ledger and M22 algebra	Verified selected archival subset and exact reconstruction	The recovery is not the whole lost archive; provenance checks do not validate all historical physical claims.
Three-dimensional M22 multiplicity spaces	Correct	M22 acts trivially on their multiplicity factors, allowing arbitrary frame matrices. They are not faithful family triplets.
Canonical decorated A5 transport and orientation cover	Computed geometry and projector splitting	Rank-three parallel projectors do not imply three light quark modes; the canonical connection has zero kernel.
Central C60 subgroup in the twelvefold cover	Supported group-theoretic construction	Central phases cancel from neutral spectral-projector overlaps; they do not select a CKM frame.
3.A6 inside 3.A7 inside 3.M22	Abstract subgroup/cover route supported by character-table data	The specific global twisted quark bundle and its explicit lifted matrices have not been built.
Exact Valentiner triplet and low-degree invariant census	Computed for the specified generators and polynomial/tensor spaces	The unique sextic is a classical invariant. Its uniqueness fixes tensor components, not all independent Wilson coefficients.
Failure of complete residual frames and the tested two-involution route	Correct scoped obstructions	They exclude those unbroken constructions, not every model with controlled symmetry breaking.
Product family symmetry $G_u \times G_d$	Added model assumption	It is not a consequence of the recovered single M22 action.
Six separately charged source flavons, singlet right-handed quarks and chosen mediators	Added field/charge assumptions	The field content already contains three column labels. Their presence is not a derivation of three families.
Global versus gauged family and shaping symmetries	UV realization unspecified	A gauge completion and its anomaly constraints have not been established. Formal selection rules alone do not supply that completion.
Product symmetry forbids bare cross-sector overlap quartics	Correct for those representations	It still permits nine dressed degree-six overlaps and other source/link couplings.
58 CP-even renormalizable scalar parameters	Specified nonsupersymmetric scalar-sector count	It is not the parameter count of a supersymmetric completion including messenger scalars, Kähler terms and soft breaking.

Input or inference	Audit status	Consequence for the search
Positive link alignment at scaled group matrices	Correct for the chosen polynomial potential	Unitarity, determinant stabilization and the degree-twelve completion were supplied; uniqueness and physical vacuum selection were not proved.
Renormalizable holomorphic sextic messenger chain	Representation counts and bilinear support the leading contraction	Other allowed self-cubics remain independent. The 17 invariant coefficients belong to the four-species messenger sector, not the whole theory.
Supersymmetric messenger superpotential versus quark link example	Separate constructions; direct amalgamation invalid	The quark reverse coupling uses the conjugate link. It cannot be promoted unchanged to a holomorphic superpotential with only the same chiral link field.
Source frames and column hierarchies	Chosen backgrounds and couplings	The CP-real basis has real orthogonal freedom. No joint scalar minimum selects its orientation or realistic mass hierarchy.
Canonical tree matching at zero Higgs background	Correct within the specified invertible heavy blocks	Thresholds, loop-induced kinetic corrections and finite-Higgs current normalization remain separate.
Scale at which the golden relation is imposed	Unspecified physical matching input	CKM parameters run above the weak scale. The standalone SM running diagnostic is not a threshold-matched evolution of the messenger construction.
Anti-linear involution on the chosen elementary fields	Exactly computed with a representation-content restriction	3.A6 has no class-inverting automorphism valid for arbitrary irreducible content. All added fields and contractions need compatible CP action.
Nonzero J and commutator invariant in the quark example	Correct conditional demonstration of weak CP violation	The group element was selected for large absolute J, not by a scalar vacuum; it gives unrealistic mixing and no prediction of the physical phase. Strong CP is unaddressed.
Modified 44,352-mode operator with three zero modes	Correct conditional mass-squared spectrum	Component choice, connection flattening and rank-three penalty are engineered breaking inputs. A chiral mass operator and the complex local model are not unified.
Observable polynomial relations	Exact consequences of the nominated CKM chart with branch conditions	Elimination creates a test of the hypothesis; it does not supply an independently justified interaction.
Reproducible artifacts and passing tests	Evidence of implementation consistency	Neither byte reproduction nor test count establishes experimental truth, radiative protection or novelty.

The rational-coupling qualification

Set $x = 2\cos(\theta)$. The minimal polynomial of $2\cos(66^\circ)$ has degree eight. A rational derivative vanishing at that root must be divisible by this polynomial, so it also vanishes at all its conjugates. The independent Chebyshev calculation reproduces rank eight for the constant plus twelve harmonics, and rank four on harmonics 2, 3, 5, 8, 9 and 12.

For unrestricted real coefficients, stationarity at one fixed interior angle is one nonzero linear equation. For example, $\cos(2\theta) + b\cos(3\theta)$ is stationary at 66 degrees when $b = -2\sin(132^\circ)/(3\sin(198^\circ))$. That example does not claim a minimum or a protecting symmetry. It isolates the coefficient-field assumption. Representation coefficients may be algebraic and independent physical couplings may be real, so the appropriate coefficient

field must be derived before promoting the rational rank to a model-building requirement. The exact CP-even deformation remains a valid obstruction in the scalar model where it is allowed.

The superpotential is not the chosen scalar alignment energy

In global supersymmetry with canonical kinetic terms, the scalar F-term potential is $\sum |dW/d\Lambda_{ij}|^2$; it is not $-\text{Re } W$. The audit reconstructs the fully symmetric sextic tensor from its exact polynomial coefficients. Its norm squared is 16 and its one-index reduced tensor is exactly $(16/3) I$. Hence, for $I6(\Lambda) = \langle T, T \text{ composed with } \Lambda \rangle$,

$I6(I) = 16$, $dI6/d\Lambda_{ij}$ at $I = 32 \delta_{ij}$, and $\sum \Lambda_{ij} dI6/d\Lambda_{ij}$ at $I = 96$.

With only $W = \kappa I6$, the radial line $\Lambda = rI$, for real positive r , has $VF = 3072 |\kappa|^2 r^{10}$. A nonzero scaled group matrix is therefore not radially stationary in this isolated model, while the origin has zero energy. Generating the sextic in W does not by itself generate the previously chosen nonzero alignment vacuum.

This is a limitation of the isolated term, not a no-go theorem for the allowed messenger EFT. The independently allowed determinant term can balance the gradient: for

$W = a \det(\Lambda) + \kappa I6(\Lambda)$, a nonzero rI is F-flat when $a + 32 \kappa r^3 = 0$.

Its group transforms are then F-flat as well. The origin remains F-flat, and the complete allowed terms, kinetic metric, source couplings and soft breaking must determine stability and occupation. This audit does not assert that those nonzero branches solve the CKM problem.

The ordinary down-quark example contains both a link and its Hermitian conjugate in its heavy mass block. That is allowed in the nonsupersymmetric theory. Holomorphy prevents copying both entries unchanged into a renormalizable superpotential with just one chiral link. A separate conjugate chiral link, a modified coupling chain, or a nonholomorphic effective interaction would require new fields or matching and a fresh operator audit. A hybrid construction may be possible; none was supplied by the prior calculations.

Updated data and the interpretation of the target

The 2026 PDG CKM review retains $|V_{us}| = .22431 \pm .00085$, and quotes combined $|V_{cb}| = .0407 \pm .0013$, $|V_{ub}| = .00389 \pm .00016$, and $\sin(2\beta) = .710 \pm .011$. Inclusive and exclusive determinations remain distinct, and the review's correlated global fit must not be treated as independent data added to these inputs.

Keeping the golden coefficient and 66-degree phase fixed while substituting only the updated central V_{cb} gives $|V_{ub}| = .00348716940$ and $\sin(2\beta) = .690881841$. These remain conditional chart predictions. The rounded combined measurement centers instead give $C_{\text{eff}} = .426088669$. First-order independent-error propagation gives about $.0222480$, putting the golden coefficient approximately 1.98 nominal standard deviations below that central combination. This diagnostic is not a joint likelihood or an exclusion significance. It establishes that the measured inputs do not currently fix the coefficient to the golden value. No new fit or posterior selection was run.

The existing retrospective search already reports that its leading formula has only .1603% posterior probability in the support-three family, its B5/smooth training evidence ratio is about .800, and removing ϕ barely changes its broad forecast. Those findings must remain inputs to our judgment. The narrow forecast conditional on choosing the golden formula does not include uncertainty over choosing that formula. The 66-degree phase was favored using measured γ ; its appearance in later constructed potentials is not independent evidence for it.

What the audit changes

The search should retain the recovered exact algebra, genuine triplet invariant tensors, canonical quark matching, explicit CP consistency calculation and the scoped obstructions. It should stop treating the golden target, product symmetry, CP-real source frames, unitary link, and three-mode modified spectrum as deductions from one common starting symmetry. They are different added inputs, and the physical maps between them remain missing.

A defensible next result would specify one field content and one Lagrangian, derive its scalar and kinetic terms, solve its joint source/link vacuum, and extract the two canonically normalized quark eigenframes without inserting the target. Allowed variations then test whether any observable relation is forced. A failed relation would still be an informative result. Recovering more elaborate arithmetic that only re-encodes the nominated angle would not repair the audited gaps.

Verification and sources

Run ``PYTHONPATH=python OPENBLAS_NUM_THREADS=1 python python/audit_flavor_inputs.py``. The deterministic receipt records the independent rational ranks, exact reduced tensor and sextic gradient, updated central-input calculation, and SHA-256 hashes of eleven audited implementation/input files. The historical training receipt is unchanged. The audit reran 65 flavor unit tests and eight Valentiner invariant/CP/global checks successfully. These checks do not independently reconstruct every archival M22 object or certify a complete UV theory.

Primary references: PDG, **CKM Quark-Mixing Matrix**, revised March 2026, <https://pdg.lbl.gov/2026/reviews/rpp2026-rev-ckm-matrix.pdf>, equations 12.8, 12.11, 12.12 and 12.20; S. P. Martin, **A Supersymmetry Primer**, <https://arxiv.org/abs/hep-ph/9709356>, sections 3.2 and 4.7; M.-C. Chen et al., **CP Violation from Finite Groups**, <https://arxiv.org/abs/1402.0507>, especially section 4's restricted-representation qualification. Existing character-table and generator references remain in the M22 and Valentiner chapters. The sextic-gradient and coefficient-field checks are independent calculations in this audit, rather than claims taken from these references.

Supersymmetric vacua and the CKM boundary

This calculation connects the previously separate link, scalar and quark constructions in a specified effective field theory. It derives an isolated group-aligned supersymmetric link branch and its complete local mass spectrum. It then proves that the minimal exact-supersymmetry source sector has only zero flavon vacua. A specified spurion-breaking extension supplies locally stable nonzero sources and a joint vacuum with physical weak CP violation. Finally an exact finite census excludes every orthogonal axis branch from realistic CKM mixing, and a permitted cross-sector interaction moves the solved quark frame.

Neither the golden coefficient nor 66 degrees enters the superpotential, source directions, breaking terms or vacuum parameters. The positive result is an explicit dynamical realization of discrete alignment and weak CP on a locally stable branch. The negative result is a sharp limitation of that branch, not a no-go theorem for broken Valentiner flavor models. The global family spectrum from M22 is not used as a physical input or claimed to be unified here.

One specified flavor effective theory

Take $G_u \times G_d$, with each factor the exact $3.A_6$ triplet group. The chiral link K transforms as $(\bar{3}_u, 3_d)$. There are three source flavons Φ_{ui} in $(\bar{3}_u, 1)$ and three Φ_{di} in $(1, \bar{3}_d)$. Each source has its own C_6 shaping charge; its corresponding right-handed conjugate quark has the opposite charge. All other family messengers in the displayed quark chain are neutral under these factors. The left-handed quark doublet Q transforms as $(3_u, 1)$. Two Higgs doublets and vectorlike quarks carry the usual Standard Model gauge representations. Matter parity and baryon conservation are assumed. The canonical kinetic metric, finite-order EFT truncation and eventual gauge/anomaly completion remain separate assumptions.

Write F_6 for the complex-conjugate polynomial coefficients of the previously derived Valentiner sextic, and I_6 for the corresponding holomorphic link invariant. On the link-and-source scalar sector, all nonconstant holomorphic invariants through field degree six give

$$W_{\text{flavor}} = a \det(K) + b I_6(K) + c \det(K)^2 + \sum_i \kappa_i F_6(\Phi_i).$$

There are nine independent contractions: the link determinant, its square, the new link sextic and six source sextics. Individual shaping charges exclude mixed link/source and mixed-source holomorphic terms at these degrees. This is a complete statement about this scalar superpotential sector, not a complete Kähler, breaking or higher-degree quark EFT enumeration. CP permits real coefficients in the checked convention. The source coefficients are independently nonzero; no golden relation among them is assumed. With fields of mass dimension one, a is dimensionless and b , c and κ have dimension minus three.

The physical Dirac row link is $L = \text{conjugate}(K)$. The quark superpotential contains $Q.H_u U^c$, $u_u U^c$, source terms $u_i^c \Phi_{ui}^T U$, $Q.H_d D_u^c$, $u_d D_u^c$, $md D_d D_d^c$, the forward term $g D_u^T K D_d^c$, and source terms $d_i^c \Phi_{di}^T D_d$. All column couplings are independent real inputs. A Higgs μ term and positive Higgs breaking masses keep the Higgs backgrounds zero in this flavor-scale calculation; electroweak breaking is not derived.

The earliest holomorphic reverse vertex is $\eta D_d^\dagger \text{adj}(K) D_u^c$. It has field degree four, with an independent coefficient of mass dimension minus one. The adjugate is a quadratic polynomial in K , and is defined even at singular K . It replaces the invalid attempt to put $K^\dagger$ in a superpotential with one chiral link. The corresponding physical heavy down-quark block is $M = [[\mu, I, g, L], [\eta \text{adj}(L), m_d, I]]$. Canonical tree matching is applied to this block, rather than copying the earlier nonsupersymmetric result. Additional higher-degree quark terms are not asserted absent.

The group-aligned branch and all nine link masses

$$r^3 = -\frac{a}{32b+2c}, \quad m_1 = |96b+6c| |r|^4, \quad m_8 = |72b| |r|^4$$

The exact symmetric tensor T has norm squared 16. Its two-index reduced tensor is $(4/3)(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk})$; all 81 entries are checked in the coefficient field. This gives $d^2 I_6(I)[E, F] = 40(\text{Tr}(E)\text{Tr}(F) + \text{Tr}(EF))$. Combined with the determinant derivatives, it establishes the F -flat branch

$r^3 = -a/(32b+2c)$, with $K = r \text{conjugate}(R(g))$.

At a nonzero r , every group transform has zero F energy. On that branch the superpotential Hessian is $r^4[(8b+2c)\text{Tr}(E)\text{Tr}(F) + 72b \text{Tr}(EF)]$. The nine complex link modes split into a singlet with fermion mass $|96b+6c||r|^4$ and eight modes with mass $|72b||r|^4$. Canonical scalar squared masses are their squares, with real multiplicities two and sixteen. Thus a nonzero branch is locally isolated in the link sector when b and $16b+c$ are nonzero, in addition to the existence conditions. No continuous family group is gauged here, so these eight massive directions are not Goldstone modes.

For $a = -.0324$, $b = 1$ and $c = .2$, the real branch has $r = .1$. Its fermion masses are .00972 and .0072. Four independent coefficient variations retain the nonzero branch and positive masses. The radius follows the coupling ratio; it is not a mixing-angle input. The origin also remains F -flat. Other F -flat link solutions and global occupation have not been classified.

There is a local protection statement worth separating from a CKM prediction. The diagonal family stabilizer of rI fixes only the scalar matrices, because $3 \text{ tensor } 3\text{bar}$ decomposes as $1+8$. An invariant superpotential gradient at rI must therefore lie in that one-dimensional fixed subspace. With a nonzero singlet Hessian, small invariant holomorphic deformations preserve a nearby scalar-radius branch by the implicit function theorem. They do not need the specially chosen coefficients of the original angular potential. This protects a local link alignment branch; source-dependent operators can remove that stabilizer and change the relative quark frame.

What exact supersymmetry does to the sources

$$\nabla F_6(\phi) = 0 \iff \phi = 0$$

The minimal source superpotential does not stabilize nonzero sources. The exact computation establishes $\text{grad } F_6(\phi) = 0$ if and only if $\phi = 0$. To verify this, write $u = x^2$, $v = y^2$ and $w = z^2$, turning F_6 into a cyclic cubic H . One-coordinate support has a nonzero derivative. The two-coordinate derivative resultant is $(67797-35235 i \sqrt{15})/32$, which is nonzero. On three-coordinate support, the two-variable Gröbner basis of H_u , H_v and H_w after $u=1$ is [1]. Cyclicity covers every projective patch. The independent test works directly with the original quintic derivatives, without the squared-coordinate reduction, and also obtains [1]. Conjugating the representation leaves the obstruction unchanged.

Consequently all six flavons vanish in the exact-SUSY, zero-matter, zero-Higgs vacuum slice for nonzero κ_i . Both Yukawa matrices then vanish. There is no CKM frame to extract. The eighteen complex source modes have zero quadratic mass, but their potential is lifted at degree ten; they are not flat moduli. This is an obstruction to this minimal field-content/truncation, not to models with driving fields, different charges or breaking interactions.

A specified breaking extension with stable nonzero sources

$$V_\phi = \kappa^2 \|\nabla F_6\|^2 - m^2 \|\phi\|^2 + 2A\kappa \text{Re } F_6$$

Add for each source $V_i = \kappa_i^2 \text{norm}(\text{grad } F_6)^2 - m_i^2 \text{norm}(\Phi_i)^2 + 2A_i \kappa_i \text{Re } F_6(\Phi_i)$. These are explicitly specified symmetry-respecting spurion-breaking EFT terms. They are not claimed to exhaust the allowed breaking action, and their origin or common coefficients are not derived. The independent column labels and their quark couplings are already field-content inputs, not a derivation of three families.

There is a stationary axis branch $\Phi = \rho e_j$ when $m_i^2 = \kappa_i^2 \rho^8 (180 + 6t)$, where $t = A_i / (\kappa_i \rho^4)$. The two transverse tensor coefficient magnitudes are exactly $\sqrt{45/2}$. In canonical real coordinates the six squared masses, divided by $\kappa_i^2 \rho^8$, are $1440 + 24t$, $-36t$, and each of $-90 - 6t$ plus/minus $(3\sqrt{10}/2)|48 + 2t|$ repeated twice. They are positive with positive m_i^2 throughout

$-30 < t < -(90 + 72\sqrt{10})/(6 + 3\sqrt{10})$.

At $t = -24$ they are 864, 864 and four copies of 54. The source vacuum energy is $-48 \kappa_i^2 \rho^{10}$, below the source origin. Every group transform of an axis has the same energy and spectrum. The $t = -26$, -24 and -22 cases are retained; direct six-dimensional real Hessians independently confirm their positive spectra. These are locally stable vacua. Other source extrema and the global winners are not classified.

Nonzero source F backgrounds also induce bilinear terms in the quark scalar potential. They cannot be ignored in a stability claim. At $\rho = .1$, $\kappa = 1$ and the stated real quark column couplings, their operator norm is at most $5.4e-5$. Adding a specified positive universal quark scalar breaking mass squared of $1e-4$ gives a rigorous lower bound $4.6e-5$ after including the nonnegative supersymmetric mass block. Explicit up and down scalar matrices have smallest squared masses approximately $9.99965e-5$ and $9.99985e-5$. Positive Higgs masses complete local stability in the declared zero-Higgs, zero-matter slice. This is not a threshold- or loop-matched complete Standard Model vacuum.

CP on a link and on a joint vacuum

For real CP-compatible coefficients, r^3 is real. The three cube-root phases differ by a central element of $3A_6$. The group link states form one family orbit. Acting with the elementary CP involution sends $rR(g)$ to $r^*R(u(g))$; a family transformation and, if needed, a central element restore the vacuum. The group link by itself therefore conserves a CP operation combined with family symmetry. Complex entries alone do not establish weak CP.

The nonzero source vacua change that conclusion. Exact field arithmetic identifies 45 axis rays, 30 unordered orthogonal frames, and two family orbits of fifteen frames each. The generator action on exact rank-one projectors, their orthogonality and the orbit partition are checked without clustering tolerances. Their 900 ordered up/down frame pairs cover every group link choice, because a group link permutes the ray and frame sets. Mass-order permutations only permute the displayed magnitudes or reverse the sign of J .

A joint stationary vacuum from the retained frame pairs has $J = -\sqrt{3}/32$, with an exactly computed invariant-product imaginary part. Canonically normalized quark matrices reproduce $J = -.0541265877$ and their CP conjugates give the opposite sign, with identical spectra and magnitudes. An independent weak-basis commutator determinant is nonzero. The worked pair has $|V_{us}| = |V_{cb}| = 1/2$, $|V_{ub}| = 1/\sqrt{2}$, and $C_{\text{eff}} = \sqrt{2}$. The frame pair was selected as the first nonzero- J witness, not by an energy preference. Its up and down spectra come from independently chosen real column couplings. This demonstrates physical weak CP in a locally stable branch; it does not select one sign or reproduce observed quark mixing. Strong CP is not solved.

A complete obstruction for the orthogonal axis branch

$$|V_{ij}| = 0 \quad \text{or} \quad |V_{ij}| \geq \frac{\sqrt{5}-1}{4} \approx 0.309$$

Every source column in this branch is a group-rotated coordinate ray. An overlap of two such rays is an entry of a group matrix, since its relative group product remains in $3A_6$. The implementation enumerates all exact squared entry values and compares them using a rational enclosure of $\sqrt{5}$. The smallest nonzero squared value is $(3-\sqrt{5})/8$, hence the smallest nonzero mixing magnitude is $(\sqrt{5}-1)/4 = .3090169944$. The ordering is certified by rational arithmetic, not inferred from a numerical minimum.

The entire orthogonal axis branch therefore has entries either zero or at least .3090169944. It cannot yield the observed nonzero $|V_{ub}|$ near .004, under any up/down frame pair, group link or mass permutation. This is considerably stronger than rejecting the displayed CP witness. It is a dynamical realization of a residual-frame obstruction: the scalar model has now produced the relevant stationary directions and positive local masses. Nonorthogonal source choices and other source extrema lie outside the obstruction and require their own joint-vacuum calculations.

The permitted interaction still moves the solved frame

$$H \Delta q = -\nabla O, \quad O = |\phi_{u1}^\dagger L \phi_{d1}|^2$$

The scalar EFT permits $\epsilon |\phi_{u1}^\dagger L \phi_{d1}|^2$ under both family factors, all six shaping charges and CP. At the solved CP witness, the first source overlap probability is $1/4$. Its constrained source torque is nonzero, with squared norm $r^4 \rho^6 B(1-B) = 1.875e-11$ in the declared units. Thus this interaction acts on an actual stable vacuum rather than on an assumed source frame.

The full 54-real-dimensional link/source Hessian is positive, with smallest eigenvalue about $5.4e-7$. Solving $H \delta x = -\text{grad } O$ gives a first-order stationary-vacuum response. Canonical quark matching along that response changes both magnitudes and weak CP: $dC_{\text{eff}}/d\epsilon$ is approximately 9.616 and $dJ/d\epsilon$ approximately .03590. These numbers use a numerical Hessian and dimensionful EFT normalization; they are not experimental error bars, a radiative coefficient estimate or a naturalness scale. The retained ϵ plus/minus $1e-6$ cases are linear continuations, not freshly minimized finite- ϵ vacua.

A mechanism protecting a precise relative frame must consequently constrain this operator or account for its effects. The product family group by itself does not do so. Generic small deformations preserve a nearby local minimum, but they need not preserve its CKM quantities exactly. This distinguishes stability of a vacuum from protection of an observable relation.

What the push establishes and what should follow

The earlier scalar-matching gap is now closed for a specified EFT: the determinant and sextic give a locally isolated nonzero link branch; a specified breaking sector gives nonzero sources and positive local spectra; a holomorphic cofactor replaces the conjugate-link reverse vertex; and physical weak CP is calculated from canonical quark matrices at a joint vacuum. The branch still fails the hierarchy and permits an explicit observable-changing cross-sector interaction. No target angle was supplied to obtain these conclusions.

The next useful search should explore nonorthogonal source vacua or new joint source/link contractions, while retaining the complete allowed breaking and kinetic operator spaces. It must derive their coefficients or demonstrate robustness across the independent ones. Adding a driving field or another representation is justified only if its operator census actually removes a demonstrated obstruction. The present calculation supplies concrete tests for that decision and avoids confusing another constructed golden root with a flavor prediction.

Reproduction and relation to established work

Run `PYTHONPATH=python OPENBLAS_NUM_THREADS=1 python python/develop_valentiner_susy_vacua.py`. Its scientific receipt records exact tensor identities, the source critical-locus reduction, parameter variations, the finite frame census, canonical CP pairs, scalar stability bounds and the allowed-response calculation. Run `python python/tests/test_valentiner_susy_vacua.py` for independent original-polynomial, Hessian, cofactor covariance, holomorphy, full fermion-mass and CP-orbit checks. Numerical and exact statements are identified separately. No new Lean theorem or experimental fit is claimed.

The classical Valentiner sextic and the general idea of supersymmetric flavon alignment are established mathematics and model-building tools. This draft does not claim their discovery. Primary background: S. P. Martin, A Supersymmetry Primer, <https://arxiv.org/abs/hep-ph/9709356> ; F. Feruglio, C. Hagedorn and L. Merlo, Vacuum Alignment in SUSY A4 Models, <https://arxiv.org/abs/0910.4058> ; C. Hagedorn, M. L. Lopez-Ibanez, M. J. Perez, M. H. Rahat and O. Vives, Flavon vacuum alignment beyond SUSY, <https://arxiv.org/abs/2312.07430> ; M.-C. Chen et al., CP Violation from Finite Groups, <https://arxiv.org/abs/1402.0507> . The cited alignment papers concern their own models, not a proof of our coefficients or CKM relation. The exact branch, spectra, finite census and response reported here are calculations for this declared construction. Publication novelty requires comparison and independent review.

The joint potential and the protection test

The nonorthogonal route is dynamically possible in the declared field content. This calculation enumerates the entire independent CP-even scalar polynomial space through degree six, activates every coefficient, and solves the full 54-real-field stationarity equations. It obtains positive local spectra, small nonzero mixing and physical weak CP. A shared-heavy-ray example also gives the ordering $|V_{ub}| < |V_{cb}| < |V_{us}|$. Neither example reproduces the measured CKM matrix, derives the quark-column hierarchy, selects a global vacuum or predicts a golden coefficient.

The decisive constraint comes from the allowed parameter space. On the retained nonorthogonal branch, independent scalar and quark couplings generate four independent physical observable directions, including after compensating the weights to hold all six quark masses fixed. Thus no exact nontrivial relation among the three mixing magnitudes and J holds throughout a neighborhood of this branch with these couplings free. Fully re-minimized one-percent scalar deformations directly change the effective coefficient, also after mass matching. The remaining task is to derive correlations among the interactions and quark weights, rather than extend an unconstrained potential search.

Field content, order and conventions

Keep the preceding $G_u \times G_d$ construction, with each factor the genuine $3.A_6$ triplet group. The physical row link L transforms as $g_u L g_d^\dagger$. Three physical source vectors u_i transform under g_u and three d_i under g_d . Their chiral counterparts are conjugates. Each source has a distinct C_6 shaping charge. CP acts through the previously checked involution X : L maps to $X \text{conjugate}(L) X^\dagger$ and each vector maps to $X \text{conjugate}(\phi)$. The quark chain includes the independent holomorphic reverse cofactor vertex. Its tree matching and right-handed canonical normalization are retained.

The word complete here has a specified scope: every real CP-even polynomial scalar contraction through total field degree six, on the zero-quark, zero-Higgs flavor slice, under these two finite family groups and all six independent shaping factors. Constants are discarded. The F potentials induced by the earlier superpotential are kept as a separately specified positive completion: link degrees four, seven and ten, and source degree ten. This is a scalar symmetry-breaking EFT. General Kähler terms, all higher-degree quark operators, loop matching, electroweak breaking and a gauge/anomaly completion are not included in this census. There are infinitely many higher-order allowed terms; no claim of an all-order scalar action is made.

Use $\rho=.1$ and express every link or source field in units of ρ , taking the reference mass scale $\Lambda=1$. Restoring it, the fields are $\Lambda \rho z$ and $V_{\text{physical}} = \Lambda^4 \rho^{10} V_{\text{scaled}}$. A degree- d coefficient c in the receipt means $c \Lambda^{(4-d)} \rho^{(10-d)}$. The raw real coordinates are $\text{Re}(z)$, $\text{Im}(z)$; canonical real squared masses are $\Lambda^2 \rho^{8/2}$ times their scaled Hessian eigenvalues. This factor is necessary because a canonical complex scalar is $(q+i p)/\sqrt{2}$.

The exact joint operator census

$$N_{\text{CP even}} = 7 + 1 + 41 + 7 + 207 = 263$$

The independent counts are as follows. The complex column counts polynomials before Hermiticity and CP; CP-even plus CP-odd real dimensions equal that complex dimension.

Field degree	CP-even real terms	Complex dimension
2	7	7
3	1	2
4	41	41
5	7	14
6	207	223
Total	263	287

An independent exact character calculation obtains the complex dimensions. For link bidegree (p,q) , Cauchy's decomposition expands $\text{Sym}^p(3 \otimes \bar{3})$ and its conjugate into Schur representations. Local Jacobi-Trudi characters are calculated in $\mathbb{Q}(\sqrt{5}, \omega)$, and averaged exactly on each 1080-element group, with the balanced source characters multiplied in. Separate C6 charges require each source to occur equally often with its conjugate at these orders, except for an individual holomorphic sixth power. The six source sextics and their conjugates are added explicitly. The receipt retains every contributing multidegree sector, not just totals.

Independence is also certified without floating rank tests. The balanced contractions have integer coefficients in real coordinates. Evaluate them at 260 reproducible integer-coordinate points modulo the prime 1000003, using formal Gaussian residues and conjugation. Full-rank square evaluation minors establish exact independence of the 7 quadratic, 41 quartic and 199 balanced sextic polynomials. Their pivot-row witnesses are retained. Distinct source/link phase bidegrees separate the remaining source and determinant sectors. The two holomorphic link sextics are independent already on $L = e_1 e_1$ -transpose, where $I_6 = 1$ but $\det(L)^2 = 0$. Multiplication by the nonzero determinant polynomial preserves independence of the quadratic factors used at degree five. Together with the character dimensions and CP graph pairing, this establishes the complete independent real basis.

The explicit contraction basis is constructed independently from trace and Gram graphs. Local finite-group tensor counts agree with $SU(3)$ through degree six on all balanced index sectors. The only additional finite-group contributions at this order are the individual source sextics and the link sextic. The group-specific sextic is therefore not used to remove any otherwise permitted balanced cross-sector interaction. CP pairs conjugate graph orientations. The two within-sector Gram triangles and the six link-dressed same-sector triangles give eight CP-odd balanced sextics; the holomorphic source and link pairs give eight more. Cubic and quintic conjugate pairs give respectively one and seven CP-odd terms. There are no CP-odd quadratic or quartic terms.

For an explicit basis, set $M_u = L L^\dagger$, $M_d = L^\dagger L$, $S = \text{Tr } M_u$, $T = \text{Tr } M_u^2$, $R = \text{Tr } M_u^3$, $D = \det L$, $N_i = \phi_i^\dagger \phi_i$, $Q_i = \phi_i^\dagger M_{\text{sector}} \phi_i$ and $Q_2 = \phi_i^\dagger M_{\text{sector}}^2 \phi_i$. Define $G_{ij} = \phi_i^\dagger \phi_j$ only for sources in the same sector. The operator receipt provides a name, degree and coefficient for every contraction.

At degree two the basis is S and the six N_i . Degree three contains $\text{Re } D$. Degree four contains S^2 and T , twelve $S N_i$ and Q_i terms, all twenty-one norm products $N_i N_j$, and the six within-sector $|G_{ij}|^2$. Degree five contains $S \text{Re } D$ and six $N_i \text{Re } D$ terms.

At degree six the link-only terms are S^3 , $S T$, R , $\text{Re } D^2$ and $\text{Re } I_6$. The twenty-four link-four/source-two terms are $S^2 N_i$, $T N_i$, $S Q_i$ and Q_2^i . The seventy-eight link-two/source-four terms comprise twenty-one $S N_i N_j$, thirty-six $Q_i N_j$, six $S |G_{ij}|^2$, six real dressed Gram triangles and nine $X_{ij}=|u_i \text{--} \dagger L d_j|^2$. The one hundred source-only terms are fifty-six triple norm products, thirty-six norm times within-sector overlap squares, two real Gram triangles and six $\text{Re } F_6(\phi_i)$. A redundant $|\det L|^2$ is eliminated by Newton's trace identity. This gives $5+24+78+100=207$ sextics.

All nine X_{ij} have independent coefficients. Their absolute squares are neutral under any additional Abelian phase assignments to the displayed fields: the charge of $u_i \text{--} \dagger L d_j$ cancels that of its conjugate. Ordinary phase charges alone therefore cannot forbid these norm-square operators, even if they constrain holomorphic vertices. A mechanism that relates or removes them needs an additional structural restriction, such as a justified interaction origin or a symmetry acting nontrivially on source labels. No such restriction is assumed here.

A bounded potential with every coefficient active

The scaled base potential is the squared gradient of $W_{\text{link}} = -32.4 \det L + I_6(L) + .2(\det L)^2$, plus the sum of six source potentials $\text{norm}(\text{grad } F_6(\phi_i))^2 - 36 \text{norm}(\phi_i)^2 - 48 \text{Re } F_6(\phi_i)$. Its nonzero link $L=I$ and all group-rotated coordinate sources reproduce the previously derived stationary branch. Sources need not be orthogonal to each other: each field independently occupies a stable axis orbit.

Add a real coefficient times every one of the 263 basis operators. The declared deterministic coefficient prescription uses seed 108066: small positive, distinct generic coefficients at the other terms, a positive .02 coefficient for S^3 , independent positive within-sector overlaps near .5 and independent negative X_{ij} coefficients near -.5. These are chosen EFT inputs. Their ratios are not derived or adjusted to the golden relation. Every coefficient is nonzero in the three mixed cases; the unmixed case is retained as the baseline. A common factor 0, .25, 1 or 4 scales the added polynomial.

The positive pure-link sextic is a deliberate boundedness condition, not a flavor target. By Cauchy-Schwarz, $|I_6(L)| \leq 16 S^3$, and $|\text{Re}(\det L)^2| \leq S^3/27$. Also $S T$ and R are bounded by S^3 . The retained pure-link sextic has a strictly positive lower-bound coefficient .0194628392 after subtracting all these absolute contributions. The established smoothness of F_6 implies a positive minimum of $\text{norm}(\text{grad } F_6)^2$ on the unit sphere, so it gives degree-ten radial growth for each source. Every mixed contraction has weighted growth $a/6 + b/10 < 1$, where a and b count link and source fields. Young's inequality absorbs such terms into the positive link-six and source-ten bounds. All positive-factor potentials are consequently coercive and possess global minima. The calculations identify local minima; they do not classify those global minima.

Nonorthogonal local vacua and physical CP

$$\text{Im}(G_{12}G_{23}G_{31}) = \frac{\sqrt{3}}{16}, \quad \det G = \frac{1}{8}$$

Start with the first full-rank axis-ray triple having a nonzero imaginary Gram triangle, and use it in both sectors. In the fixed enumeration its ray indices are 0,3,10. Exact field arithmetic gives $G_{12} G_{23} G_{31} = (1+i \sqrt{3})/16$ and $\det G = 1/8$. The individual source directions are genuine

group-rotated stationary axes, not a copied CKM chart. Each is stable before adding the mixed operators. The phase invariant exists despite real potential and quark coefficients.

There is a useful exact CP identity for any shared full-rank column matrix C . Write $H_u = C \text{diag}(a_i) C^\dagger$ and $H_d = C \text{diag}(b_i) C^\dagger$, with real weights, and $G = C^\dagger C$. Direct multiplication gives $[H_u, H_d] = C A C^\dagger$, where $A_{ij} = (a_i b_j - b_i a_j) G_{ij}$ and $A_{ii} = 0$. Therefore

$$\det[H_u, H_d] = 2i \det(G) (a_1 b_2 - b_1 a_2)(a_2 b_3 - b_2 a_3)(a_3 b_1 - b_3 a_1) \text{Im}(G_{12} G_{23} G_{31}).$$

For the retained triple this is $i \sqrt{3}/64$ times the three weight differences. The raw weighted matrices violate weak CP when that product is nonzero. On a scalar group-aligned link, the canonical up and down matching maps are strictly increasing spectral functions of these raw matrices, so their left projectors and physical J are unchanged. This proves the mechanism's CP content without identifying a complex entry with a physical phase. The perturbed link is handled by the full holomorphic mediator block and canonical matching, and CP is checked again there.

The quark-column inputs are $(.007, .05, .9)$ in the up sector and $(.02, .2, .85)$ in the down sector. These explicitly imposed hierarchies are not outputs of the scalar symmetry. The full 54-coordinate minimization, followed by Newton refinement, gives stationarity residuals below $4e-11$ across the four cases. Their raw Hessian minima range from 108 at the baseline to 102.70 at factor four. The independent weights preserve three nondegenerate quark modes in each sector.

At factor one the matched magnitudes are $|V_{us}| = .00139934$, $|V_{cb}| = .01621072$ and $|V_{ub}| = .00038814$, with $J = 8.64911e-9$ and standard phase 100.7496 degrees. The effective coefficient is 17.11066. These magnitudes escape the .309 lower bound of the orthogonal branch, but their ordering is not the observed Cabibbo hierarchy. They are a local capability example, not a data fit. The independent commutator determinant is nonzero, about $1.38e-34$ in this dimensionful normalization. CP conjugation preserves potential energy and magnitudes and reverses J .

The nonzero source F backgrounds also require a matter stability check. Their weighted Frobenius norm gives an upper bound approximately $7.55e-5$ on the source-induced quark-scalar bilinear operator norm. The specified positive universal matter breaking mass squared $1e-4$ leaves a rigorous positive lower bound approximately $2.45e-5$ after adding the nonnegative supersymmetric mass block. This check concerns the displayed matter action; arbitrary additional matter EFT operators are not included. Electroweak vacuum selection remains separate.

Shared heavy directions and the ordering of the mixing angles

A second illustration keeps the dominant source ray common to both sectors and varies the two lighter down-sector rays. There are 946 unordered pairs to inspect; 902 pass the declared determinant cutoff .05, and 467 give nonzero J with $V_{ub} < V_{cb} < V_{us}$ at the stated quark weights. These are numerical background diagnostics, not a complete classification of all nonorthogonal triples or a proof of energy selection.

The smallest- V_{us} pair within this explicitly ordered sample uses down rays 18, 21, 10. This selection is phenomenological and is reported as such; no measured anchor, target phase or golden coefficient is fitted. Fully minimizing the same 263-coefficient potential from that

starting point gives $|V_{us}|=.425322$, $|V_{cb}|=.0216236$, $|V_{ub}|=.0112854$ and $J=6.19010e-5$. Its stationarity residual is below $9e-12$ and its raw Hessian minimum is 106.15. The phase is 41.2339 degrees and $C_{\text{eff}}=1.22692$. The largest angle remains too large and the coefficient is not golden. The source sectors are nonorthogonal and the shared heavy direction only becomes approximate after relaxation.

This demonstrates how shared dominant directions and hierarchical weights can suppress mixing involving the heavy family while retaining a larger light-family angle and physical CP. It does not establish that the present symmetry explains either the weights or the measured hierarchy. The same potential has other local branches and no cosmological occupation rule. The example supplies a concrete test case for a proposed source-label interaction.

Why nonorthogonality changes the protection problem

$$H\Delta q = -\nabla O, \quad \text{rank} \frac{\partial(|V_{us}|, |V_{cb}|, |V_{ub}|, J)}{\partial c} = 4$$

There is an exact elementary theorem behind the numerical sensitivity. For a fixed full-rank column matrix C , let $H(t)=\sum_i t_i c_i c_i^\dagger$, with all t_i independently positive. If its unordered nondegenerate left eigenprojectors stay fixed on an open parameter set, then differentiating H with respect to each t_i shows that every rank-one matrix $c_i c_i^\dagger$ is diagonal in that same eigenbasis. They therefore commute. Two nonzero rank-one projectors commute only when their vectors are parallel or orthogonal; full rank excludes parallel columns. The columns must be pairwise orthogonal. Conversely orthogonal columns share their projectors throughout every fixed mass-order region.

Thus nonorthogonal sources solve one problem and introduce another obligation: the quark-column coefficients affect the mixing frame. The scalar potential on the zero-matter/Higgs slice does not determine those coefficients. This theorem applies directly to the universal up mediator and to both sectors on a scalar aligned link. On the deformed link the fully matched numerical calculation establishes the response. The theorem concerns the full frame; it does not by itself rule out a specially protected single observable identity at another exceptional configuration.

At the actual factor-one vacuum, calculate the response to six same-sector overlap coefficients, all nine X_{ij} and all six logarithmic quark weights. Scalar variations solve $H \delta x = -\text{grad } O$ using the full 54-dimensional Hessian. The resulting Jacobian of $(|V_{us}|, |V_{cb}|, |V_{ub}|, J)$ has rank four. Because J is small, the raw matrix is poorly scaled. After dividing each observable by its nonzero reference magnitude and normalizing the parameter columns, its singular values are approximately 3.75815, 2.06946, 1.60725, .10196. The fourth direction is well separated from numerical roundoff in these coordinates.

The implicit function theorem supplies nearby scalar vacua, and the inverse function theorem applied to a nonsingular four-column observable minor supplies a neighborhood of physically independent observable values. On this regular local branch with the specified couplings free, an exact golden amplitude relation or a fixed CP phase therefore cannot be an identity. This is a local numerical-rank statement supported by explicit positive masses, not a theorem excluding every possible branch, new representation or correlated UV completion.

The stronger calculation holds the six quark singular values fixed. Write the six log-mass response as $M_{\text{scalar}} \delta c + M_{\text{weights}} \delta_{\log} \lambda$. The weight block is invertible, with singular values between .9845 and .9999. Set $\delta_{\log} \lambda = -M_{\text{weights}}^{-1} M_{\text{scalar}} \delta c$ and substitute it into the mixing response. The remaining fifteen scalar directions still generate all four physical mixing directions. With the same fractional-observable and column

normalization, their singular values are 3.26373,1.59728,1.33898,.06274. Thus the freedom is not removed by calibrating all six quark masses. These reference masses are the illustrative masses of this vacuum, not experimental measurements.

Fully re-minimized finite deformations

$$C_{\text{eff}} = \frac{|V_{ub}|(1 - |V_{ub}|^2)}{|V_{us}||V_{cb}|}, \quad C_{\text{eff}}^2 - 3C_{\text{eff}} + 1 = 0$$

The nominated coefficient is $C_{\text{eff}} = |V_{ub}|(1 - |V_{ub}|^2)/(|V_{us}||V_{cb}|)$. The golden condition requires $C_{\text{eff}}^2 - 3C_{\text{eff}} + 1 = 0$ with $0 < C_{\text{eff}} < 1$. At the retained vacuum, the golden polynomial has nonzero derivatives along independent allowed directions. This is not an artifact of an off-vacuum source frame or an unnormalized quark matrix.

Perturb the X_{02} coefficient by plus and minus one percent of its absolute value, and solve the stationarity equations again. C_{eff} changes from 17.11066 to approximately 18.22351 and 15.99458. Perturbing the B_{02} same-sector overlap gives approximately 15.88999 and 18.32730. All four fresh vacua have positive Hessians and stationarity residuals below $2e-10$. Their receipt contains full observables and stability diagnostics. These are dimensionful EFT perturbations of the explicitly chosen coefficient values, not experimental error ranges or estimates of radiative corrections.

After re-solving each vacuum, independently refit all six positive quark-column weights to recover the baseline six masses. Their maximum log-mass residual is below $2e-12$. The X_{02} deformations still give C_{eff} approximately 18.22682 and 15.99157; the B_{02} deformations give 15.89963 and 18.31621. Since the tree-level zero-matter/Higgs scalar potential is independent of these quark weights, this compensation preserves scalar stationarity. This supplies finite examples of observable change at fixed masses, in addition to the local differential-rank test.

The next interaction must constrain both sectors

An unconstrained joint-potential search can now be distinguished from a flavor prediction. The declared field content admits nonorthogonal stable vacua, weak CP and hierarchy mechanisms. Its complete low-degree operator space and independent quark weights leave all four physical mixing parameters adjustable on the tested branch. Finding a golden-looking point in that space would select couplings rather than explain a relation.

The next construction needs a justified non-Abelian action on source labels or a UV interaction whose unique invariant contractions correlate the quark-column weights and the nine mixed scalar coefficients. It must retain its allowed norm-square operators and kinetic corrections in the audit. A larger Abelian phase group cannot forbid the X_{ij} . A proposed M22-derived interaction must specify which of these actual contractions it relates, rather than supply an arbitrary matrix on a multiplicity space. New fields are useful only if their full operator space reduces the demonstrated physical freedom. No specific new representation is asserted here to achieve that reduction.

Verification and established context

Run `OPENBLAS_NUM_THREADS=1 PYTHONPATH=python python python/develop_valentiner_joint.py` to reproduce the exact multidegree census, complete coefficient list, four mixed-potential cases, CP partner, shared-heavy-ray witness, 21-parameter response and four freshly minimized deformations. The field coordinates and every independent scalar coefficient are retained. The tests independently check all operator covariance under family generators, all six shaping factors and CP, every directional derivative, an independently contracted link F potential, the shared-column CP determinant identity, full stationary spectra, finite deformation replay and the eigenframe weight theorem.

Cross-coupling-driven reorientation and flavon alignment are established model-building problems. Relevant primary comparisons are M. Holthausen and M. A. Schmidt, Natural Vacuum Alignment from Group Theory: The Minimal Case, <https://arxiv.org/abs/1111.1730> ; and I. de Medeiros Varzielas, M.-S. Liu, A. Sengupta and J. Talbert, Residual Symmetries and Scalar Multiplet Vacuum Alignment in Non-Abelian Flavour Models, <https://arxiv.org/abs/2512.19789> . Their group models differ from this construction. The present results are the explicit operator census, nonorthogonal branch calculations, CP invariant and local freedom test for this specified EFT. Publication novelty and robustness beyond the declared truncation require independent review.

CP-even kinetic freedom at fixed quark masses

Result and assumptions

A complete Hermitian quark metric can be generated by nine CP-compatible polynomial covariants with real coefficients. When their CP triangle is nonzero, they span all nine real metric directions at the reference vacuum. Small positive changes of the bare right-handed metric can preserve a previously solved scalar vacuum, its superpotential and all six quark masses while changing every physical CKM parameter. This is an algebraic statement about allowed operators, rather than a claim that quantum corrections have a particular size. It makes a protecting model responsible for kinetic matching as well as scalar and Yukawa interactions.

The assumptions are linear unitary internal flavor transformations, flavor-covariant functions X and Z with the same left flavor representation, a full-rank X with distinct singular values, a nonzero CP triangle, and independent real coefficients for the complete allowed matter kinetic operator space. X and Z may have different right representations and ordinary phase charges: those cancel from $R=X^\dagger X$ and $S=X^\dagger Z Z^\dagger X$. The degree bound below requires polynomial covariants. Matched analytic Yukawa functions obey the spanning argument but require an effective-field-theory expansion before a polynomial degree can be assigned. Degenerate spectra, CP-conserving vacua, coefficient restrictions imposed by a UV theory and additional restrictions beyond these internal transformation laws are outside this theorem.

The numerical demonstration uses the already minimized nonorthogonal joint vacuum from commit f76767e. The golden-angle test is a separately initialized attack fixture. It does not turn the nominated angle or golden coefficient into a dynamical prediction. The accompanying common-label construction supplies a justified interaction and a stable heavy-family branch, but has not been solved as a complete joint three-family model.

Nine real polynomial covariants

$$R = X^\dagger X, \quad S = X^\dagger Z Z^\dagger X, \quad \det \mathcal{M} = \tau^3$$

Let $R=X^\dagger X$ and $S=X^\dagger Z Z^\dagger X$. Both transform by conjugation on the right flavor space of X . Under generalized CP they transform into their complex conjugates conjugated by the corresponding unitary CP matrix. Fix the three positive distinct eigenvalues $r_1 < r_2 < r_3$ of R at a reference vacuum as real numerical constants. Define $P_i(R)$ as the product over j unequal to i of $(R - r_j I)/(r_i - r_j)$. The P_i are genuine matrix polynomials with fixed coefficients, rather than eigenprojectors recomputed as the fields vary. At the reference vacuum they are the three orthogonal rank-one spectral projectors.

Take the three P_i and, for each pair $i < j$, the two Hermitian covariants $B_{ij}=P_i S P_j + P_j S P_i$ and $D_{ij}=P_i S^2 P_j + P_j S^2 P_i$. Every coefficient is real, every expression is Hermitian, and each expression has the required family and CP transformation law. In particular, there is no explicitly imaginary coefficient inserted to manufacture a missing phase direction.

In an R eigenbasis put $\tau = \text{Im}(S_{12} S_{23} S_{31})$. For the pairs (12),(13),(23), the real-coordinate determinants of the two complex off-diagonal entries S_{ij} and $(S^2)_{ij}$ are respectively

$-\tau, +\tau, -\tau$. The diagonal entries of S only add real multiples of S_{ij} and cancel from these determinants. The three diagonal projectors span the diagonal subspace. Consequently the nine-generator coordinate determinant is τ -cubed in coordinates consisting of the three diagonals followed by real and imaginary off-diagonal entries. It is 8τ -cubed if both off-diagonal coordinates are multiplied by square-root two to obtain Hilbert-Schmidt orthonormal coordinates. This proves completeness when τ is nonzero. The exact symbolic identities are independently checked in the driver and focused tests.

An invariant certificate of physical CP

$$\text{Im Tr}(R^2 S^2 R S) = \Delta(R) \tau$$

The reference-basis condition can be written without choosing an eigenbasis: $\text{Im Tr}(R\text{-squared } S\text{-squared } R S) = \Delta(R) \tau$, where $\Delta(R) = (r_2 - r_1)(r_3 - r_1)(r_3 - r_2)$. This identity is checked by symbolic expansion of a general Hermitian S . A nonzero left-hand side certifies that the covariants do not admit a simultaneous CP-real frame.

If X and Z are the canonically normalized up and down Yukawa functions, S in the R eigenbasis is the up singular-value diagonal matrix times the down left mass-squared matrix times the same diagonal matrix. Its triangle imaginary part is the product of the three squared up singular values times the triangle imaginary part of the down matrix in the up mass basis. For nondegenerate spectra that latter quantity is proportional to the physical Jarlskog invariant and the down spectral Vandermonde. Thus nonzero physical weak CP supplies the required spanning condition for the matched Yukawa functions. For the finite polynomial covariants used in the explicit heavy-mediator example, the condition is checked directly; it is not inferred merely from the existence of a phase in their entries.

Positive completion and finite degree

$$B(\Phi) = (I + T(\Phi))^2 + \eta I \geq \eta I$$

At the reference vacuum orthonormalize the three off-diagonal pairs using real Hilbert-Schmidt inner products. This uses fixed real normalization constants. Together with the three projectors the resulting Q_a form a real orthonormal Hermitian basis. Given any positive target right-handed metric B_0 , choose a positive η strictly smaller than its least eigenvalue, set $T_0 = \text{square-root}(B_0 - \eta I) - I$, and expand T_0 as the sum of the nine Q_a times the real coefficients $\text{Tr}(Q_a T_0)$. Use the same coefficients and fixed polynomial recipes at every field value to define $T(\phi)$.

The metric $B(\phi) = (I + T(\phi))^2 + \eta I$ equals B_0 at the reference point and is bounded below by ηI at every field value. Positivity follows because T is Hermitian. This is a positive completion of the right-handed block. It is not a global positivity theorem for the entire Kahler metric, whose mixed matter-flavon blocks appear away from zero matter. Near zero matter backgrounds those mixed blocks vanish and positivity of the complete metric follows locally by continuity.

If X has flavor-field degree p and Z degree q , then R has degree $2p$, S has degree $2p+2q$, and each interpolating P_i has degree at most $4p$. The highest-degree generator $P_i S\text{-squared } P_j$ plus its adjoint has degree at most $12p+4q$. Squaring the Hermitian completion gives an upper bound $24p+8q$ on the flavor-field degree of B . A matter bilinear adds two matter fields to the operator degree. The explicit down-sector example uses $X = L C_d$ and $Z = C_u$, hence $p=2, q=1$ and a maximum flavor-field degree of 56. High-degree coefficients and reference spectral

denominators can be large; no naturalness or loop-size estimate follows from this existence proof.

Fixed masses with full heavy matching

$$B = Y_0^\dagger H_{\text{target}}^{-1} Y_0 - A^\dagger A, \quad Y Y^\dagger = H_{\text{target}}$$

The bare light right-handed quarks mix with heavy quarks through a fixed holomorphic null frame $N=(I,-A)$. Let Y_0 be the corresponding unnormalized Yukawa matrix. With the old identity bare metric the effective metric is $K_0=I+A^\dagger A$ and the matched Yukawa is $Y_0 K_0$ to the power $-1/2$. Replace only the bare light-quark metric by B . The effective null-frame metric becomes $K=B+A^\dagger A$.

For any sufficiently nearby positive target left mass-squared matrix H_{target} , choose $B=Y_0^\dagger H_{\text{target}}^{-1} Y_0 - A^\dagger A$. Then $K=Y_0^\dagger H_{\text{target}}^{-1} Y_0$ and, for invertible Y_0 , the canonically matched Yukawa obeys $Y Y^\dagger = H_{\text{target}}$ exactly. At the reference target this B is the identity, so positivity persists for a neighborhood of targets. Taking $H_{\text{target}}=U H_{\text{down}} U^\dagger$ preserves all down masses. The up sector is untouched. Relative unitary rotations cover the four physical mixing directions after rephasings, at a generic three-family point.

Every such B can be realized by the CP-compatible positive polynomial completion above. In the chiral-superfield convention its transposed right-handed metric is used if required by the definition of the physical right-handed column; the positive spectrum and covariance statement are unchanged. This matching is performed before eliminating the heavy fields. It does not impose an arbitrary matrix after canonical normalization and call that matrix a symmetry-derived contraction.

At zero quark, heavy-quark and Higgs backgrounds, the additional Kahler term is a matter bilinear and vanishes. Its mixed derivatives vanish there as well, and its flavon block does not change. Therefore the existing scalar stationary point and flavor Hessian remain unchanged at this tree-level background. The claim does not include an electroweak vacuum calculation or a radiatively corrected scalar vacuum.

Constructive checks on the solved joint vacuum

The driver reconstructs the unchanged up and down superpotential matrices and the complete down heavy null frame of the prior joint solution. It uses the polynomial covariants $X=L C d$ and $Z=C u$ on the bare down-quark label space. Six anti-Hermitian rotations in the up mass basis, each evaluated with both signs at magnitude 0.00001, produce twelve positive bare metrics. All twelve are realized with real polynomial coefficients and matched again through their heavy null-frame metrics.

The four singular values of the fractional-observable, column-normalized response matrix are 1.94061, 1.18728, 0.736859, 0.530491. These normalization-dependent numbers certify four independent observable directions; they are not physical sensitivity eigenvalues. The greatest logarithmic residual among all six quark singular values is below 7 times 10 to the power -13 in double-precision SVD. The nine-basis orthonormality error and positive metric reconstruction errors are below 10 to the power -75 in 90-digit arithmetic. After explicitly Hermitianizing the floating-point target mass matrix, the exact-matching residual is likewise checked at multiprecision. The driver records each metric, its eigenvalue bound, its nine real coefficients and all resulting observables.

A separate golden-chart fixture starts with $C=(3\text{-square-root five})/2$, physical phase 66 degrees, $|V_{us}|=0.2253$ and $|V_{cb}|=0.0409$. Holding both anchors, the phase and all six masses fixed, two CP-compatible positive metrics change C by -0.1% and $+0.1\%$. Their operator-norm distance from the identity is about 0.00343545, and their least eigenvalue exceeds 0.99657. This is a concrete allowed deformation of an initialized exact relation. It neither predicts these inputs nor establishes the typical size of kinetic corrections in a natural UV completion.

A shared non-Abelian source label

$$Z_8 = Z - \frac{\text{Tr} Z}{3} I, \quad V_{\text{eff}} = -\frac{k^2}{M^2} \|Z_8\|^2$$

The positive interaction candidate replaces three independently charged source columns in each sector by a single matrix source with one common phase charge. Introduce a common label group $H=3.A_6$ and sources $A=(3_u, 3_{\bar{H}})$, $B=(3_d, 3_{\bar{H}})$, together with one right-handed H triplet per quark charge sector. The common group acts on source labels as a representation, rather than permitting three arbitrary column coefficients. The exact singlet multiplicity in triplet times conjugate triplet is one, so the corresponding quark-source contraction has one coupling per sector.

For the link $L=(3_u, 3_{\bar{d}})$, $Z=A\text{-dagger} L B$ transforms by conjugation under H . The endomorphism representation decomposes as $1+8$. The two quadratic dressed channels are the singlet norm $|\text{Tr} Z|^2$ and the traceless norm $\text{Tr}(Z\text{-dagger} Z) - |\text{Tr} Z|^2/3$. Exact averaging over all 1080 group elements gives singlet multiplicities 1 in triplet times conjugate triplet, 1 in adjoint squared and 2 in adjoint cubed. These are representation-derived coupling counts.

A complex adjoint scalar S_H , with the phase charge of Z , permits $M\text{-squared} \|S_H\|^2 - 2k \text{Re Tr}(S_H\text{-dagger} Z_8)$, where $Z_8=Z-(\text{Tr} Z)I/3$. Eliminating S_H at this leading order gives $-(k\text{-squared}/M\text{-squared})\|Z_8\|^2$. This correlates the two norm coefficients through the fixed $-1/3$ trace-subtraction ratio. The ratio is a Clebsch coefficient and is not the golden coefficient. The source-link-mediator term is quartic in elementary scalar fields. Additional allowed mediator quartics and mediator norm couplings to the source and link bilinears must be included in a complete UV matching; the leading exchange is not a license to omit them.

A stable heavy-family branch

The same sextic invariant tensor gives a matrix-source invariant $I_6(A)$. Include the second allowed holomorphic sextic through $W_A=I_6(A)+c \det(A)\text{-squared}$, with real c . In dimensionless units consider $V_A=\|\text{gradient } W_A\|^2 - (180+6t)\|A\|^2 + 2t \text{Re } W_A$. The rank-one configuration $A=\text{diag}(1,0,0)$ is stationary, as are its finite group rotations. The determinant begins at second order around rank one, so its square and its first three derivatives vanish there. Hence its coefficient c does not alter the local Hessian.

In canonically normalized real coordinates the eighteen mass coefficients are: radial $1440+24t$; phase $-36t$; four copies each of $-90-6t$ minus and plus $3\text{ square-root ten times }|24+t|$; and four copies each of $-171-6t$ minus and plus $3|24+t|$. For $-36<t<-33$ all are positive and the energy $-144-4t$ lies below the origin. At $t=-34$ the energy is -8 . At $t=-34$, the spectrum is 3 with multiplicity four, $114-30\text{ square-root ten}$ with multiplicity four, 63 with multiplicity four, $114+30\text{ square-root ten}$ with multiplicity four, 624 and 1224. An independent eighteen-dimensional finite-difference Hessian reproduces this spectrum.

With the unique universal quark-source contraction this branch has one heavy family and two exactly massless light families before rank lifting. It supplies a stable hierarchy starting point without assigning three independent column weights. The complete joint potential, its rank-lifting operators and its CKM output have not been solved on this new field content. The kinetic theorem still applies if the completed branch has full-rank nondegenerate spectra and the required nonzero CP triangle.

What the result changes

A viable next UV construction must match its scalar potential, Yukawa contractions and matter metric together. The shared label group gives a justified contraction count and a heavy-family source branch, but internal symmetry with unrestricted allowed kinetic coefficients does not by itself protect an exact golden CKM equation at a CP-breaking vacuum. A finite-order prediction remains possible if omitted terms are controlled; an exact all-order relation requires stronger coefficient correlations. Merely adding phase charges, changing the group order or selecting a scalar stationary point does not supply those correlations for the operators constructed here.

The general importance of canonical normalization is established prior work. King and Peddie, Canonical normalisation and Yukawa matrices, Physics Letters B586 (2004) 83-94, arXiv:hep-ph/0312237, explicitly demonstrate changes in flavor textures. Espinosa and Ibarra, Flavour Symmetries and Kahler Operators, JHEP 0408 (2004) 010, arXiv:hep-ph/0405095, analyze its physical mixing effects and find subdominant quark effects under their phenomenological assumptions. Chen, Ramos-Sanchez and Ratz, A note on the predictions of models with modular flavor symmetries, arXiv:1909.06910, discuss additional Kahler parameters and predictivity. The contribution established in this repository is the explicit nine-polynomial spanning construction, its positive finite-degree completion, fixed-spectrum bare-mediator matching and reproducible application to the current joint vacuum. Publication priority has not been established by this targeted literature check.

Reproduction and certificates

Run `OPENBLAS_NUM_THREADS=1 PYTHONPATH=python python python/develop_flavor_kinetic.py` from the repository root. The inputs are the retained joint-vacuum and exact invariant-tensor receipts; the output is `receipts/m22_interactions/flavor_kinetic.json`. The module `python/perfectpower/flavor_kinetic.py` contains the reusable interpolating covariants, positive completion and bare matching. The focused tests cover exact symbolic identities, nine-dimensional spanning, family/phase/CP covariance away from the reference point, positive completion away from the vacuum, independent heavy-null-space matching, the retained four-direction response, the fixed-anchor golden deformation and all eighteen source masses. A fresh extraction repeats the scientific receipt byte for byte. The complete monograph and source archive retain the previous calculations and their stated boundaries.