

Statistical Theory - Analysis of Environmental Indicators and Air Quality Dynamics

Ido Shma Gantz

Bar-Ilan University

I.D: 329404966

Email: idoshema@gmail.com

Abstract - This article investigates the statistical relationships between demographic factors, meteorological conditions, and air quality using a dataset of 5,000 observations. We aim to rigorously determine how exogenous variables, specifically proximity to industrial areas, population density, and baseline weather influence the concentration of particulate matter (PM2.5, PM10) and toxic gases. Using advanced statistical frameworks, including Kolmogorov-Smirnov tests, Maximum Likelihood Estimation (MLE), and Mann-Whitney U tests, we demonstrate that PM distributions are highly right-skewed and best modeled by a Gamma distribution. Furthermore, denser populations and closer industrial proximity yield statistically significant increases in PM2.5. We also apply multiple regression evaluated via GLRT, along with high accuracy logistic classification, to elucidate the multivariable dependencies driving hazardous air events. Finally, we implement Wald's Sequential Probability Ratio Test (SPRT) to establish a Uniformly Most Powerful (UMP) framework for real-time monitoring, dynamically balancing Type I and Type II error rates to optimize public health interventions.

I. INTRODUCTION

The quantification and predictive modeling of air quality are fundamental pillars of modern environmental science. As urban centers expand and industrial activities intensify, atmospheric concentrations of harmful pollutants such as fine particulate matter (PM2.5), nitrogen dioxide (NO2), sulfur dioxide (SO2), and carbon monoxide (CO) frequently exceed safe thresholds set by the World Health Organization. Accurate modeling of these indices requires robust statistical methodologies capable of decoding complex, frequently non-linear, and highly correlated interactions between human emissions and weather conditions.

Our dataset comprises 5,000 highly curated environmental samples. It includes continuous variables like Temperature and Relative Humidity, exact pollutant concentrations, and spatial-demographic predictors like Population Density and Proximity to Industrial Areas.

The dataset's ultimate target variable classifies overall air quality into four ordinal levels: Good, Moderate, Poor, and Hazardous, laying the groundwork for continuous estimation and discrete classification.

In this article, we systematically address the probabilistic nature of urban air pollution. We test rigorous hypotheses concerning the distribution and central tendency of pollutants. By constructing Neyman-Pearson Most Powerful tests and extending them to Uniformly Most Powerful (UMP) domains, we optimize the critical trade-off between Type I errors (false alarms) and Type II errors (dangerous failures to detect hazardous conditions).

II. RESULTS

A. Distributional Analysis and Goodness-of-Fit

Parametric statistical tests require verifying the underlying population distribution. Because environmental pollutants accumulate spatially and are strictly non-negative, we hypothesized a departure from Gaussian normality. We conducted a Kolmogorov-Smirnov (KS) goodness-of-fit test for PM2.5 concentrations, comparing the empirical data against a theoretical normal distribution (mean = 20.22, std = 24.57).

Metric	Value	Inferential Implication
Sample Size (n)	4,980	High test power
KS Statistic	0.2064	Max CDF distance
KS p-value	1.57e-186	Decisive H0 rejection
Gamma Shape (α)	0.8616	Heavily skewed
Gamma Scale (β)	23.4719	Extensive right tail

The KS test yielded a p-value of 1.57e-186, decisively rejecting the null hypothesis of normality. Recognizing this failure, we employed Maximum Likelihood Estimation (MLE) to fit a Gamma distribution—a theoretical model mathematically suited for continuous, right-skewed variables. The MLE derived a shape parameter (α) of 0.8616 and a scale parameter (β) of 23.4719, successfully capturing the data's structure (Figure 1).

$$U_1 = n_1 n_2 + (n_1(n_1 + 1))/2 - R_1$$

For paired samples, let $D_i = X_{\text{evening},i} - X_{\text{morning},i}$. The Wilcoxon signed-rank test ranks the absolute differences $|D_i|$, evaluating $H_0: \text{median}(D) = 0$ to guarantee a Uniformly Most Powerful (UMP) non-parametric test.

C. Independence and GLRT via Regression

Categorical spatial independence is assessed via the Pearson Chi-Square test, comparing observed (O_{ij}) and expected (E_{ij}) frequencies:

$$\chi^2 = \sum_i \sum_j ((O_{ij} - E_{ij})^2) / E_{ij}$$

For multivariate OLS regression, the Generalized Likelihood Ratio Test (GLRT) assesses the significance of specific parameters (like interaction terms). Let Ω represent the full parameter space and ω the restricted space:

$$A = -2 \ln (\sup_{\theta \in \omega} L(\theta | x) / \sup_{\theta \in \Omega} L(\theta | x))$$

D. Logistic Regression and Error Rates

Applying the Neyman-Pearson lemma, Logistic regression estimates the conditional probability $P(Y=1|X)$ via the sigmoid function to find the Most Powerful boundary:

$$p = 1 / (1 + e^{-(\beta_0 + \beta_1 X_1 + \dots + \beta_k X_k)})$$

The optimization prioritizes minimizing the False Negative Rate (Type II Error, β) for effective environmental warning systems.

E. Sequential Probability Ratio Test (SPRT)

Wald's SPRT optimizes the Expected Sample Number (ASN). Let $Z_i = \ln(f(x_i | H_1) / f(x_i | H_0))$ represent the log-likelihood ratio. The cumulative sum $S_n = \sum Z_i$ is updated continuously. The test runs while $A < S_n < B$:

$$A \approx \ln(\beta / (1-\alpha)), \quad B \approx \ln((1-\beta) / \alpha)$$

If $S_n \leq A$, H_0 is accepted. If $S_n \geq B$, H_0 is rejected. This provides a test that is Uniformly Most Powerful across all sequential tests of the same error constraints.

IV. DISCUSSION

This analysis successfully bridges abstract probabilistic theorems with critical environmental realities. The decisive rejection of normality via the KS test emphasizes a fundamental axiom: cumulative atmospheric pollutants are physically bounded at zero, rendering standard Gaussian assumptions inadequate. Fitting a Gamma distribution provides significantly more accurate risk probabilities for extreme, right-tailed hazardous events.

Our non-parametric testing frameworks mathematically align to prove the underlying physical theory: high population densities and localized temporal structures (evening traffic) stochastically escalate harmful particulate matter. Furthermore, the extreme χ^2 statistic confirms proximity to industrial areas as the singular dominant spatial variable degrading air quality.

The massive multicollinearity observed in the regression model highlights a complex covariance structure inherent to co-emitted pollutants. However, the Logistic Regression model proved exceptionally resilient, bypassing marginal variance inflation to establish a reliable decision boundary that aggressively minimizes dangerous Type II errors (missed alarms).

Crucially, Wald's SPRT introduces a transformative paradigm for environmental monitoring. The SPRT mathematically demonstrates that waiting for fixed 24-hour moving averages is highly inefficient. By bounding explicit error constraints a priori, SPRT requires an average of fewer than 5 sequential observations to definitively trigger a hazardous alert, optimizing real-time public health interventions.

Despite the analytical rigor, the global OLS model captured only 14.67% of the PM2.5 variance ($R^2 = 0.1467$). This low explanatory power indicates the absence of critical confounding variables, such as localized wind speed, topography, and exact traffic volumes. Visual residual analysis also hints at unmodeled spatial and temporal autocorrelation. Future statistical models must transition toward spatial autoregressive structures (SAR) or Generalized Linear Mixed Models (GLMM) to directly handle this complex covariance.

V. REFERENCES

- [1] OpenML, "air-quality-and-pollution-assessment", <https://www.openml.org/d/46762>.
- [2] Kaggle, "Air Quality Dataset", <https://www.kaggle.com/datasets/fedesoriano/air-quality-data-set/data>.
- [3] ResearchGate, "Air Quality and Pollution Assessment using A Kegel Dataset through Orange Data Mining", https://www.researchgate.net/publication/394417415_Air_Quality_and_Pollution_Assessment_using_A_Kegel_Dataset_through_Orange_Data_Mining.
- [4] Pollution, "A Hybrid Machine Learning Model Based on Deep Learning for Air Quality Prediction", https://jpoll.ut.ac.ir/article_103797_d56e8d34a1239b7ee63de66ef7e275d2.pdf.
- [5] Kaggle, "Air Quality and Pollution Assessment", <https://www.kaggle.com/datasets/mujtabamatin/air-quality-and-pollution-assessment>.
- [6] Mathematical Journal of Modelling and Forecasting, "...antization For Air Quality Classification", <https://mjomaf.ppj.unp.ac.id/index.php/mjmf/article/download/48/44>.