

---

**Final Project of HPPL course**

# **Crank-Nicolson technique for solving the Schrodinger equation**

Ivan Gurev  
Nikolay Kotoyants

# Problem statement

---

The time-dependent Schrödinger equation (TDSE) is the fundamental equation of non-relativistic quantum mechanics. The equation has found widespread application in many fields of physical science, for example in atomic, molecular, optical, nuclear and solid-state physics, among others.

$$i\hbar \frac{\partial}{\partial t} \Psi(x, t) = \hat{H} \Psi(x, t)$$

# Importance/Relevance

---

For TDSE, it is challenging to acquire their analytic solutions. A forward difference Euler method has been used to compute the TDSE numerical solutions. Nevertheless, the Euler scheme is instability in some cases. That is why the Crank-Nicolson technique, which is unconditional stable was proposed to overcome this disadvantage

# The method itself

The Crank-Nicolson method, when applied to the Schrödinger equation, performs the following approximation at each time step

$$(2 + i\Delta t \mathbf{H}/\hbar) \vec{u}^{j+1} = (2 - i\Delta t \mathbf{H}/\hbar) \vec{u}^j.$$

which results in the  $(N_x + 1 \times N_x + 1)$ -matrix system

$$\begin{pmatrix} A_1 & 1 & 0 & 0 & \dots \\ 1 & A_2 & 1 & 0 & \dots \\ 0 & 1 & A_2 & 1 & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & A_{L-1} \end{pmatrix} \begin{pmatrix} \Psi_1^{n+1} \\ \Psi_2^{n+1} \\ \Psi_3^{n+1} \\ \dots \\ \Psi_{L-1}^{n+1} \end{pmatrix} = \begin{pmatrix} B_1 \\ B_2 \\ B_3 \\ \dots \\ B_{L-1} \end{pmatrix}$$

Initial state  $\Psi_i^0$   
Boundary conditions  $\Psi_0^n, \Psi_L^n$

# Thomas algorithm

Simplified Gaussian elimination, complexity  $O(n)$

Forward step:

$$U_1 = \frac{1}{A_1} \text{ and } R_1 = B_1 U_1$$

$$U_i = \frac{1}{A_i - U_{i-1}} \text{ and } R_i = (B_i - R_{i-1}) U_i$$

Backward step:

$$\Psi_i^{n+1} = \begin{cases} R_{L-1} & i = L - 1 \\ R_i - U_i \Psi_{i+1}^{n+1} & i < L - 1 \end{cases}$$

$$\begin{bmatrix} 1 & U_1 & 0 & \dots & 0 \\ 0 & 1 & U_2 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix} \begin{bmatrix} \Psi_1^{n+1} \\ \Psi_2^{n+1} \\ \Psi_3^{n+1} \\ \dots \\ \Psi_{L-1}^{n+1} \end{bmatrix} = \begin{bmatrix} R_1 \\ R_2 \\ R_3 \\ \dots \\ R_{L-1} \end{bmatrix}$$

# Our implementation

---

Initial condition - Gaussian wave packet:

$$\Psi(t=0) = \frac{1}{\pi^{1/4}\sqrt{\sigma}} \exp\left(\frac{-(x-\mu)^2}{2\sigma^2}\right) \exp(ip_0x/\hbar)$$

$$x \in [-10,10]$$

$$t \in [0,50]$$

$$m = 1$$

$$\Delta x = 0.1$$



$$\Delta t = 0.5$$

$$p_0 = 0$$


Boundary conditions:  $\Psi(0, t) = \Psi(L, t) = 0$

# Our implementation

Potential  $V(x) = 0$



# Double-well potential



Potential:


$$V(x) = \frac{a}{x_0^4}(x^2 - x_0^2)^2$$




# Complexity

Potential  $V(x) = 0$

$O(n)$



# Speed up



# 2D case


$$\mathbf{A} \cdot \mathbf{x} = \begin{pmatrix} a_{00} & -r_x & 0 & 0 & \cdots & 0 & -r_y & 0 & 0 & \cdots & 0 \\ -r_x & a_{10} & -r_x & 0 & \cdots & 0 & 0 & -r_y & 0 & \cdots & 0 \\ 0 & -r_x & a_{20} & -r_x & & & & -r_y & & & \\ 0 & 0 & -r_x & \ddots & \ddots & & & & \ddots & & \\ \vdots & \vdots & & \ddots & & & & & & \ddots & \\ 0 & & & & & & & & & & \\ -r_y & 0 & & & & & & & & & \\ 0 & -r_y & & & & & & & & & \\ 0 & 0 & -r_y & & & & & & \ddots & & \\ \vdots & \vdots & & \ddots & & & \ddots & \ddots & & & \\ 0 & 0 & & & & & \ddots & \ddots & -r_x & & \\ & & & & & & -r_x & a_{(N-1),(N-1)} & & & \end{pmatrix} \begin{pmatrix} \psi_{0,0}^{n+1} \\ \psi_{1,0}^{n+1} \\ \psi_{2,0}^{n+1} \\ \vdots \\ \psi_{i,j}^{n+1} \\ \vdots \\ \psi_{(N-1),(N-1)}^{n+1} \end{pmatrix}.$$

For 2D problem we have considered a 2-dimensional spatial grid (the xy plane) of N points in the x direction and N points in the y direction.


# Our implementation

---

Time evolution of free potential particle in two-dimensional case:



# Speed up



# Literature

---

1. David G. Robertson. “Solving the Time-Dependent Schrodinger Equation.”
2. Medium: “Quantum Mechanics with Python. Solving the 1D Time Independent Schrödinger Equation Numerically”
3. B. Ducomet AND A. Zlotnik “On stability of the Crank-Nicholson scheme with approximate transparent boundary condition for the Schrodinger equation”.
4. Amin Khan. “Numerical Solution of Schrödinger Equation by Crank–Nicolson Method”

# thx.

Skoltech

