

EXAM 4

MATH 1B, Bach

Tejas Patel, Roll #30

1 Identifying Quadric Surfaces

I

Horizontal Traces (xy plane): Ellipse
Vertical Traces (xz plane): Parabola
Vertical Traces (yz plane): Parabola
Surface: Elliptic Paraboloid

II

Horizontal Traces(xy plane): Ellipse
Vertical Traces (xz plane): Hyperbola
Vertical Traces (yz plane): Hyperbola
Surface: Hyperboloid of One Sheet

III

Horizontal Traces (xy plane): Hyperbola
Vertical Traces (xz plane): Ellipse
Vertical Traces (yz plane): Ellipse
Surface: Elliptic Cone

IV

Horizontal Traces(xy plane): Hyperbola
Vertical Traces(xz plane): Hyperbola
Vertical Traces(yz plane): Ellipse
Surface: Hyperboloid of Two Sheets

V

Horizontal Traces(xy plane): Hyperbola
Vertical Traces(xz plane): Parabola
Vertical Traces(yz plane): Parabola
Surface: Hyperbolic Paraboloid

VI

Horizontal Traces(xy plane): Ellipse
Vertical Traces(xz plane): Ellipse
Vertical Traces(yz plane): Ellipse
Surface: Ellipsoid

1.1 For equation (a): $\frac{x^2}{36} + \frac{y^2}{9} + \frac{z^2}{16} = 1$

Let $k = z$ (to test the x/y plane shape, the horizontal tracing)

$$\frac{x^2}{36} + \frac{y^2}{9} = 1 - \frac{z^2}{16} \text{ is an equation for an } \boxed{\text{ellipse}}$$

$$1 - \frac{z^2}{16} \neq 0$$

$$1 \neq \frac{z^2}{16}$$

$$16 \neq z^2$$

$$k \in \mathbb{R} \quad z \neq \pm 4$$

Now, let $k = y$ (to test the x/z plane shape, the vertical tracing)

$$\frac{x^2}{36} + \frac{k^2}{9} + \frac{z^2}{16} = 1$$

$$\frac{x^2}{36} + \frac{z^2}{16} = 1 - \frac{k^2}{9} \text{ is an equation for an } \boxed{\text{ellipse}}$$

$$1 - \frac{k^2}{9} \neq 0$$

$$1 - \frac{k^2}{9} \neq 0$$

$$1 \neq \frac{k^2}{9}$$

$$9 \neq k^2$$

$$k \in \mathbb{R} \quad k \neq \pm 3$$

Now, let $k = x$ (to test the y/z plane trace, the vertical tracing)

$$\frac{k^2}{36} + \frac{y^2}{9} + \frac{z^2}{16} = 1$$

$$\frac{y^2}{9} + \frac{z^2}{16} = 1 - \frac{k^2}{36} \text{ is an equation for an } \boxed{\text{ellipse}}$$

$$1 - \frac{k^2}{36} \neq 0$$

$$1 \neq \frac{k^2}{36}$$

$$36 \neq k^2$$

$$k \in \mathbb{R} \quad k \neq \pm 6$$

As the traces on all 3 planes are ellipses,
the shape is an **ellipsoid**, **figure VI (6)**

For equation (b): $\frac{x^2}{4} + \frac{y^2}{16} - z = 0$

Let $k = z$ (to test the x/y plane shape, the horizontal tracing)

$$\frac{x^2}{4} + \frac{y^2}{16} - k = 0$$

$$\frac{x^2}{4} + \frac{y^2}{16} = k \text{ This is an equation for an } \boxed{\text{Ellipse}}$$

$$k \in \mathbb{R} \quad k \neq 0$$

Now, let $k = x$ (to test the y/z plane shape, the vertical tracing)

$$\frac{k^2}{4} + \frac{y^2}{16} - z = 0$$

$$\frac{y^2}{16} - z = -\frac{k^2}{4} \text{ This is an equation for a } \boxed{\text{Parabola}}$$

$$-\frac{k^2}{4} \neq 0$$

$$k^2 \neq 0$$

$$k \in \mathbb{R} \quad k \neq 0$$

Now, let $k = y$ (to test the x/z plane shape, the vertical tracing)

$$\frac{x^2}{4} + \frac{k^2}{16} - z = 0$$

$$\frac{x^2}{4} - z = -\frac{k^2}{16} \text{ This is an equation for a } \boxed{\text{Parabola}}$$

$$0 \neq -\frac{k^2}{16}$$

$$0 \neq k^2$$

$$k \in \mathbb{R} \quad k \neq 0$$

Since the equation $\frac{x^2}{4} + \frac{y^2}{16} - z = 0$ can be
rewritten as $\frac{x^2}{4} + \frac{y^2}{16} = \frac{z}{1}$ the equation represents an

Elliptic Paraboloid where $c = \pm 1$, $a = \pm 2$, and $b = \pm 4$
and matches with **Figure I**

For equation (c): $\frac{x^2}{4} + \frac{y^2}{16} - \frac{z^2}{4} = 1$

Let $k = z$ (to test the x/y plane shape, the horizontal tracing)

$$\frac{x^2}{4} + \frac{y^2}{16} - \frac{k^2}{4} = 1$$

$$\frac{x^2}{4} + \frac{y^2}{16} = 1 + \frac{k^2}{4} \text{ is an equation for an } \boxed{\text{Ellipse}}$$

$$1 + \frac{k^2}{4} \neq 0$$

$\frac{k^2}{4} \neq -1$ has no solutions in the real number spectrum

Meaning $k \in \mathbb{R}$

Now, let $k = x$ (to test the y/z plane shape, the vertical tracing)

$$\frac{k^2}{4} + \frac{y^2}{16} - \frac{z^2}{4} = 1$$

$$\frac{y^2}{16} - \frac{z^2}{4} = 1 - \frac{k^2}{4} \text{ is an equation for a } \boxed{\text{Hyperbola}}$$

$$1 - \frac{k^2}{4} \neq 0$$

$$1 \neq \frac{k^2}{4}$$

$$4 \neq k^2$$

$$k \in \mathbb{R} \quad k \neq \pm 2$$

Now, let $k = y$ (to test the x/z plane shape, the vertical tracing)

$$\frac{x^2}{4} + \frac{k^2}{16} - \frac{z^2}{4} = 1$$

$$\frac{x^2}{4} - \frac{z^2}{4} = 1 - \frac{k^2}{16} \text{ is an equation for a } \boxed{\text{Hyperbola}}$$

$$1 - \frac{k^2}{16} \neq 0$$

$$1 \neq \frac{k^2}{16}$$

$$16 \neq k^2$$

$$k \in \mathbb{R} \quad k \neq \pm 4$$

The equation $\frac{x^2}{4} + \frac{y^2}{16} - \frac{z^2}{4} = 1$ is an equation for a **Hyperboloid of One Sheet** where

$a = c = \sqrt{4} = 2$ and $b = \sqrt{16} = 4$, matching

Figure II

For equation (d): $x^2 - 4y^2 - z^2 = 4$

Let $k = z$ (to test the x/y plane shape, the horizontal tracing)

$$x^2 - 4y^2 - k^2 = 4$$

$$x^2 - 4y^2 = 4 + k^2 \text{ is an equation of a } \boxed{\text{Hyperbola}}$$

$$4 + k^2 \neq 0$$

$$k^2 \neq -4 \text{ is true for all real numbers}$$

$$k \in \mathbb{R}$$

Now, let $k = x$ (to test the y/z plane shape, the vertical tracing)

$$k^2 - 4y^2 - z^2 = 4$$

$$-4y^2 - z^2 = 4 - k^2$$

$$4y^2 + z^2 = k^2 - 4 \text{ is an equation for an } \boxed{\text{Ellipse}}$$

$$k^2 - 4 \neq 0$$

$$k^2 \neq 4$$

$$k \in \mathbb{R} \quad k \neq \pm 2$$

Now, let $k = y$ (to test the x/z plane shape, the vertical tracing)

$$x^2 - 4k^2 - z^2 = 4$$

$$x^2 - z^2 = 4 + 4k^2 \text{ is an equation of a } \boxed{\text{Hyperbola}}$$

$$4 + 4k^2 \neq 0$$

$$4k^2 \neq -4$$

$$k^2 \neq -1 \text{ is true for all real numbers}$$

$$k \in \mathbb{R}$$

The equation $x^2 - 4y^2 - z^2 = 4$ can be rewritten as $\frac{x^2}{4} - y^2 - \frac{z^2}{4} = 1$ making the quadric a

Hyperboloid of Two Sheets. The x and z variables are both negative, making the graph center go through the x axis, corresponding to **Figure IV**

For equation (e): $9y^2 - 4x^2 - 9z = 0$

Let $k = z$ (to test the x/y plane shape, the horizontal tracing)

$$9y^2 - 4x^2 - 9k = 0$$

$9y^2 - 4x^2 = 9k$ is an equation of a Hyperbola

$$9k \neq 0$$

$$k \in \mathbb{R} \quad k \neq 0$$

Now, let $k = x$ (to test the y/z plane shape, the vertical tracing)

$$9y^2 - 4k^2 - 9z = 0$$

$9y^2 - 9z = 4k^2$ is an equation of a Parabola

$$4k^2 \neq 0$$

$$k \in \mathbb{R} \quad k \neq 0$$

Now, let $k = y$ (to test the x/z plane shape, the vertical tracing)

$$9k^2 - 4x^2 - 9z = 0$$

$-4x^2 - 9z = -9k^2$ is an equation of a Parabola

$$-9k^2 \neq 0$$

$$k^2 \neq 0$$

$$k \in \mathbb{R} \quad k \neq 0$$

The equation $9y^2 - 4x^2 - 9z = 0$ carries the format of a **Hyperbolic Paraboloid**, represented in **Figure V**, where the z axis value is the non-exponential and the y variable is the positive squared variable

$$2 \quad \sum_{n=1}^{\infty} (-1)^n \frac{(x+5)^n}{n \cdot 3^n}$$

a) Finding the radius of convergence

Method: Ratio Test $R = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ where the series converges if $|R| < 1$

and where a_n denotes the n_{th} term of the series and r is the radius of convergence

$$a_n = (-1)^n \frac{(x+5)^n}{n \cdot 3^n}$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} \frac{(x+5)^{n+1}}{(n+1) \cdot 3^{n+1}}}{(-1)^n \frac{(x+5)^n}{n \cdot 3^n}} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{(x+5)^{n+1}}{(n+1) \cdot 3^{n+1}} \cdot \frac{n \cdot 3^n}{(x+5)^n} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{n \cdot (x+5)}{3 \cdot (n+1)} \right|$$

$$R = |x+5| \lim_{n \rightarrow \infty} \left| \frac{n}{3 \cdot (n+1)} \right|$$

$$R = |x+5| \cdot \left| \frac{1}{3} \right|$$

$$\left| \frac{x+5}{3} \right| = 1$$

$$|x+5| = 3$$

$$\boxed{r = 3}$$

The radius of convergence is 3

and the interval of convergence is $-8 < x \leq -2$

b) Finding the Interval of Convergence

The series converges if $\left| \frac{x+5}{3} \right| < 1$

$$-1 < \frac{x+5}{3} < 1$$

$$-3 < x+5 < 3$$

$$-8 < x < -2$$

Testing the endpoints:

Lower: $\sum_{n=1}^{\infty} (-1)^n \frac{(-8+5)^n}{n \cdot 3^n}$

$$\sum_{n=1}^{\infty} (-1)^n \frac{(-3)^n}{n \cdot 3^n}$$

The $\frac{(-3)^n}{3^n}$ can be reduced to $(-1)^n$

$$\sum_{n=1}^{\infty} (-1)^n \frac{(-1)^n}{n}$$

The $(-1)^n(-1)^n$ can be reduced to $(-1)^{2n}$, which can be reduced to 1^n , which is equal to 1 for all real numbers

$\sum_{n=1}^{\infty} \frac{1}{n}$ diverges by the Harmonic Series Test, meaning -8 can not be included in the interval of convergence

Upper: $\sum_{n=1}^{\infty} (-1)^n \frac{(-2+5)^n}{n \cdot 3^n}$

$$\sum_{n=1}^{\infty} (-1)^n \frac{3^n}{n \cdot 3^n}$$

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{ is } \boxed{\text{convergent by the Alternating Series Test}},$$

meaning that -2 is part of the interval of convergence

After testing endpoints, we can conclude the interval of convergence is $\boxed{-8 < x \leq -2}$

$$\mathbf{3} \quad f(x) = \frac{x^2}{3x+4}$$

Conversion to a power series

To be easily converted to a power series $\sum_{n=0}^{\infty} ar^n$, the equation will need the form of $\frac{a}{1-r}$

We can first pull out an x^2 from the fraction and get $f(x) = x^2 \frac{1}{3x+4}$

Separating out a 4 from the denominator get $f(x) = x^2 \cdot \frac{1}{4} \cdot \frac{1}{\frac{3x}{4} + 1}$

Which can be rearranged as $f(x) = \frac{x^2}{4} \cdot \frac{1}{1 + \frac{3x}{4}}$

When $a = 1$, $r = -\frac{3x}{4}$, the power series can be represented as $f(x) = \frac{x^2}{4} \sum_{n=0}^{\infty} \left(-\frac{3x}{4}\right)^n$

$$f(x) = \frac{x^2}{4} \sum_{n=0}^{\infty} \left(-\frac{3x}{4}\right)^n$$

Distributing the power $f(x) = \frac{x^2}{4} \sum_{n=0}^{\infty} \frac{(-1)^n \cdot 3^n \cdot x^n}{4^n}$

Since $x^2 \cdot x^n = x^{n+2}$ and $4 \cdot 4^n = 4^{n+1}$, substituting them in we get $f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot 3^n \cdot x^{n+2}}{4^{n+1}}$

Similar to section 11.9 example 3, we can shift the indices $f(x) = \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 3^{n-2} \cdot x^n}{4^{n-1}}$

Plugging this back in,
$$\frac{x^2}{3x+4} = \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 3^{n-2} \cdot x^n}{4^{n-1}}$$

Finding radius of convergence

Method: Ratio Test $R = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ where the series converges if $|R| < 1$

and where a_n denotes the n_{th} term of the series and r is the radius of convergence

$$a_n = \frac{(-1)^{n-2} \cdot 3^{n-2} \cdot x^n}{4^{n-1}}$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n-1} \cdot 3^{n-1} \cdot x^{n+1}}{4^n}}{\frac{(-1)^{n-2} \cdot 3^{n-2} \cdot x^n}{4^{n-1}}} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n-1} \cdot 3^{n-1} \cdot x^{n+1}}{4^n} \cdot \frac{4^{n-1}}{(-1)^{n-2} \cdot 3^{n-2} \cdot x^n} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{-3x}{4} \right|$$

$$R = \left| \frac{-3x}{4} \right|$$

$$\left| \frac{-3x}{4} \right| = 1$$

$$|-3x| = 4$$

$$3x = 4$$

$$\boxed{r = \frac{4}{3}}$$

Finding interval of convergence

The series converges if $\left| \frac{-3x}{4} \right| < 1$

$$-1 < \frac{3x}{4} < 1$$

$$-4 < 3x < 4$$

$$-\frac{4}{3} < x < \frac{4}{3}$$

Testing the endpoints:

Upper Endpoint: $x = \frac{4}{3}$

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 3^{n-2} \cdot \frac{4^n}{3}}{4^{n-1}} \\ &= \sum_{n=2}^{\infty} \frac{(-3)^{n-2} \cdot \frac{4^n}{3}}{4^{n-1}} \\ &= \sum_{n=2}^{\infty} \frac{(-3)^{n-2} \cdot \frac{4^{n-2}}{3} \cdot \frac{4^2}{3}}{4^{n-2} \cdot 4} \\ &= \sum_{n=2}^{\infty} \frac{(-4)^{n-2} \cdot \frac{4^2}{3}}{4^{n-2} \cdot 4} \\ &= \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 16}{4 \cdot 9} \\ &= \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 4}{9} \end{aligned}$$

This series is divergent by the Ratio Test, where the ratio is 1, meaning the upper endpoint can not be included in the interval of convergence

Lower Endpoint: $x = -\frac{4}{3}$

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{(-1)^{n-2} \cdot 3^{n-2} \cdot -\frac{4^n}{3}}{4^{n-1}} \\ &= \sum_{n=2}^{\infty} \frac{(-3)^{n-2} \cdot -\frac{4^n}{3}}{4^{n-1}} \\ &= \sum_{n=2}^{\infty} \frac{(-3)^{n-2} \cdot -\frac{4^{n-2}}{3} \cdot -\frac{4^2}{3}}{4^{n-2} \cdot 4} \\ &= \sum_{n=2}^{\infty} \frac{4^{n-2} \cdot \frac{4^2}{3}}{4^{n-2} \cdot 4} \\ &= \sum_{n=2}^{\infty} \frac{\left(-\frac{4}{3}\right)^2}{4} \\ &= \sum_{n=2}^{\infty} \frac{16}{4 \cdot 9} \\ &= \sum_{n=2}^{\infty} \frac{4}{9} \end{aligned}$$

This series is divergent by the Ratio Test, where the ratio is 1, meaning the lower endpoint can not be included in the interval of convergence

$$\boxed{\text{Radius of Convergence: } \frac{4}{3}}$$

$$\boxed{\text{Interval of Convergence: } -\frac{4}{3} < x < \frac{4}{3}}$$

$$4 \quad f(x) = 3^{-2x}$$

Finding the first 6 coefficients $\frac{f^{(n)}(0)}{n!}$

$$0: f(x) = 3^{-2x} = 9^{-x} = \frac{1}{9^x} \text{ at } x = 0$$

$$\frac{1}{9^0} = 1 \text{ then dividing by } 0! \quad \frac{1}{0!} = \boxed{1}$$

$$1: f'(x) = \frac{d}{dx} 3^{-2x} = \frac{d}{dx} 3^{-2x} = \frac{d}{dx} \frac{1}{3^{2x}}$$

$$\text{using the quotient rule } \frac{d}{dx} \frac{1}{3^{2x}} = \frac{0 \cdot 3^{2x} - 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-2 \ln 3}{3^{2x}} \text{ where } x = 0$$

$$= \frac{-2 \ln 3}{3^{2 \cdot 0}} = -2 \ln 3$$

$$\text{dividing by } 1! \text{ we get } \boxed{\frac{-2 \ln 3}{1!}}$$

$$2: f''(x) = \frac{d}{dx} \frac{-2 \ln 3}{3^{2x}}$$

Using the quotient rule, again

$$\frac{d}{dx} \frac{-2 \ln 3}{3^{2x}} = \frac{0 \cdot 3^{2x} - (-2 \ln 3) \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{2 \ln 3 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{(2 \ln 3)(2 \ln 3)}{3^{2x}} \text{ where } x = 0$$

$$= (2 \ln 3)(2 \ln 3) \text{ or } (2 \ln 3)^2$$

$$\text{Dividing by } 2! \text{ we get } \boxed{\frac{(2 \ln 3)^2}{2!}}$$

$$3: f'''(x) = \frac{d}{dx} \frac{(2 \ln 3)^2}{3^{2x}}$$

Using the quotient rule, again

$$\frac{d}{dx} \frac{(2 \ln 3)^2}{3^{2x}} = \frac{0 \cdot 3^{2x} - (2 \ln 3)^2 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-(2 \ln 3)^2 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-(2 \ln 3)^3}{3^{2x}} \text{ where } x = 0$$

$$= -(2 \ln 3)^3$$

$$\text{dividing by } 3! \text{ we get } \boxed{\frac{-(2 \ln 3)^3}{3!}}$$

$$4: f^{(4)}(x) = \frac{d}{dx} \frac{-(2 \ln 3)^3}{3^{2x}}$$

$$\text{Using the quotient rule, again } \frac{d}{dx} \frac{-(2 \ln 3)^3}{3^{2x}} = \frac{0 \cdot 3^{2x} - (-(2 \ln 3)^3) \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{(2 \ln 3)^3 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{(2 \ln 3)^4}{3^{2x}} \text{ where } x = 0, \text{ evaluates to } (2 \ln 3)^4$$

$$\text{Dividing by } 4! \text{ we get } \boxed{\frac{(2 \ln 3)^4}{4!}}$$

$$5: f^{(5)}(x) = \frac{d}{dx} \frac{(2 \ln 3)^4}{3^{2x}}$$

Using the quotient rule, for the last time

$$\frac{d}{dx} \frac{(2 \ln 3)^4}{3^{2x}} = \frac{0 \cdot 3^{2x} - (2 \ln 3)^4 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{0 \cdot 3^{2x} - (2 \ln 3)^4 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-(2 \ln 3)^4 \cdot 3^{2x}(2 \ln 3)}{3^{4x}}$$

$$= \frac{-(2 \ln 3)^5 \cdot 3^{2x}}{3^{4x}}$$

$$= \frac{-(2 \ln 3)^5}{3^{2x}} \text{ where } x = 0, \text{ evaluates to } -(2 \ln 3)^5$$

$$\text{Dividing by } 5! \text{ we get } \boxed{\frac{-(2 \ln 3)^5}{5!}}$$

First 6 coefficients:

0. 1

1. $\frac{-2 \ln 3}{1!}$

2. $\frac{(2 \ln(3))^2}{2!}$

3. $\frac{-(2 \ln(3))^3}{3!}$

4. $\frac{(2 \ln(3))^4}{4!}$

5. $\frac{-(2 \ln(3))^5}{5!}$

The emergent pattern seems to be $\boxed{\frac{f^{(n)}(0)}{n!} = (-1)^n \frac{(2 \ln(3))^n}{n!}}$

The Maclaurin Series

Using the known coefficient pattern, we can multiply the general term by x^n and make it a summation

$$\boxed{\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = \sum_{n=0}^{\infty} (-1)^n \frac{(2 \ln(3))^n}{n!} x^n}$$

5 Line L passes through $A(4, -3, 5)$ and $B(-2, -1, 8)$

Vector Equation

For the purposes of this problem, I will use the vector going in the direction from point A to point B

We can create a vector \vec{v} that is parallel to \overrightarrow{AB}

The variable t will represent a scalar parameter to any equation in the scope of this problem that takes in a parameter as an input

This vector \vec{v} can be found using the formula $B - A$ and is calculated using $\vec{v} = \langle B_x - A_x, B_y - A_y, B_z - A_z \rangle$
 $\vec{v} = \langle -2 - 4, -1 - (-3), 8 - 5 \rangle = \langle -6, 2, 3 \rangle$

Now using $\vec{r} = \vec{r}_0 + t\vec{v}$, where r_0 is the tail point of the vector, denoted by point $A(4, -3, 5)$ for this example, we can compute the tip of the vector given a parameter and the following vector equation:

$$\vec{r} = \langle 4, -3, 5 \rangle + t\langle -6, 2, 3 \rangle$$

$$\vec{r} = \langle 4, -3, 5 \rangle + \langle -6t, 2t, 3t \rangle$$

$$\boxed{\vec{r} = \langle 4 - 6t, 2t - 3, 3t + 5 \rangle}$$

Parametric Equations

Using Theorem #2 in section 12.5 of the textbook, Parametric equations for a line through the point (x_0, y_0, z_0) and parallel to the direction vector $\langle a, b, c \rangle$ are:

$$x = x_0 + at \quad y = y_0 + bt \quad z = z_0 + ct$$

(x_0, y_0, z_0) will be the tail of the vector, point $A(4, -3, 5)$, and $\langle a, b, c \rangle$ will be the respective components of defined vector \vec{v} ,

$$\vec{v} = \langle -6, 2, 3 \rangle$$

By Theorem #2 in section 12.5, $\boxed{x = 4 - 6t \quad y = -3 + 2t \quad z = 5 + 3t}$

Symmetric Equation

To find a symmetric equation, we can solve all 3 of our parametric equations for the variable t , assuming none of the components of $\langle a, b, c \rangle$ are 0, which would introduce division by zero

$$x = 4 - 6t \quad y = -3 + 2t \quad z = 5 + 3t$$

$$x - 4 = -6t \quad y + 3 = 2t \quad z - 5 = 3t$$

$$\boxed{\frac{x - 4}{-6} = t}$$

$$\boxed{\frac{y + 3}{2} = t}$$

$$\boxed{\frac{z - 5}{3} = t}$$

Combining these into a single equation to eliminate the parameter t , we get the Symmetric Equation

$$\boxed{\frac{x - 4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}}$$

5.1 Intersecting the planes

The line $\frac{x - 4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}$ intersects the xy plane when $z = 0$

$$\frac{x - 4}{-6} = \frac{y + 3}{2} = \frac{-5}{3}$$

$$\frac{x - 4}{-6} = \frac{-5}{3}$$

$$x - 4 = \frac{-5 \cdot -6}{3}$$

$$x - 4 = 10$$

$$\boxed{x = 14}$$

$$\frac{y + 3}{2} = \frac{-5}{3}$$

$$y + 3 = \frac{-5 \cdot 2}{3}$$

$$y + 3 = \frac{-10}{3}$$

$$y = \frac{-10}{3} - 3$$

$$\boxed{y = \frac{-19}{3}}$$

Making the xy plane intersection coordinates $\boxed{(14, \frac{-19}{3}, 0)}$

The line $\frac{x - 4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}$ intersects the xz plane when $y = 0$

$$\frac{x - 4}{-6} = \frac{3}{2} = \frac{z - 5}{3}$$

$$\frac{x - 4}{-6} = \frac{3}{2}$$

$$x - 4 = \frac{-6 \cdot 3}{2}$$

$$x - 4 = \frac{-18}{2}$$

$$x - 4 = -9$$

$$\boxed{x = -5}$$

$$\frac{3}{2} = \frac{z - 5}{3}$$

$$\frac{3 \cdot 3}{2} = z - 5$$

$$\frac{9}{2} = z - 5$$

$$\boxed{z = \frac{19}{2}}$$

Making the intersection point to the xz plane $\boxed{(-5, 0, \frac{19}{2})}$

The line $\frac{x - 4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}$ intersects the yz plane when $x = 0$

$$\frac{-4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}$$

$$\frac{-4}{-6} = \frac{y + 3}{2} = \frac{z - 5}{3}$$

$$\frac{4}{6} = \frac{y + 3}{2} = \frac{z - 5}{3}$$

$$\frac{4}{6} = \frac{y+3}{2}$$

$$\frac{8}{6} = y+3$$

$$\frac{8}{6} - 3 = y$$

$$y = \frac{-10}{6}$$

$$\boxed{y = \frac{-5}{3}}$$

$$\frac{-4}{-6} = \frac{z-5}{3}$$

$$\frac{-4 \cdot 3}{-6} = z-5$$

$$\frac{-12}{-6} = z-5$$

$$2 = z-5$$

$$\boxed{z = 7}$$

Making the intersection point to the yz plane $\boxed{(0, \frac{-5}{3}, 7)}$

5.2 Distance to Point $C(-1, 4, -6)$

This can be calculated using $d = \frac{|\vec{v} \times \vec{BC}|}{|\vec{v}|}$ where

$$\vec{v} = \langle -6, 2, 3 \rangle \text{ and } \vec{BC} = C - B = \langle -1 - (-2), 4 - (-1), -6 - 8 \rangle = \langle 1, 5, -14 \rangle \quad \vec{v} \times \vec{BC}$$

$$\text{Starting off with the matrix } \vec{v} \times \vec{BC} = \begin{vmatrix} i & j & k \\ -6 & 2 & 3 \\ 1 & 5 & -14 \end{vmatrix}$$

$$\vec{v} \times \vec{BC} = i(2 \cdot -14 - 3 \cdot 5) - j(-6 \cdot -14 - 3 \cdot 1) + k(-6 \cdot 5 - 2 \cdot 1)$$

$$\vec{v} \times \vec{BC} = i(-28 - 15) - j(84 - 3) + k(-30 - 2)$$

$$\vec{v} \times \vec{BC} = i(-43) - j(81) + k(-32)$$

$$\vec{v} \times \vec{BC} = \langle -43, -81, -32 \rangle$$

Finding the magnitude is easy using the Pythagorean theorem for 3 dimensions

$$|\vec{v} \times \vec{BC}| = \sqrt{(-43)^2 + (-81)^2 + (-32)^2}$$

$$|\vec{v} \times \vec{BC}| = \sqrt{1849 + 6561 + 1024}$$

$$|\vec{v} \times \vec{BC}| = \boxed{\sqrt{9434}}$$

Now to divide it by the magnitude of \vec{v}

$$|\vec{v}| = \sqrt{(-6)^2 + 2^2 + 3^2}$$

$$|\vec{v}| = \sqrt{36 + 4 + 9}$$

$$|\vec{v}| = \sqrt{49}$$

$$|\vec{v}| = 7$$

$$\boxed{d = \frac{\sqrt{9434}}{7}} \text{ units}$$

6 Maclaurin Series for $\tan^{-1} x = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{2n+1}}{2n+1}$

Power Series for $f(x) = x^3 \cdot \tan^{-1}(x^2)$

Given that we know the Maclaurin Series for $\tan^{-1}(x)$, modifying the power series for our new equation is straightforward

Modifying the x As x is not the index variable in the infinite series, it is fully permeable through the membrane of the sigma function. This will be accomplished by directly substituting the variable, similar to what was shown in example 11 in Section 11.10 in the textbook

$$\tan^{-1}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{2n+1}}{2n+1}$$

$$x^3 \cdot \tan^{-1}(x^2) = x^3 \sum_{n=0}^{\infty} \frac{(-1)^n \cdot (x^2)^{2n+1}}{2n+1}$$

Multiplying together the stacked exponents: $x^3 \cdot \tan^{-1}(x^2) = x^3 \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+2}}{2n+1}$ as $2 \cdot 2n + 1 = 4n + 2$

Moving the x^3 across the sigma function membrane, we get $x^3 \cdot \tan^{-1}(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+2} \cdot x^3}{2n+1}$

Simplifies to $x^3 \cdot \tan^{-1}(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+2+3}}{2n+1}$

$$x^3 \cdot \tan^{-1}(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+2+3}}{2n+1}$$

$$x^3 \cdot \tan^{-1}(x^2) = \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+5}}{2n+1}$$

Checking for Convergence in the interval of integration

Using the Ratio test

Using $a_n = \frac{(-1)^n \cdot x^{4n+5}}{2n+1}$ we can do the ratio test $R = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$ and the series converges for $R < 1$

$$R = \lim_{n \rightarrow \infty} \left| \frac{\frac{(-1)^{n+1} \cdot x^{4(n+1)+5}}{2(n+1)+1}}{\frac{(-1)^n \cdot x^{4n+5}}{2n+1}} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} \cdot x^{4n+9}}{2(n+1)+1} \cdot \frac{2n+1}{(-1)^n \cdot x^{4n+5}} \right|$$

$$R = \lim_{n \rightarrow \infty} \left| \frac{-x \cdot (2n+1)}{2n+3} \right|$$

$$R = |-x| \lim_{n \rightarrow \infty} \left| \frac{(2n+1)}{2n+3} \right|$$

$$R = |-x|$$

$$-1 < -x < 1$$

$$1 > x > -1$$

Since the series is for sure convergent for all values between, but not including $x = -1$ and $x = 1$, we know it is convergent for the interval $[0, \frac{1}{2}]$

Integrating $\int_0^{\frac{1}{2}} x^3 \cdot \tan^{-1}(x^2) dx$

Given the power series for $x^3 \cdot \tan^{-1}(x^2)$ as $\sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+5}}{2n+1}$

We can start off with $\int_0^{\frac{1}{2}} x^3 \cdot \tan^{-1}(x^2) dx = \int_0^{\frac{1}{2}} \sum_{n=0}^{\infty} \frac{(-1)^n \cdot x^{4n+5}}{2n+1} dx$

We can move everything not bound to the variable x out of the integrand, including the sigma function

$$= \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \int_0^{\frac{1}{2}} x^{4n+5} dx$$

Evaluating the integral using the power rule

$$\begin{aligned} &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \int_0^{\frac{1}{2}} x^{4n+5} dx \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \frac{x^{4n+6}}{4n+6} \Big|_{x=0}^{x=\frac{1}{2}} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \frac{(\frac{1}{2})^{4n+6}}{4n+6} - \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \frac{0^{4n+6}}{4n+6} \rightarrow 0 \\ &= \boxed{\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} \frac{(\frac{1}{2})^{4n+6}}{4n+6}} \end{aligned}$$

Using a calculator, we can calculate the values of each term to find where $|\text{Error}| < 1.0 \cdot 10^{-9}$ or $\frac{1}{1,000,000,000}$

$$0. \frac{1}{384} = 0.0026041666$$

$$1. \frac{-1}{30720} = -0.000032552083$$

$$2. \frac{1}{1146880} = 8.7193080357 \cdot 10^{-7}$$

$$3. \frac{-1}{33030144} = -3.0275375124 \cdot 10^{-8}$$

$$4. \frac{1}{830472192} = 1.204134237886 \cdot 10^{-9}$$

$$5. \frac{-1}{19193135104} = -5.210196221625 \cdot 10^{-11} - \text{This term is the error as it is the first term below } 10^{-9}$$

Giving us the series: $\boxed{\frac{1}{384} - \frac{1}{30720} + \frac{1}{1146880} - \frac{1}{33030144} + \frac{1}{830472192}}$ Or: $\boxed{\sum_{n=0}^4 \frac{(-1)^n}{2n+1} \frac{(\frac{1}{2})^{4n+6}}{4n+6}}$

The output from this summation is $\boxed{0.00257246}$

$$7 \quad A(2, 6, 5), B(-1, 3, 4), C(-1, -5, 7)$$

7.1 Side Lengths

$$|\vec{AB}|$$

$$\vec{AB} = B - A$$

$$= \langle -1 - 2, 3 - 6, 4 - 5 \rangle$$

$$= \langle -3, -3, -1 \rangle$$

$$\text{Now for magnitude} = \sqrt{(-3)^2 + (-3)^2 + (-1)^2} = \boxed{\sqrt{19}}$$

$$|\vec{AC}|$$

$$\vec{AC} = C - A$$

$$= \langle -1 - 2, -5 - 6, 7 - 5 \rangle$$

$$= \langle -3, -11, 2 \rangle$$

$$\text{Now for magnitude} = \sqrt{(-3)^2 + (-11)^2 + (2)^2} = \sqrt{9 + 121 + 4} = \boxed{\sqrt{134}}$$

$$|\vec{BC}|$$

$$\vec{BC} = C - B$$

$$= \langle -1 - (-1), -5 - 3, 7 - 4 \rangle$$

$$= \langle 0, -8, 3 \rangle$$

$$\text{Now for magnitude} = \sqrt{(0)^2 + (-8)^2 + (3)^2} = \sqrt{0 + 64 + 9} = \boxed{\sqrt{73}}$$

7.2 Angles

$$\text{Theorem: } |A \times B| = |A||B|\sin(\theta)$$

$$|\vec{AB} \times \vec{BC}| = |\vec{AB}||\vec{BC}|\sin(\theta)$$

$$\vec{AB} \times \vec{BC} = \begin{vmatrix} i & j & k \\ -3 & -3 & -1 \\ 0 & -8 & 3 \end{vmatrix}$$

$$\vec{AB} \times \vec{BC} = i(-3 \cdot 3 - (-1) \cdot -8) - j(-3 \cdot 3 - (-1) \cdot 0) + k(-3 \cdot -8 - 3 \cdot 0)$$

$$\vec{AB} \times \vec{BC} = i(-17) - j(-9) + k(24)$$

$$\vec{AB} \times \vec{BC} = \langle -17, 9, 24 \rangle$$

$$|\vec{AB} \times \vec{BC}| = \sqrt{(-17)^2 + 9^2 + (24)^2}$$

$$|\vec{AB} \times \vec{BC}| = \sqrt{(-17)^2 + 9^2 + (24)^2}$$

$$|\vec{AB} \times \vec{BC}| = \sqrt{946}$$

$$\sqrt{946} = \sqrt{19}\sqrt{73}\sin\theta$$

$$\frac{\sqrt{946}}{\sqrt{19}\sqrt{73}} = \sin\theta$$

$$\arcsin \frac{\sqrt{946}}{\sqrt{19}\sqrt{73}} = \theta$$

$$55.68^\circ = \theta$$

However, since that input exceeds the maximum possible angle that arcsin can provide (90°), we can subtract the output from 180° to get the final angle. This is true since $\sin(x)$ is horizontally symmetric around the points $x = \frac{\pi}{2} + n\pi$

$$180^\circ - 55.68^\circ = \boxed{\theta = 124.32^\circ} \text{ for angle } \angle ABC$$

$$|\vec{AC} \times \vec{BC}| = |\vec{AC}||\vec{BC}|\sin(\theta)$$

$$\vec{AC} \times \vec{BC} = \begin{vmatrix} i & j & k \\ -3 & -11 & 2 \\ 0 & -8 & 3 \end{vmatrix}$$

$$\vec{AC} \times \vec{BC} = i(-11 \cdot 3 - 2 \cdot -8) - j(-3 \cdot 3 - 2 \cdot 0) + k(-3 \cdot -8 - 11 \cdot 0)$$

$$\vec{AC} \times \vec{BC} = i(-17) - j(-9) + k(24)$$

$$\vec{AC} \times \vec{BC} = \langle -17, 9, 24 \rangle$$

$$|\vec{AC} \times \vec{BC}| = \sqrt{(-17)^2 + (9)^2 + (24)^2}$$

$$|\vec{AC} \times \vec{BC}| = \sqrt{946}$$

$$\sqrt{946} = \sqrt{134}\sqrt{73}\sin\theta$$

$$\frac{\sqrt{946}}{\sqrt{134}\sqrt{73}} = \sin\theta$$

$$\arcsin \frac{\sqrt{946}}{\sqrt{134}\sqrt{73}} = \theta$$

$$\boxed{\theta = 18.12^\circ} \text{ for angle } \angle ACB$$

$$|\vec{AB} \times \vec{AC}| = |\vec{AB}||\vec{AC}|\sin(\theta)$$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} i & j & k \\ -3 & -3 & -1 \\ -3 & -11 & 2 \end{vmatrix}$$

$$\vec{AB} \times \vec{AC} = i(-3 \cdot 2 - (-1) \cdot -11) - j(-3 \cdot 2 - (-1) \cdot -3) + k(-3 \cdot -11 - (-3) \cdot -3)$$

$$\vec{AB} \times \vec{AC} = i(-6 - 11) - j(-6 - 3) + k(33 - 9)$$

$$\vec{AB} \times \vec{AC} = i(-17) - j(-9) + k(24)$$

$$\vec{AB} \times \vec{AC} = \langle -17, 9, 24 \rangle$$

$$|\vec{AB} \times \vec{AC}| = \sqrt{(-17)^2 + (9)^2 + 24^2}$$

$$|\vec{AB} \times \vec{AC}| = \sqrt{946}$$

$$\sqrt{946} = \sqrt{19}\sqrt{134}\sin\theta$$

$$\frac{\sqrt{946}}{\sqrt{19}\sqrt{134}} = \sin\theta$$

$$\arcsin \frac{\sqrt{946}}{\sqrt{19}\sqrt{134}} = \theta$$

$$\boxed{\theta = 37.56^\circ}$$

7.3 Linear Equation of Plane Containing points A,B,C

Using equation $a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$ where a, b, c are components of the normal vector \vec{n} and (x_0, y_0, z_0) are coordinates for a reference points, in this case point $A(2, 6, 5)$

$$\vec{AB} \times \vec{AC} = \begin{vmatrix} i & j & k \\ -3 & -3 & -1 \\ -3 & -11 & 2 \end{vmatrix}$$

$$\vec{AB} \times \vec{AC} = i(-3 \cdot 2 - (-1) \cdot -11) - j(-3 \cdot 2 - (-1) \cdot -3) + k(-3 \cdot -11 - (-3) \cdot -3)$$

$$\vec{AB} \times \vec{AC} = i(-6 - 11) - j(-6 - 3) + k(33 - 9)$$

$$\vec{AB} \times \vec{AC} = i(-17) - j(-9) + k(24)$$

$$\vec{AB} \times \vec{AC} = \langle -17, -9, 24 \rangle$$

$$\text{Let } \vec{n} = \langle a, b, c \rangle = \langle -17, -9, 24 \rangle$$

Plugging things in, we get $-17(x - 2) + 9(y - 6) + 24(z - 5) = 0$

$$-17x + 34 + 9y - 54 + 24z - 120 = 0$$

$$\boxed{-17x + 9y + 24z = 140}$$

7.4 Coplanarity with $D(-2, 6, -4)$

To see if point $D(-2, 6, -4)$ lies on the plane $-17(x - 2) + 9(y - 6) + 24(z - 5) = 0$, we can plug in the coordinates

$$-17(-2 - 2) + 9(6 - 6) + 24(-4 - 5) = 0$$

$$-17(-4) + 9(0) + 24(-9) = 0$$

$$68 - 216 = 0$$

$$-148 = 0 \text{ This is obviously not true, meaning point } \boxed{D \text{ is not coplanar with points } A, B, C}$$

7.5 Volume of the Parallelepiped

Volume can be found using the formula provided by the textbook $V = |\vec{a} \cdot (\vec{b} \times \vec{c})|$

Lets refresh on our points and vectors:

Points: $A(2, 6, 5), B(-1, 3, 4), C(-1, -5, 7), D(-2, 6, -4)$

Vectors: $\vec{AB} = \langle -3, -3, -1 \rangle, \vec{AC} = \langle -3, -11, 2 \rangle, \vec{BC} = \langle 0, -8, 3 \rangle$

Calculating vector \vec{AD}

$$\vec{AD} = D - A$$

$$\vec{AD} = \langle -2 - 2, 6 - 6, -4 - 5 \rangle$$

$$\vec{AD} = \langle -4, 0, -9 \rangle$$

Now we can use the Volume Formula using $\vec{a} = \vec{AB}, \vec{b} = \vec{AC}, \vec{c} = \vec{AD}$

Calculating $\vec{b} \times \vec{c}$

$$\vec{b} \times \vec{c} = \vec{AC} \times \vec{AD}$$

$$\vec{AC} \times \vec{AD} = \begin{vmatrix} i & j & k \\ -3 & -11 & 2 \\ -4 & 0 & -9 \end{vmatrix}$$

$$\vec{AC} \times \vec{AD} = i(-11 \cdot -9 - 2 \cdot 0) - j(-3 \cdot -9 - (-4) \cdot 2) + k(-3 \cdot 0 - (-11) \cdot -4)$$

$$\vec{AC} \times \vec{AD} = i(99) - j(35) + k(-44)$$

$$\vec{AC} \times \vec{AD} = \langle 99, -35, -44 \rangle$$

$$\vec{b} \times \vec{c} = \langle 99, -35, -44 \rangle$$

Calculating the dot product with $\vec{a} \cdot \vec{b} \times \vec{c}$

$$\langle -3, -3, -1 \rangle \cdot \langle 99, -35, -44 \rangle$$

$$= -3 \times 99 + -3 \times -35 + -1 \times -44$$

$$= -297 + 105 + 44$$

$$= -148$$

To top it all off, take the absolute value of the result $V = |-148| = \boxed{148 \text{ units}^3}$