

MATH 2 Lecture Notes

Tejas Patel

Tuesday, 14 January, 2025

Contents

1	Chapter 1	2
1.1	Terminology	2
1.2	Some Mathematical Models	2
2	First-Order Differential Equations	4
2.1	Preliminary Theory	4
2.2	Separable Variables	4
2.3	Homogenous Equations	4
2.4	Exact Equations	4
2.5	Preliminary Theory	4
3	Example Problems with Solutions	5
3.1	5

1 Chapter 1

1.1 Terminology

Definition A differential equation is an equation containing the derivatives or differentials of one or more dependent variables, with respect to one or more independent variables.

· An Ordinary Differential Equation (ODE) involves only ordinary derivatives

· A Partial Differential Equation (PDE) involves partial derivatives.

Definition The order of a DE is the order of the highest-order derivative that appears in the DE

Notation $F(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2})$

Definition A linear DE is any DE that can be written in form:

$$a_0(x)y + a_1(x)y' + a_2(x)y'' \cdots + a_n(x)y^{(n)} = b(x)$$


For a DE to be linear:

1. Y and all of its derivatives must be of the 1st degree
2. Any term that does not include y or any of its derivatives must be a function of x

1.2 Some Mathematical Models

I. Free-falling body

Goal: Find $s(t)$.



Set up a differential equation in S, model it, then solve

$$ma = mg$$


$$\frac{d^2s}{dt^2} = g$$

$$v = \frac{ds}{dt}, g = \frac{dv}{dt}$$

What if there is air resistance. Assume force scales linear with velocity

$$\frac{dv}{dt} = g - \frac{kv}{m} \rightarrow \frac{dv}{dt} = g - \frac{k}{m} \cdot \frac{ds}{dt}$$

II: Series Circuit



Voltage drops:

$$V = L \frac{dI}{dt}, V = L \frac{d^2 q}{dt^2}$$

$$V = IR, V = R \frac{dq}{dt}$$

$$V = \frac{q}{C}$$

$$E(t) = L \frac{d^2 q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C}$$

III: Population Growth

$P = P(t)$ = population at time t — use exponential model

$$\frac{dp}{dt} \propto P \rightarrow \frac{dp}{dt} = kP \rightarrow C e^{kt} \text{ where } C \text{ is the initial population}$$

IV: Population Growth with Finite Capacity

"Carrying Capacity" = N — uses the logistic growth model

$$\frac{dp}{dt} \propto \text{both } P \text{ and amount to carrying capacity } (N-P)$$

$$\frac{dp}{dt} = kP(N - P)$$

V: Chemical Reaction

$A + B \rightarrow C$ Concentrations of A and B decreases by amount of C formed

Can we write DE governing the concentration of C $x(t)$?

The rate at which the reaction takes place \propto Product of the remaining concentrations of A and B

α initial concentration of A

β initial concentration of B

$$\frac{dx}{dt} = k(\alpha - x)(\beta - x)$$

2 First-Order Differential Equations

2.1 Preliminary Theory

Example DE: $y' = 3y \Rightarrow \boxed{y = Ce^{3x}}$ the general solution where C is an arbitrary constant

Add initial condition $y(0) = 5$ plug in $x=0$ to $5 = Ce^{3*0}$, $5 = C * 1$, $C = 5 \Leftarrow$ Initial Value Problem
 $y = 5e^{3x}$ is the general solution for the Initial Value Problem

Theorem $f(x) = \begin{cases} \frac{dy}{dx} = f(x, y) & \text{Differential Equation} \\ y(x_0) = y_0 & \text{Initial Condition} \end{cases}$

Let R be a rectangular region in the xy -plane defined by $a \leq x \leq b, c \leq y \leq d$, that contains the point (x_0, y_0) in its interior.

If $f(x, y)$ and $\partial f / \partial y$ are continuous on R , then there exists an interval I centered at x_0 , and on this interval I there exists a unique solution $y(x)$ for this IVP

Key Questions:

Does every IVP have at least one solution?

If an IVP has a solution is it the only solution?

2.2 Separable Variables

2.3 Homogenous Equations

2.4 Exact Equations

2.5 Preliminary Theory

3 Example Problems with Solutions

3.1

$$\begin{cases} \frac{dy}{dx} = 2xy^{\frac{2}{3}} \\ y(0) = 0 \end{cases} \quad y = 0 \text{ and } y = \frac{x^6}{27} \text{ are solutions}$$

$$\frac{dy}{dx} \frac{x^6}{27} = 2x \cdot \frac{x^4}{9} = y^{\frac{2}{3}}$$

$$\begin{cases} \frac{dy}{dx} = 2yx^{\frac{2}{3}} \\ y(0) = 0 \end{cases} \quad \text{and } y = 0 \text{ is the only solution. This IVP satisfies a certain condition and that makes}$$

it have a unique solution

$$\begin{cases} \frac{dy}{dx} = xy^{\frac{1}{2}} \\ y(0) = 0 \end{cases}$$

Does the IVP have a unique solution? When on \mathbb{R}^2 is $\frac{\partial f}{\partial y}$ continuous? $\frac{\partial f}{\partial y} = \frac{1}{2}xy^{-\frac{1}{2}} = \frac{x}{2\sqrt{y}}$

$$\begin{cases} \frac{dy}{dx} = 3y \\ y(0) = 5 \end{cases} \quad \text{Yes there is a unique solution, } \frac{\partial f}{\partial y} = 3$$

Determine the region R for which the DE would have a unique solution through a point (x_0, y_0) in the region $\frac{dy}{dx} = \sqrt{xy}$

Where on \mathbb{R}^2 is $\frac{\partial f}{\partial y}$ continuous? $\frac{\partial f}{\partial y} = \frac{1}{2}(xy)^{-1/2} * \frac{\partial}{\partial y}(xy) = \frac{x}{2\sqrt{xy}}$

DIY

$$\frac{dy}{dx} - y = x$$