

RH 1.9

MATH 5, Jones

Tejas Patel

Refrigerator Homework

10

$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ first reflects points through the vertical x_2 -axis and then rotates points $\frac{3\pi}{2}$ radians.

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \rightarrow \text{Reflection through the } x_2 \text{ axis from textbook table} \rightarrow \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\frac{3\pi}{2} \text{ rad rotation (90 degree clockwise)} \rightarrow \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

$$\text{Multiplying them together: } \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \boxed{\begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}}$$

17

Show that T is a linear transformation by finding a matrix that implements the mapping. Note that x_1, x_2, \dots are not vectors but are entries in vectors.

$$T(x_1, x_2, x_3, x_4) = (0, x_1 + x_2, x_2 + x_3, x_3 + x_4)$$

$$T = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$$

Since all values inside the matrix are real and make the transformation linear and nothing other than linear, the transformation can be considered linear

23

True. The transformation matrix is calculated using the identity matrix, meaning Identity Matrix \rightarrow Transformation Matrix

25

Yes. Rotations are linear transformations as they scale by 1 and don't have any nonlinear effect on the original point

33

$A = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}$ only has 3 pivots so it only maps into \mathbb{R}^3 , not \mathbb{R}^4 so it is neither onto and also means its not one-to-one since there's a row of zeroes

35

$T \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & -5 & 4 \\ 0 & 1 & -6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ makes the transformation matrix $A = \begin{bmatrix} 1 & -5 & 4 \\ 0 & 1 & -6 \end{bmatrix}$. It is onto since there are 2 pivots, one for each row, but is not one-to-one since its not true that the only element of each row is the pivot

43

For $\mathbb{R}^n \rightarrow \mathbb{R}^m$ A linear transformation T is onto if its image spans the entire codomain \mathbb{R}^m , which means the rank of the transformation matrix A must be equal to m (i.e., the number of linearly independent rows must be m). That wasy $n \geq m$

A linear transformation T is one-to-one if the null space of A is trivial, meaning the only solution to $Ax = 0$ is the zero vector. This occurs when A has full column rank, meaning: $n \leq m$

For T to be both onto ($n \geq m$) and one-to-one ($n \leq m$), we require: $n = m$

Computer Homework: Next 10 Pages

Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework

Question 1, 1.9.3 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Assume that T is a linear transformation. Find the standard matrix of T.

T: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$, rotates points (about the origin) through $\frac{\pi}{6}$ radians.

A =
$$\begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$$

(Type an integer or simplified fraction for each matrix element. Type exact answers, using radicals as needed.)

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework

Question 2, 1.9.5

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Assume that T is a linear transformation. Find the standard matrix of T.

T: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a vertical shear transformation that maps e_1 into $e_1 - 18e_2$ but leaves the vector e_2 unchanged.

A =
$$\begin{bmatrix} 1 & 0 \\ -18 & 1 \end{bmatrix}$$

(Type an integer or simplified fraction for each matrix element.)

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework

Question 3, 1.9.7 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Assume that T is a linear transformation. Find the standard matrix of T.


T: $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ first rotates points through $-\frac{\pi}{3}$ radians and then reflects points through the horizontal x_1 -axis.

A =
$$\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}$$

(Type an integer or simplified fraction for each matrix element. Type exact answers, using radicals as needed.)

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10



Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.9 Homework

Question 4, 1.9.13

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list

Question 1

Question 2

Question 3

Question 4

Question 5

Question 6

Question 7

Question 8

Question 9

Question 10

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation such that $T(\mathbf{e}_1)$ and $T(\mathbf{e}_2)$ are the vectors shown in the figure. Using the figure, sketch the vector $T(2,1)$.

Choose the correct graph below.

A.

B.

C.

D.



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Search tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework

Question 5, 1.9.17 HW Score: 100%, 10 of 10 points
Points: 1 of 1


Show that T is a linear transformation by finding a matrix that implements the mapping. Note that x_1, x_2, \dots are not vectors but are entries in vectors.

$T(x_1, x_2, x_3, x_4) = (x_1 + 9x_2, 0, 2x_2 + x_4, x_2 - x_4)$

A =
$$\begin{bmatrix} 1 & 9 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 1 \\ 0 & 1 & 0 & -1 \end{bmatrix}$$
 (Type an integer or decimal for each matrix element.)

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.9 Homework Question 6, 1.9.33 Part 2 of 2 HW Score: 100%, 10 of 10 points Points: 1 of 1 Save

Question list

Determine if the specified linear transformation is (a) one-to-one and (b) onto. Justify your answer.

$T(x_1, x_2, x_3, x_4) = (x_2 + x_3, x_1 + x_2, x_2 + x_3, 0)$

Question 1

a. Is the linear transformation one-to-one?

A. T is not one-to-one because the columns of the standard matrix A are linearly independent.
 B. T is not one-to-one because the standard matrix A has a free variable.
 C. T is one-to-one because $T(\mathbf{x}) = \mathbf{0}$ has only the trivial solution.
 D. T is one-to-one because the column vectors are not scalar multiples of each other.

Question 2

b. Is the linear transformation onto?

A. T is not onto because the columns of the standard matrix A span \mathbb{R}^4 .
 B. T is not onto because the fourth row of the standard matrix A is all zeros.
 C. T is onto because the columns of the standard matrix A span \mathbb{R}^4 .
 D. T is onto because the standard matrix A does not have a pivot position for every row.


Question 6

Question 7

Question 8

Question 9

Question 10



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework Question 7, 1.9.35 Part 2 of 2 HW Score: 100%, 10 of 10 points Points: 1 of 1 Save

Determine if the specified linear transformation is (a) one-to-one and (b) onto. Justify each answer.

$T(x_1, x_2, x_3) = (x_1 - 5x_2 + 3x_3, x_2 - 6x_3)$

(a) Is the linear transformation one-to-one?

A. T is one-to-one because $T(x) = 0$ has only the trivial solution.
 B. T is not one-to-one because the columns of the standard matrix A are linearly independent.
 C. T is not one-to-one because the columns of the standard matrix A are linearly dependent.
 D. T is one-to-one because the column vectors are not scalar multiples of each other.

(b) Is the linear transformation onto?

A. T is onto because the columns of the standard matrix A span \mathbb{R}^2 .
 B. T is onto because the standard matrix A does not have a pivot position for every row.
 C. T is not onto because the standard matrix A does not have a pivot position for every row.
 D. T is not onto because the columns of the standard matrix A span \mathbb{R}^2 .

Question list ←

Question 1

Question 2

Question 3

Question 4

Question 5


Question 6

Question 7

Question 8

Question 9

Question 10



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 HW

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.9 Homework

Question 8, 1.9.37

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Save

Question list

Describe the possible echelon forms of the standard matrix for a linear transformation T where $T: \mathbb{R}^3 \rightarrow \mathbb{R}^4$ is one-to-one.

Give some examples of the echelon forms. The leading entries, denoted \blacksquare , may have any nonzero value. The starred entries, denoted $*$, may have any value (including zero). Select all that apply.

A.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \end{bmatrix}$$


B.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

C.
$$\begin{bmatrix} \blacksquare & * & * \\ 0 & \blacksquare & * \\ 0 & 0 & * \\ 0 & 0 & \blacksquare \end{bmatrix}$$

D.
$$\begin{bmatrix} \blacksquare & * & * \\ 0 & \blacksquare & * \\ 0 & 0 & \blacksquare \\ 0 & 0 & 0 \end{bmatrix}$$

E.
$$\begin{bmatrix} 0 & \blacksquare & * \\ 0 & 0 & \blacksquare \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

F.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$



Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.9 Homework

Question 9, 1.9.38

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Describe the possible echelon forms of the standard matrix for a linear transformation T where $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ is onto.

Give some examples of the echelon forms. The leading entries, denoted \blacksquare , may have any nonzero value; the starred entries, denoted $*$, may have any value (including zero). Select all that apply.

A.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

B.
$$\begin{bmatrix} 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

C.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

D.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & 0 & \blacksquare \end{bmatrix}$$

E.
$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

F.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & 0 & 0 \end{bmatrix}$$


G.
$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & * \end{bmatrix}$$

H.
$$\begin{bmatrix} 0 & \blacksquare & * & * \\ 0 & \blacksquare & * & * \\ 0 & \blacksquare & * & * \end{bmatrix}$$

I.
$$\begin{bmatrix} 0 & 0 & \blacksquare & * \\ 0 & 0 & 0 & \blacksquare \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10



Linear Algebra and Its Applications | IWM - \$225.88 | Robinhood | Section 1.9 Homework | Do Homework - Section 1.9 H +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355177&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google Scholar tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden Oldie DRIP Flight Schedule Flight Schedule Math 2 Math 5 Physics 43

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.9 Homework

Question 10, 1.9.45 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Let T be the linear transformation whose standard matrix is given. Decide if T is a one-to-one mapping. Justify your answer.

$$\begin{bmatrix} -5 & 7 & -6 & -15 \\ 7 & 4 & -8 & 2 \\ 6 & 9 & 3 & 27 \\ -3 & 2 & 8 & 12 \end{bmatrix}$$

Choose the correct answer below.

A. The transformation T is not one-to-one because the equation $T(\mathbf{x}) = \mathbf{0}$ has only the trivial solution.
 B. The transformation T is not one-to-one because the equation $T(\mathbf{x}) = \mathbf{0}$ has a nontrivial solution.
 C. The transformation T is one-to-one because the equation $T(\mathbf{x}) = \mathbf{0}$ has only the trivial solution.
 D. The transformation T is one-to-one because the equation $T(\mathbf{x}) = \mathbf{0}$ has a nontrivial solution.

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10

