## RH 1.4

## MATH 5, Jones

Tejas Patel

## Refrigerator Homework

5

$$\begin{bmatrix} 1 & 3 & 1 & 0 \\ -4 & -9 & 2 & 0 \\ 0 & -3 & -6 & 0 \end{bmatrix} \rightarrow R_2 + = 4R_1 \rightarrow \begin{bmatrix} 1 & 3 & 1 & 0 \\ 0 & 3 & 6 & 0 \\ 0 & -3 & -6 & 0 \end{bmatrix} \rightarrow R_3 + = R_2, R_2 / = 3 \rightarrow \begin{bmatrix} 1 & 3 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$R_1 - = 3R_2 \rightarrow \begin{bmatrix} 1 & 0 & -5 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \rightarrow x_1 = 5x_3 \quad x_2 = -2x_3 \quad x_3 = x_3 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = x_3 \begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix}$$

7

$$\begin{bmatrix} 1 & 3 & -3 & 7 & 0 \\ 0 & 1 & -4 & 5 & 0 \end{bmatrix} \rightarrow R_1 - = 3R_2 \rightarrow \begin{bmatrix} 1 & 0 & 9 & -8 & 0 \\ 0 & 1 & -4 & 5 & 0 \end{bmatrix} \ x_3 = s, \ x_4 = t$$

$$x_1 = 8t - 9s, \ x_2 = 4s - 5t, \ x_3 = s, \ x_4 = t$$
 
$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = x_3 \begin{bmatrix} 9 \\ -4 \\ 1 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 8 \\ -5 \\ 0 \\ 1 \end{bmatrix}$$

20

$$\begin{bmatrix} 1 & 3 & -5 & 4 \\ 1 & 4 & -8 & 7 \\ -3 & -7 & 9 & -6 \end{bmatrix} \to R_2 - = R_1, \ R_3 + = 3R_1 \to \begin{bmatrix} 1 & 3 & -7 & 4 \\ 0 & 1 & -3 & 3 \\ 0 & 2 & -6 & 6 \end{bmatrix}$$

$$R_3 - 2R_2, R_1 - 3R_2 \rightarrow \begin{bmatrix} 1 & 0 & 2 & -5 \\ 0 & 1 & -3 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} x_3 = t$$

$$x_1 = -5 - 2t, \ x_2 = 3 + 3t$$
  $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -2 \\ 3 \\ 1 \end{bmatrix}$ 

**23** 

Line is y=-x-2 so the equation is 
$$x = \begin{bmatrix} 0 \\ -2 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

28

False: Nontrivial means at least 1 is nonzero, not all of them

40

No. The origin is where b = 0 in all dimensions, so b would need to equal 0

## 1 Computer Homework

- 1.1
- 1.2
- 1.3
- 1.4
- 1.5
- 1.6
- 1.7
- 1.8
- 1.9
- 1.10