

RH 1.8

MATH 5, Jones

Tejas Patel

Refrigerator Homework

17

Since T is linear $T(a\mathbf{x}) = aT(\mathbf{x})$ and $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$.

Given $T(\mathbf{u}) = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $T(\mathbf{v}) = \begin{bmatrix} -1 \\ 3 \end{bmatrix}$.

$$3u = 3 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \end{bmatrix} \quad 2v = 2 \begin{bmatrix} -1 \\ 3 \end{bmatrix} = \begin{bmatrix} -2 \\ 6 \end{bmatrix} \quad 3u + 2v = \begin{bmatrix} 6 - 2 \\ 3 + 6 \end{bmatrix} = \begin{bmatrix} 4 \\ 9 \end{bmatrix}$$

20

$$\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \mathbf{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

The first column of A is $T(\mathbf{e}_1)$, and the second column is $T(\mathbf{e}_2)$.

$$T(x_1, x_2) = x_1 v_1 + x_2 v_2$$

Given that $T(x_1, x_2) = x_1 v_1 + x_2 v_2$, then for the standard basis vectors $\mathbf{e}_1 = (1, 0)$ and $\mathbf{e}_2 = (0, 1)$ we have $T(\mathbf{e}_1) = v_1$, $T(\mathbf{e}_2) = v_2$.

$$A = [v_1 \ v_2] = \boxed{\begin{bmatrix} -2 & 7 \\ 5 & -3 \end{bmatrix}}$$

21

True. A linear transformation is indeed a function (one that takes vectors as inputs and returns vectors as outputs) but it satisfies two special properties:

1. Additivity: $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$.


2. Homogeneity: $T(c\mathbf{u}) = cT(\mathbf{u})$ for any scalar c

Any function that meets both of these properties is called a linear transformation, so it's a specialized type of function.

24

False. The set of all linear combinations of the columns of A is the range (or image) of the transformation function, a subset of its codomain. When we write $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ via $T(\mathbf{x}) = A\mathbf{x}$, the usual codomain is \mathbb{R}^m . The range—the set of all possible outputs that T actually attains—coincides with the column space of A . So the column space is a subspace of the codomain, not the entire codomain itself

31



32

1. Every $\mathbf{x} \in \mathbb{R}^n$ can be written as a linear combination of the spanning vectors v_1, \dots, v_p . In symbols,

$$\mathbf{x} = c_1 v_1 + c_2 v_2 + \dots + c_p v_p.$$

2. Apply T and use linearity:

$$T(\mathbf{x}) = T(c_1 v_1 + c_2 v_2 + \dots + c_p v_p) = c_1 T(v_1) + c_2 T(v_2) + \dots + c_p T(v_p).$$

3. But each $T(v_i) = \mathbf{0}$, by assumption. Hence every term on the right is zero, so

$$T(\mathbf{x}) = \mathbf{0}.$$

Because \mathbf{x} is an arbitrary vector that can span \mathbb{R}^n , it follows that $T(\cdot)$ sends every vector to $\mathbf{0}$.

39

Because v_1, v_2, v_3 is dependent, there exist scalars c_1, c_2, c_3 , not all zero, such that

$$c_1 v_1 + c_2 v_2 + c_3 v_3 = \mathbf{0}.$$

Apply the linear map T to both sides:

$$T(c_1 v_1 + c_2 v_2 + c_3 v_3) = c_1 T(v_1) + c_2 T(v_2) + c_3 T(v_3) = T(\mathbf{0}) = \mathbf{0}.$$

This shows that

$$c_1 T(v_1) + c_2 T(v_2) + c_3 T(v_3) = \mathbf{0},$$

with $(c_1, c_2, c_3) \neq (0, 0, 0)$, meaning $T(v_1), T(v_2), T(v_3)$ is also linearly dependent.

Computer Homework: Next 10 Pages

1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet D github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

☰ Homework: Section 1.8 Homework Question 1, 1.8.4 Part 2 of 2 HW Score: 100%, 10 of 10 points ✓ Points: 1 of 1 Save

Question list ←

Question 1

If T is defined by $T(\mathbf{x}) = A\mathbf{x}$, find a vector \mathbf{x} whose image under T is \mathbf{b} , and determine whether \mathbf{x} is unique. Let $A = \begin{bmatrix} 1 & -4 & 3 \\ 0 & 1 & -3 \\ 3 & -13 & 9 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} -5 \\ -9 \\ -3 \end{bmatrix}$.

Find a single vector \mathbf{x} whose image under T is \mathbf{b} .

$\mathbf{x} = \begin{bmatrix} -50 \\ -12 \\ -1 \end{bmatrix}$

Is the vector \mathbf{x} found in the previous step unique?

A. No, because there is a free variable in the system of equations.
 B. Yes, because there are no free variables in the system of equations.
 C. No, because there are no free variables in the system of equations.
 D. Yes, because there is a free variable in the system of equations.

Question 2

Question 3

Question 4

Question 5

Question 6

Question 7

Question 8

Question 9

Question 10

1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

☰ Homework: Section 1.8 Homework Question 2, 1.8.6 Part 2 of 2 HW Score: 100%, 10 of 10 points ✓ Points: 1 of 1 Save

Question list ←

✓ Question 1

✓ Question 2

✓ Question 3

✓ Question 4

✓ Question 5

✓ Question 6

✓ Question 7

✓ Question 8

✓ Question 9

✓ Question 10


Find a vector \mathbf{x} whose image under T , defined by $T(\mathbf{x}) = A\mathbf{x}$, is \mathbf{b} , and determine whether \mathbf{x} is unique. Let $A = \begin{bmatrix} 1 & 4 & 7 \\ 2 & 12 & 18 \\ 0 & 1 & 1 \\ -2 & -9 & -15 \end{bmatrix}$, $\mathbf{b} = \begin{bmatrix} 14 \\ 36 \\ 2 \\ -30 \end{bmatrix}$.

Find a single vector \mathbf{x} whose image under T is \mathbf{b} .

$\mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix}$

Is the vector \mathbf{x} found in the previous step unique?

A. Yes, because there are no free variables in the system of equations.
 B. Yes, because there is a free variable in the system of equations.
 C. No, because there is a free variable in the system of equations.
 D. No, because there are no free variables in the system of equations.



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.8 Homework

Question 3, 1.8.9 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Find all \mathbf{x} in \mathbb{R}^4 that are mapped into the zero vector by the transformation $\mathbf{x} \mapsto A\mathbf{x}$ for the given matrix A.

$$A = \begin{bmatrix} 1 & -5 & 16 & -9 \\ 0 & 1 & -5 & 3 \\ 3 & -12 & 33 & -18 \end{bmatrix}$$

Select the correct choice below and fill in the answer box(es) to complete your choice.

A. There is only one vector, which is $\mathbf{x} = \boxed{\quad}$.


B. $x_3 \begin{bmatrix} 9 \\ 5 \\ 1 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -6 \\ -3 \\ 0 \\ 1 \end{bmatrix}$

C. $x_3 \boxed{\quad}$

D. $x_1 \boxed{\quad} + x_2 \boxed{\quad} + x_4 \boxed{\quad}$

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.8 Homework

Question 4, 1.8.11 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list

◀ Question 1

Is \mathbf{b} in the range of the linear transformation? Why or why not?

A. No, \mathbf{b} is not in the range of the linear transformation because the system represented by the augmented matrix $[A \ \mathbf{b}]$ is consistent.

B. Yes, \mathbf{b} is in the range of the linear transformation because the system represented by the augmented matrix $[A \ \mathbf{b}]$ is consistent.

C. Yes, \mathbf{b} is in the range of the linear transformation because the system represented by the augmented matrix $[A \ \mathbf{b}]$ is inconsistent.

D. No, \mathbf{b} is not in the range of the linear transformation because the system represented by the augmented matrix $[A \ \mathbf{b}]$ is inconsistent.

Question 2

Question 3

Question 4

Question 5


Question 6

Question 7

Question 8

Question 9

Question 10



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.8 Homework

Question 5, 1.8.12 HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list

Question 1

Question 2

Question 3

Question 4

Question 5

Question 6

Question 7

Question 8


Question 9

Question 10

Let $\mathbf{b} = \begin{bmatrix} -11 \\ 2 \\ -3 \\ 0 \end{bmatrix}$, and let A be the matrix $\begin{bmatrix} 1 & 4 & 11 & 5 \\ 1 & 0 & 3 & -4 \\ 0 & 1 & 2 & 3 \\ -2 & 1 & -4 & 1 \end{bmatrix}$. Is \mathbf{b} in the range of the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$? Why or why not?

Is \mathbf{b} in the range of the linear transformation? Why or why not?

A. No, \mathbf{b} is not in the range of the linear transformation because the system represented by the appropriate augmented matrix is inconsistent.
B. Yes, \mathbf{b} is in the range of the linear transformation because the system represented by the appropriate augmented matrix is consistent.
C. Yes, \mathbf{b} is in the range of the linear transformation because the system represented by the appropriate augmented matrix is inconsistent.
D. No, \mathbf{b} is not in the range of the linear transformation because the system represented by the appropriate augmented matrix is consistent.



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

<mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework>

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.8 Homework

Question 6, 1.8.13
Part 3 of 3

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list ←

Use a rectangular coordinate system to plot $\mathbf{u} = \begin{bmatrix} 6 \\ 1 \end{bmatrix}$, $\mathbf{v} = \begin{bmatrix} -1 \\ 4 \end{bmatrix}$, and their images under the given transformation T. Describe geometrically what T does to each vector \mathbf{x} in \mathbb{R}^2 .

Question 1

$T(\mathbf{x}) = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$

Question 2

Question 3

Question 4

Question 5

Question 6

Which graph below shows \mathbf{u} and its image under the given transformation?

A.

B.

C.

D.

Which graph below shows \mathbf{v} and its image under the given transformation?

A.

B.

C.

D.

What does T do geometrically to each vector \mathbf{x} in \mathbb{R}^2 ?

A. A shear transformation

B. A dilation transformation over the x_1 -axis

C. A projection onto the x_1 -axis

D. A reflection through the origin



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

☰ Homework: Section 1.8 Homework Question 7, 1.8.17 Part 3 of 3 HW Score: 100%, 10 of 10 points ✓ Points: 1 of 1 Save

Question list ←

Question 1

Question 2

Question 3

Question 4

Question 5

Question 6

Question 7

Question 8

Question 9


Question 10

Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation that maps $\mathbf{u} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$ into $\begin{bmatrix} 6 \\ 1 \end{bmatrix}$ and maps $\mathbf{v} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$ into $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$. Use the fact that T is linear to find the images under T of $4\mathbf{u}$, $3\mathbf{v}$, and $4\mathbf{u} + 3\mathbf{v}$.

The image of $4\mathbf{u}$ is $\begin{bmatrix} 24 \\ 4 \end{bmatrix}$.

The image of $3\mathbf{v}$ is $\begin{bmatrix} -3 \\ 6 \end{bmatrix}$.

The image of $4\mathbf{u} + 3\mathbf{v}$ is $\begin{bmatrix} 21 \\ 10 \end{bmatrix}$.



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet D github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

☰ Homework: Section 1.8 Homework Question 8, 1.8.41 Part 4 of 4 HW Score: 100%, 10 of 10 points ✓ Points: 1 of 1 Save

Question list <

Show that the transformation T defined by $T(x_1, x_2) = (3x_1 - 4x_2, x_1 + 4, 5x_2)$ is not linear.

If T is a linear transformation, then $T(\mathbf{0}) = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ and $T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v})$ for all vectors \mathbf{u}, \mathbf{v} in the domain of T and all scalars c, d.

(Type a column vector.)

Check if $T(\mathbf{0})$ follows the correct property to be linear.

$T(0,0) = (3(0) - 4(0), (0) + 4, 5(0))$ Substitute.
= $(0, 4, 0)$ Simplify.

What is true about $T(\mathbf{0})$?


A. $T(\mathbf{0}) = \mathbf{0}$
 B. $T(\mathbf{0}) \neq \mathbf{0}$
 C. $T(\mathbf{0}) = (1,1,1)$
 D. $T(\mathbf{0}) = 4$

Therefore, T is not linear.

Question 8

Question 9

Question 10



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet github hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel ?

Homework: Section 1.8 Homework

Question 9, 1.8.45

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list

- Question 1
- Question 2
- Question 3
- Question 4
- Question 5
- Question 6
- Question 7
- Question 8
- Question 9
- Question 10

The given matrix determines a linear transformation T. Find all x such that $T(x) = \mathbf{0}$.

$$\begin{bmatrix} 4 & -2 & 6 & -5 \\ -1 & 3 & -8 & 0 \\ -5 & 7 & 6 & 4 \\ 7 & -5 & 8 & -8 \end{bmatrix}$$


Select the correct choice below and fill in the answer box within your choice.

A. $x = x_2 + x_3 + x_4$

B. There is only one vector, which is $x = \underline{\hspace{2cm}}$.

C. $x = x_4 \begin{bmatrix} 1.5 \\ 0.5 \\ 0 \\ 1 \end{bmatrix}$

D. $x = x_1 + x_4$



1.8 RH x P Do Homework - Section 1.8 H x Investing | Robinhood x | Linear Algebra and Its Applications x +

mylab.pearson.com/Student/PlayerHomework.aspx?homeworkId=690355176&questionId=1&flushed=false&cld=8147608&back=DoAssignments.aspx?view=homework

cache: - Google S... tpate0562/Quizlet GitHub hosted das... Schedule of Classes LaTeX Manual.pdf Diamine Golden O... DRIP Flight Schedule Flight Schedule Math 2 Math 5

Math 5(4314) Spring 2025 Jones

Tejas Patel

Homework: Section 1.8 Homework

Question 10, 1.8.31
Part 2 of 2

HW Score: 100%, 10 of 10 points
Points: 1 of 1

Question list ←

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the linear transformation that reflects each point through the x_1 -axis, such that $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ where $\mathbf{x} \mapsto A\mathbf{x}$. Make two sketches that illustrate the two properties of a linear transformation.

Choose the correct graph below that shows the property $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$.

A.

B.

C.

D.

Choose the correct graph below that shows the property $T(c\mathbf{u}) = cT(\mathbf{u})$.

A.

B.

C.

D.

Textbook Ask my instructor