

Project 1, Transformations

Tejas Patel

25 March, 2025

1

Matrices defined: $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 2 & -1 \\ 1 & 1 & 10 \end{bmatrix}$, $V = \begin{bmatrix} 1 & 4 & 0 \end{bmatrix}$

When `show(A)`, `show(B)`, `show(V)` was input into the python terminal, the output from SageMath was $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $\begin{bmatrix} 0 & 2 & -1 \\ 1 & 1 & 10 \end{bmatrix}$, $\begin{bmatrix} 1 & 4 & 0 \end{bmatrix}$ respectively. The matrices were repeated back to me as I originally entered them

$$\text{For } QQ \rightarrow RR \Rightarrow \begin{bmatrix} 1.0000000000000000 & 3.0000000000000000 \\ 2.0000000000000000 & 4.0000000000000000 \end{bmatrix}$$

$$\text{For } QQ \rightarrow RDF \Rightarrow \begin{bmatrix} 1.0 & 2.0 \\ 3.0 & 4.0 \end{bmatrix}$$

RR and *RDF* seem to be data types for the matrices. *RR* appears to be an arbitrary precision floating point, outputting the maximum number of decimal places allowed. It won't be as precise as *QQ*, which is symbolic math, nor will it be as fast as *RDF*, which seems to be reduced decimal form, but it will get you a precise decimal answer when you're looking for one.

$$\text{RREF}(B) = \begin{bmatrix} 1 & 0 & \frac{21}{2} \\ 0 & 1 & -\frac{1}{2} \end{bmatrix}$$

The command `I4 = identity matrix(n)` creates an identity matrix of the degree of the input `n`.

$$\text{For the original command: } \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ For input = 2: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ For input = 5: } \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

2

New matrices defined: $C = \begin{bmatrix} 3 & 7 \\ 1 & 6 \end{bmatrix}$

a: In SageMath, to add: $A+C$ results in $\begin{bmatrix} 4 & 9 \\ 4 & 10 \end{bmatrix}$, to subtract $A-C$ results in $\begin{bmatrix} -2 & -5 \\ 2 & -2 \end{bmatrix}$

Lastly, to multiply: $A*C$ results in $\begin{bmatrix} 5 & 19 \\ 13 & 45 \end{bmatrix}$, bonus: $\det(A*C)$ results in -22

b:

$$B^T = \begin{bmatrix} 0 & 1 \\ 2 & 1 \\ -1 & 10 \end{bmatrix} \text{ shows that } \forall x \in B \{x_{ij} \rightarrow x_{ji}\}$$

$A*C==C*A$ returned **false**, meaning $AC \neq CA$

$(A*C).transpose()==A.transpose()*C.transpose()$ returned **false**, meaning $(AC)^T \neq A^T C^T$

$(A+C).transpose()==A.transpose()+C.transpose()$ returned **true**, meaning $(A+C)^T = A^T + C^T$

$(c*A).transpose()==c*A.transpose()$ returned **true** for $c = 6$ and $c = 10$, meaning $(cA)^T = c * A^T$

The left number of columns in the left matrix must equal the number of rows in the right matrix. If m is the number of columns in A and p is the number of rows in B, then $m = p$ must be true. The output of matrix multiplication is going to be of dimension $n \times q$

3

a: The output from the code is 4 graphs, each one containing 2 vectors. The blue arrows represent $P1, P2, P3, P4$ and the purple arrows represent $Q1, Q2, Q3, Q4$.

b: reflects across the y axis (or x_2 axis). Vector $(2, -4)$ becomes $(-2, -4)$

c: Reflects across $y = x$, $(3, 7)$ becomes $(7, 3)$ and $(1, 0)$ becomes $(0, 1)$

d: Reflects across $y = -x$, $(3, 7)$ becomes $(-7, -3)$ and $(1, 0)$ becomes $(0, -1)$

e: Maintains x value, sets y value to 0. $(3, 7)$ becomes $(3, 0)$ and $(0, 1)$ becomes $(0, 0)$

f: Pretty much using the reciprocal of the columns of M, the matrix $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ resolves the reflection across the x_1 axis, where $(3, 7)$ becomes $(-3, 7)$

g: The transformation $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ rotates the vector it is multiplied by by angle θ counterclockwise. $(3, 7)$ becomes approx. $(-2.828, 7.071)$, creating a 45° dihedral angle

4

4.1 Derivation

If $\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = a + b$, then $x_1a + x_2b = a + b$, meaning $x_1 = 1$, $x_2 = 1$

If $\begin{bmatrix} x_3 & x_4 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = a - b$, then $x_3a + x_4b = a - b$, meaning $x_3 = 1$, $x_4 = -1$

Combining these together, we get transformation matrix $A = T = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$

4.2 Check/Proof

Checking $\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ with the Identity Matrix $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$. The expected answer is $\begin{bmatrix} 1+0 \\ 0-1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \cdot 1 + 1 \cdot 0 \\ 1 \cdot 0 + -1 \cdot 1 \end{bmatrix} = \begin{bmatrix} 1+0 \\ 0-1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

We can observe the answer matches the expected, showing the transformation matrix A is the correct transformation. The answer was check with SageMath

5 Determining the Transformation Matrix J

$$\begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ x_{31} & x_{32} \\ x_{41} & x_{42} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 - 5x_2 \\ 0 \\ x_1 + 4x_2 \\ x_2 \end{bmatrix}$$

5.1 Determining individual rows

Row 1: $x_1x_{11} + x_2x_{12} = 2x_1 - 5x_2 \Rightarrow x_{11} = 2$ and $x_{12} = -5$

Row 2: $x_1x_{21} + x_2x_{22} = 0 \Rightarrow x_{21} = 0$ and $x_{22} = -0$

Row 3: $x_1x_{31} + x_2x_{32} = x_1 + 4x_2 \Rightarrow x_{31} = 1$ and $x_{32} = 4$

Row 4: $x_1x_{41} + x_2x_{42} = x_2 \Rightarrow x_{41} = 0$ and $x_{42} = 1$

5.2 Combining the results

Combining the results back into the original matrix form: $J = \begin{bmatrix} 2 & -5 \\ 0 & 0 \\ 1 & 4 \\ 0 & 1 \end{bmatrix}$