## University of Waterloo MATH 213, Spring 2015 Assignment 6

## Question 1

Use the Laplace transform to find the particular solution of the following functions

$$y' + y = \sin(t - 3)u(t - 3) + tu(t - 3) - 3u(t - 3), y(0) = 0$$

Where

$$u(t) = \begin{cases} 0 & \text{when } t < 0\\ 1 & \text{when } t \ge 0 \end{cases}$$

Take the Laplace of both sides:

$$y' + y = \sin(t - 3)u(t - 3) + tu(t - 3) - 3u(t - 3)$$

$$y' + y = \sin(t - 3)u(t - 3) + (t - 3)u(t - 3)$$

$$sY(s) + Y(s) = \frac{e^{-3s}}{s^2 + 1} + \frac{e^{-3s}}{s^2}$$

$$(s + 1)Y(s) = e^{-3s}(\frac{1}{s^2 + 1} + \frac{1}{s^2})$$

$$Y(s) = e^{-3s}(\frac{1}{(s^2 + 1)(s + 1)} + \frac{1}{s^2(s + 1)})$$

Use partial fraction decomposition:

$$\frac{1}{(s^2+1)(s+1)} = \frac{As}{s^2+1} + \frac{B}{s^2+1} + \frac{C}{s+1}$$

$$1 = As^2 + As + Bs + B + Cs^2 + C$$

$$A + C = 0, A + B = 0, B + C = 1$$

$$\therefore A = -0.5, C = 0.5, B = 0.5$$

$$\frac{1}{s^2(s+1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s+1}$$

$$1 = As^2 + As + Bs + B + Cs^2$$

$$A + C = 0, A + B = 0, B = 1$$

$$\therefore A = -1, B = 1, C = 1$$

Take the inverse Laplace:

$$\begin{split} Y(s) &= e^{-3s} (\frac{1}{(s^2+1)(s+1)} + \frac{1}{s^2(s+1)}) \\ Y(s) &= e^{-3s} (-\frac{1}{2} \frac{s}{s^2+1} + \frac{1}{2} \frac{1}{s^2+1} + \frac{1}{2} \frac{1}{s+1} - \frac{1}{s} + \frac{1}{s^2} + \frac{1}{s+1}) \\ y(t) &= (-\frac{1}{2} cos(t-3) + \frac{1}{2} sin(t-3) + \frac{3e^{-(t-3)}}{2} - 1 + (t-3))u(t-3) \end{split}$$

## Question 2