

# Improving Inferences from Randomized Trials: Using per-protocol analyses obtain better estimated of HIV treatment effects.

---

by Timothy Feeney  
SER June 20 2024

e: [feeney@unc.edu](mailto:feeney@unc.edu), bsky: [@tfeend.bsky.social](https://bsky.app/profile/tfeend.bsky.social)



**GILLINGS SCHOOL OF  
GLOBAL PUBLIC HEALTH**

# Outline

---

RCTs

Per-protocol effects

Example using ACTG 5202 Trial

- Population

- Analysis Plan

- Results

- Limitations and Future

# Randomized Trials are a gold standard

---

- Require clear enrollment criteria
- Unambiguous intervention protocol
- Exchangeability:  $Y^a \perp\!\!\!\perp A$  for  $A \in \{0, 1\}$
- Consistency<sup>1</sup>:  $Y = Y^{a=1}A + Y^{a=0}(1 - A)$
- Positivity<sup>2</sup>:  $Pr(A = a | \mathbf{L}) > 0, \forall \ell$  where  $f(a, \ell) > 0$

👉 This allows for unbiased estimation of treatment effects.

---

<sup>1</sup>also known as treatment variation irrelevance

<sup>2</sup> $\mathbf{L}$  is covariate vector

# RCT estimands

---

- Intention-to-treat (ITT) effect:  $E[Y^{r=1} - Y^{r=0}]$
- This is the effect of treatment assignment on outcomes
- Public health focused
  
- Per-protocol (PP) effect:  $E[Y^{r=1, \bar{a}=1} - Y^{r=0, \bar{a}=0}]$
- This is the effect of treatment assignment and adherence on outcomes
- Patient focused<sup>1</sup>

---

<sup>1</sup>Hernan and Robins, NEJM 2016

<sup>2</sup> $\bar{a}$  is the history of treatment adherence

# There is no *one* per-protocol effect

---

- Accounts for adherence
- "Doc, what if I take all my doses like you tell me to?"
- There are six per-protocol parameters that can be estimated<sup>1</sup>
- This can also depend on how the investigator(s) define adherence.

$$E[Y^{r=1, \bar{a}=1}] - E[Y^{r=0, \bar{a}=1}]$$

$$E[Y^{r=1, \bar{a}=1}] - E[Y^{r=0, \bar{a}=0}]$$

$$E[Y^{r=0, \bar{a}=1}] - E[Y^{r=0, \bar{a}=0}]$$

$$E[Y^{r=1, \bar{a}=1}] - E[Y^{r=1, \bar{a}=0}]$$

$$E[Y^{r=0, \bar{a}=1}] - E[Y^{r=1, \bar{a}=0}]$$

$$E[Y^{r=1, \bar{a}=0}] - E[Y^{r=0, \bar{a}=0}]$$



---

<sup>1</sup>Rudolph *et al.* Epidemiology 2020

<sup>2</sup> $r$  = randomization,  $\bar{a}$  = treatment history

# Per-protocol effects can be biased

---


- Frequently done by excluding those not adhering<sup>1</sup>
  - Estimation of the per-protocol effect is much like an *observational study*
  - Thus, bias can arise from non-exchangeability, non-consistency, or non-positivity
- 👉 This can lead to biased estimates

---

<sup>1</sup>Cole *et al.* JAMA Net. Open 2023, Dodd *et al.* Trials 2012

# Per-protocol effects can be biased

---



Can be addressed: inverse probability weighting (IPW), the g-formula, TMLE, can be used to account for selection bias (see next talk)



---

$^0\mathbb{X}$  : vector of covariates

# Per Protocol Causal Identification

---

- Conditional Exchangeability:  
 $Y^g \perp\!\!\!\perp (A_t, C_{t+1}) \mid (\bar{A}_{t-1} = \bar{a}_{t-1}^g, \bar{L}_t = \bar{\ell}_t, C_t = Y_t = 0) \quad \forall t$
- Consistency: if  $\bar{A}_t = \bar{A}_t^g$  then  $\bar{Y}_t = \bar{Y}_t^g$
- Positivity:  $f(a_t^g, C_t = 0 \mid \bar{a}_t^g, \bar{\ell}_t, C_t = Y_t = 0) > 0$  where  $f(\bar{a}_t^g, \bar{\ell}_t, C_t = Y_t = 0) > 0 \quad \forall t$
- No interference:  $\bar{A}_{it}^g \perp\!\!\!\perp \bar{Y}_{jt}^g$  where  $i \neq j$
- No missclassification and correct model specification


Where,  $C$ =censoring,  $t$ =time point from  $0 \dots t$ ,  $g$  is a deterministic treatment strategy, overbar denotes history of values

---

<sup>0</sup>Wen *et al.* Biometrics 2019




# More on Protocol Definition



# An example using an HIV Trial: AIDS Clinical Trial Group (ACTG) 5202

# Role of adherence in HIV treatment efficacy

- Adherence needed for viral suppression varies by treatment regimen.
  - Blanket recommendations fail to capture differences.
- 👉 Understanding of how adherence impacts efficacy is *critical*<sup>1</sup> for:
- Developing new treatments.
  - Maximizing current treatments.



<sup>1</sup> Adimora, Cole and Eron CID 2017

# Example: ACTG 5202 Reanalysis

---

## Overview:

- Phase 3b RCT
- 59 sites, US and Puerto Rico

## Objective:

- Per-protocol analysis modulating protocol definition.
- Target population: The population of HIV+ individuals in the United States
- Estimand:  $E[Y^{r=1, \bar{a}=1}] - E[Y^{r=0, \bar{a}=0}]$  at 48 weeks

## Outcomes: Virologic Failure defined as:


- plasma HIV-1 RNA level  $\geq 1000$  copies /mL between 16 weeks and 24 weeks
- or  $\geq 200$  copies/mL at or after 24 weeks

# ACTG 5202 Study Population

	ABC/3TC		TDF/FTC	
	+ATV	+EFV	+ATV	+EFV
n	463	465	465	464
Female (%)	75 (16.2)	98 (21.1)	78 (16.8)	71 (15.3)
Age Group (%)				
0-25 year	51 (11.0)	43 (9.2)	52 (11.2)	46 (9.9)
25-49 years	353 (76.2)	362 (77.8)	339 (72.9)	356 (76.7)
≥ 50 years	59 (12.7)	60 (12.9)	55 (15.9)	62 (13.4)
Baseline log RNA (med [IQR])	4.64 [4.31, 5.14]	4.68 [4.34, 4.96]	4.55 [4.31, 5.05]	4.65 [4.35, 4.91]
Baseline CD4 (med [IQR])	236 [73, 345]	225 [103, 324]	224 [87, 327]	233 [104, 334]
AIDS Hx (%)	84 (18.1)	88 (18.9)	69 (14.8)	71 (15.3)



# DAG



$j$ : follow up time,  $\mathbf{L}_j$ : vector of covariates at follow up time  $j$   
 $A_j$ : tx adherence at follow up time  $j$ ,  $R$ : randomization,  $Y$ : viral failure or death

# Adherence and Protocol

---

Adherence evaluated in-person at 8, 24, 48, 72, 96, then every 24 weeks and either at the final study evaluation or after virologic failure.

Last Medication	Time Missed	How Close Was Dose Schedule Followed
Never		Never
>3 months ago		Some of the time
1-3 months ago		About half the time
2-4 weeks ago		Most of the time
1-2 weeks ago		All the time
Within the past week		

Adherence Definition	Definition of Variable
0 dose missed OK	No report of missed medication doses
1 dose missed OK	Participant with only one report of missed medication doses
⋮	⋮
4 doses missed OK	Participant with $\geq 10$ reported missed medication doses without overlap in reported timing

# Deviation from Defined Protocols


---

Treatment Group	Censored	1 Dose	2 Dose	3 Dose	4 Dose	5 Dose	Total
ABC/3TC	234	276	110	57	18	7	928
TDF/FTC	211	263	79	38	23	7	929







## Risk of censoring by treatment group



# Cumulative Incidence of Composite Outcome



# TDF/FTC v ABC/3TC at 48 Weeks



# Limitations and Future Plans

---

1. Completed with public access data<sup>1</sup>
2. Reliance on coarse, self-reported medication adherence
3. Assume identification conditions met.<sup>2</sup>
4. Future directions include repeating analysis with g-formula, considering additional protocols.

---

<sup>1</sup>Approved for more granular data from ACTG, awaiting dataset

<sup>2</sup>NB: not guaranteed in per-protocol setting even though it is a trial

# Thank you!


---



**GILLINGS SCHOOL OF  
GLOBAL PUBLIC HEALTH**

I'd like to acknowledge:

- Steve Cole
- Paul Zivich
- Catherine Li
- ACTG 5202
- ACTG 5202 Participants
- Cole Lab Members



My website where you can find a link to my github.