

# When is an ecological network complex? Connectance drives degree distribution and emerging network properties

Timothée Poisot & Dominique Gravel

December 20, 2013

<sup>1</sup> **Authors affiliation:** Université du Québec à Rimouski, Département de Biologie, G5L3A1 Rimouski  
<sup>2</sup> (QC), Canada *and* Québec Centre for Biodiversity Sciences

<sup>3</sup> **Contact:** `t.poisot@gmail.com`

<sup>4</sup> Connectance and degree distributions are important components of the structure of eco-  
<sup>5</sup> logical networks. In this contribution, we use a statistical argument and simple network  
<sup>6</sup> generating models to show that properties of the degree distribution are driven by net-  
<sup>7</sup> work connectance. We discuss the consequences of this finding for (1) the generation of  
<sup>8</sup> random networks in null-model analyses, and (2) the interpretation of network structure  
<sup>9</sup> and ecosystem properties in relationship with degree distribution.

# 1 Introduction

Ecologists developed a strong interest for network theory, as it allowed to make sense of some of the complexity of ecological communities. In contrast to early approaches on “community modules” (groups of a few species within a large community, Holt, 1997) a network level approach allows one to account for the whole community scale (J. A. Dunne, 2006), thus integrating all direct and indirect interactions (Berlow et al., 2009). Ecological networks have often been called “complex” (R. J. Williams & Martinez, 2000), on account of the fact that they represent objects (ecological communities) with complex dynamics (i.-e. non-linear, sensitive to indirect interactions). Because networks are multi-faceted objects with a rich range of structure, ecologists have been looking for emerging properties that can be easily measured and analyzed, and that relate to ecological properties and processes.

Early in the ecological network literature, connectance, *i.e.* the proportion of realized ecological interactions among the potential ones (most often the squared species richness), has been recognized as a central network property (May, 1972; Yodzis, 1980; N. D. Martinez, 1992). In part, this success can be attributed to the relationship between connectance and early definitions of network complexity (Pimm, 1982), and to the fact that connectance predicts reasonably well key dynamical properties of ecological networks (J. A. Dunne, Williams, & Martinez, 2002; 2002) including their stability (May, 1972). More recently, attention shifted from connectance, a community-averaged property, to the degree distribution, that is the statistical properties of the distribution of number of interactions per species. Variation of degree distribution among networks has often been taken as evidence that assembly or interaction mechanisms differ (Vázquez, 2005; Williams, 2011), and increasingly refined methods to estimate degree distribution have been devised (Williams, 2009). Some authors proposed that degree distribution, rather than connectance, is driving higher level network properties such as nestedness or modularity, which are important drivers of network dynamics (Miguel Angel Fortuna et al., 2010).

However, it is worth asking if we were not too quick in focusing most of our research effort on degree

distribution, in detriment to more fundamental work on connectance and its effects. A network, ecological or otherwise, can be viewed as a physical space that edges (interactions) occupy. The size of this space is limited by the number of nodes. This means that there are physical constraints on the filling of a network, due to the fact that placing the first edge will limit the number of ways to place the remaining edges, and so on. For example, there is only one way to have a fully connected network, and there are a limited number of ways to have a network with the lowest possible connectance. For this reason, and given the rising importance of degree distribution in the literature, it is important that we clearly understand how constrained this distribution actually is in relation to connectance. In this contribution, using an argument from combinatorial statistics and simulations of pseudo-random networks under two different models, we present strong evidence that degree distribution, along with other emerging network properties, are constrained (and can be predicted to a certain extent) by connectance. We discuss the consequences of our results for the comparison of different ecological networks, and for the generation of random networks in null-model analyses.

## 2 Statistical argument

Assuming an ecological network made of  $n$  species, and assuming undirected interactions with no self-edges (e.g. no species can interact with itself), there can be at most  $M = n(n - 1)/2$  interactions in this network, in which case it is a complete graph (the results presented below hold qualitatively for both directed graphs, and graphs in which self-edges are allowed). We note this maximal number of links  $M_n$ . With this information in hand, it is possible to know the total number of possible networks given a number  $l$  of interactions. Ecologists are often more familiar with networks being represented as their adjacency matrix, *i.e.* (with minimal simplifications) a matrix with as many rows and columns as there are species, and a 1 at the intersection of two species that interact. In an undirected network, the existence of an edge between species  $A$  and  $B$  imply *two* interactions (*i.e.*  $A \rightarrow B$  and  $B \leftarrow A$ ), and so assuming no self-edges, the total number of ones in the adjacency matrix of a complete, undirected graph, is  $n(n - 1)$ . Throughout this paper, we represent networks as graphs, and not as adjacency

matrices. We consider only the situation of unipartite networks, *i.e.* that can be represented by a square matrix. While the shape of the matrix (*i.e.* the ratio of columns numbers over rows number) will have an impact on the results, we conducted preliminary simulations showing that the results hold qualitatively in bipartite networks of varying shapes.

If we term  $S_n$  the set of all possible  $M_n$  edges in a  $n$ -node network, then the number  $G_{n,l}$  of possible networks with  $l$  links is the number of  $l$ -combinations of  $S_n$ , *i.e.* how many possibilities are there to pick  $l$  edges among  $M_n$ . Formally, this is expressed as  $G_{n,l} = C_l^{M_n}$ , (where  $C_x^y$  is the binomial coefficient, *i.e.* the number of possible ways to pick  $x$  elements among a set of  $y$  elements) or

$$G_{n,l} = \frac{M_n!}{l!(M_n - l)!}$$

Note that this number of possible networks include some graphs in which nodes have a degree of 0, and that in most ecological studies, such nodes will be discarded. We therefore have to evaluate how many of such networks will be found withing  $G_{n,l}$ . In addition, in a null-model context (Bascompte et al., 2003; Miguel A. Fortuna & Bascompte, 2006), having unconnected nodes in random replicates will change the richness of the community, thus possibly biasing the value of randomized emerging properties. As some measures of network structure covary with species richness, if one is to generate a randomly expected distribution of the values of these properties, then it is important to hold species richness constant. Finding out the number of networks in which a given node has a degree of 0 is similar to finding out how many networks exist with  $l$  links between the  $n - 1$  nodes. If one node is removed from the network, there are  $C_{n-1}^n$  possible combinations of nodes (which is  $(n)!/((n-1)!(n-(n-1))!)$ , which further simplifies to  $n$ ). For each of these, there are  $G_{n-1,l}$  possible networks configurations. Note that these networks will also include situations in which *more* than one species has a degree of 0, so that by recurrence, evaluating  $G_{n-2,l}$  and so forth is not necessary (all networks with more than one node of null degree are within the set of the networks with at least one node of null degree). Calling  $R_{n,l}$  the number of networks with  $n$  nodes and  $l$  edges in which all nodes have at least one edge attached, we can write that the number of networks with all nodes

85 having at least one edge is the total number of networks minus the number of networks having at  
 86 least one node of null degree (evaluated for each node), or

$$R_{n,l} = G_{n,l} - C_{n-1}^n \times G_{n-1,l}$$

87 We call the quantities  $R$  and  $G$ , respectively, the *realized* and *total* network space. They measure how  
 88 many networks of  $n$  nodes and  $l$  edges exists, either allowing or preventing the existence of nodes  
 89 with no interactions. Based on this reasoning, we can make two predictions.

90 **Prediction 1:** Because  $C_x^y = C_{y-x}^y$ , it comes that the total network space is largest when  $l = M_n/2$ . As  
 91 in this context the maximal number of edges is  $M_n$ , we define effective connectance as  $Co = l/M_n$ ,  
 92 so  $\max(G_{n,l})$  is reached at  $Co = 1/2$ . The algebraic expression of the maximum value of  $R_{n,l}$  is hard  
 93 to find, but simulations show that it also occurs around  $Co = 1/2$ . In other words, regardless of the  
 94 number of nodes in a network, the “degrees of freedom” of network structure, as indicated by the  
 95 size of the realized and total network spaces, is maximized at intermediate connectance.

96 **Prediction 2:**  $R_{n,l}$  will become asymptotically closer to  $G_{n,l}$  when  $l$  is close to  $M_n$ . In other words,  
 97 there is only one way to fill a network of  $n$  nodes with  $M_n$  interactions, and in this situation there  
 98 is no possibility to have nodes with a degree of 0. In the situation in which  $l = M_n$ ,  $G_{n,l} = C_{M_n}^{M_n} = 1$ ,  
 99 given that  $M_n > M_{n-1}$ , it comes that  $G_{n,l} = R_{n,l} = 1$ . Intuitively enough, this implies that ecological  
 100 systems in which connectance is high will display very little variation from one another, as far as the  
 101 distribution of emergent network properties (e.g. variance of the degree distribution, nestedness, ...) is  
 102 concerned.

103 We now illustrate these predictions using networks of 10 nodes, with a number of edges varying from  
 104 10 to  $M_{10}$  (i.e. 45 edges). As illustrated in Fig. 1, the size of the network space has a hump-shaped  
 105 relationship with connectance, and the size of the realized network space becomes closer to the size  
 106 of the total network space when connectance increases.

107 In Fig. 2, we show that regardless of the network size, the relative size of the realized network space  
 108 increases with connectance. The rate at which it occurs increases with network size. However, in all




Figure 1: Size of the total and realized network space for  $n = 10$ . As predicted in the main text, (1) the size of network spaces peaks at  $Co = 1/2$ , and (2) the size of the realized network space becomes asymptotically closer to the size of the total network space when connectance increases.

109 cases, when connectance is low, there are only a very small proportion of the total network space in  
 110 which all nodes have at least one edge. This suggest that the structure of extremely sparse networks is  
 111 also strongly constrained. This is congruent with historical findings by Erdos & Rényi (1959), namely  
 112 that the probability of each node being connected to the graph largest connected component (*i.e.* any  
 113 set of vertices of which any two are connected by at least one path) increases with average degree  
 114 (thus for high connectances, all nodes are likely to be connected to the giant component, hence no  
 115 node has a degree of 0). In the context of ecology, in which most networks have a low connectance, it  
 116 implies that generating random networks to test null hypotheses can be a computationally intensive  
 117 task, as the realized network space is (proportionally) small.




Figure 2: Relative size of the realized network space compared to the total network space when connectance increases, for four different network sizes.

### 3 Simulations

In the previous part, we show mathematically that connectance (the number of realized *vs.* possible interactions), relative to the network size, determines the size of the *network space*, *i.e.* how many possible network combinations exist. Based on this, we can therefore expect that the degree distribution will be contingent upon network connectance. Specifically, we expect that the variance of the degree distribution, which is often related to ecosystem properties and other network structures (Miguel Angel Fortuna et al., 2010), will display a hump-shaped relationship with connectance. The mean, kurtosis, and skewness of the degree distribution should all vary in a monotonous way with connectance.

In the simulations below, we use networks of 30 nodes, filled with 35 to  $M_{30}$  interactions, by steps of 10. We use two different routines to generate random networks, that are contrasted in the way they distribute edges among nodes. First, we generate Erdős-Rényi (ER, undirected) graphs, meaning that every potential interaction has the same probability of being realized (Erdos & Rényi, 1959). We use an algorithm inspired by Knuth (1997), allowing to fix the number of edges in the graph rather than the probability of an edge occurring, although the generated graphs have the same properties as the original ER model. A total of 19000 networks are generated this way. Second, we use the niche model of food webs (R. J. Williams & Martinez, 2000), which generates (directed) networks under rules representing hypothesized mechanisms of prey-selection in empirical ecosystems (Gravel et al., 2013). This particular model assumes that the existence of interactions is constrained by the position of species along a “niche” axis, for example body size. Other randomization methods for food webs exists, but given that Stouffer et al. (2005) showed that they yield similar degree distributions to the niche model, we will not use them here. A total of 500 replicates for each value of  $l$  are generated. All networks generated with the two models are free of self-edges and nodes with a null degree.

For each replicate, we measure the degree distribution and report its variance, coefficient of variation, kurtosis, and skewness. In addition, for each network, we fit a power-law distribution on the sorted degree distribution using the least-squares method; we report the power-law exponent.






Figure 3: Statistical descriptors of the degree distribution of randomized networks,  $n = 30$ , increasing connectance. These results show that central properties of the degree distribution are contingent upon connectance, at a given network size, and under a given network generation model. ER networks are in blue, niche-model networks are in red. Each point represent a single generated network.

Qualitatively, the random graphs and the niche networks behave exactly the same. With the exception of the kurtosis, *all* statistical descriptors of the degree distribution were influenced by the effective connectance (Fig. 3). As predicted in the previous part, variance of the degree distribution is hump-shaped with regard to connectance, which implies that as average degree increases with connectance, the coefficient of variation of the degree distribution decreases at high connectances. Note also that the range of variances in the degree distribution is higher at intermediate connectances, but lower at the extreme. Due to the fact that the Erdős-Rényi graphs we simulate are essentially Poisson random graphs, it is expected that the variance of their degree distribution would be lower than for the niche model, which in contrast *forces* strong difference in the degree of species according to their niche position.

Kurtosis is independent of connectance, while skewness decreases with connectance. This result is expected. Positively skewed distribution have longer or fatter right tails, indicating mostly low values (low degree): unconnected networks are made mostly of species with a weak generality (Schoener, 1989). On the other hand, negative skewness indicate that most of the values in the distribution are high. Ecologically, it means that most species are wide-range generalists, which happens in densely connected networks. This bears important ecological consequences, as it indicates that due to physical constraints acting on the filling of interactions within the graphs, networks with intermediate connectances are expected to have species with both low and high generality (Schoener, 1989).

The estimate of the power-law exponent increases when connectance increases (Fig. 4). This indicates that the degree distribution flattens when connectance increases. Taken with the elements presented above, we show that all of the estimators of the degree distribution vary strongly with connectance of the network. Although power-laws should be truncated as soon as the probability of a species having a degree of  $2 \times n$  (or  $n$  in undirected networks) is not negligible, and as such the fitting of power-laws should not be done on highly connected networks for practical purposes, this result emphasizes the key role of connectance in driving central network structure properties.




Figure 4: The estimate of the power-law exponent increases with connectance, reaching a flat distribution for complete graphs.

## 4 Practical consequences

Randomized null models are often used to estimate how much a given network property deviates from its random expectation (Flores et al., 2011). Our results show two things. First, except for extremely high or low connectance, the proportion of the network space that will be explored using  $10^3$  or  $10^4$  replicates (typical values in null models analyses) is orders of magnitude smaller than the *realized* network space. Although this is somewhat compensated by the fact that a part of these networks are isomorphic, the risk of inferring deviation from the random expectation based on a drastically small sampling of the network space is real, and un-addressed; at the very least, it seems that that intensity of the replication should be dependent upon the connectance of the network one tries to replicate. On the other hand, when connectance is high, the number of unique network combinations decreases, and there is a risk to generate a number of replicates that is larger than the realized network space, thus decreasing the information content of the randomizations. To the best of our knowledge, these issues have seldom be addressed in the literature on ecological network randomization.

Second, generating null models with a low connectance is a computationally intensive task. When connectance decreases, the *realized* network space decreases faster than the *total* network space, meaning that the probability of picking a network with no 0 degree nodes (which is simply  $R_{n,l}/G_{n,l}$ ) goes toward zero. For this reason, classical rejection sampling (accept the random network if no nodes have no edges, reject it if not) is bound to take an unreasonable amount of time in networks with low connectance. In addition, there is a risk of selecting some particular types of networks. It makes intuitive sense that networks with extremely skewed degree distributions have less chance of being generated this way, as when a few nodes collect most of the edges, the probability than the remaining nodes each have at least one edge decreases. To the best of our knowledge, this source of bias has not received important attention in the literature. For this reason, using a purely random matrix shuffling as a starting point, then swapping interactions until no free nodes remain, seems to be a promising way to address this problem. Given the important of null-model approaches in network analysis, the generation of efficient and unbiased algorithms is a fruitful research avenue.

## 5 Conclusions

Connectance is an extremely intuitive property of network, expressing how much of the potential interactions are realized. Through statistical reasoning and simple simulations using models of random networks, we show that for a given number of species, connectance drives (i) how many different networks exist, and (ii) some key elements of the degree distribution. We observed both among and between model quantitative changes in degree distribution along a connectance gradient. The niche model is a particularly striking example of this, with the variance in the degree distribution increasing 50-fold when connectance moves from 0.1 to 0.5. This result has practical implications for network comparisons. As descriptors of degree distribution vary with connectance, connectance should be factored out from all analyses. So as to avoid colinearity issues, this can be done by either working on the residuals of the degree distributions' property of interest. To some extent, the impact of connectance is lesser in the 0.05-0.3 range where most empirical food webs lies (although bipartite networks can have much higher connectances), but the effect is high enough that it should not be ignored: at equal number of species, networks with different connectances are expected to have different degree distributions.

Finally, this analysis raises interesting ecological questions. Early analyses focusing on degree distribution argued that ecological mechanisms were responsible for the shape of the distribution (Vázquez, 2005; Miguel Angel Fortuna et al., 2010; Williams, 2011). In this contribution, we show that connectance will impose a lower and higher limit for the shape of the degree distribution. Given this information, it is time to bring the debate full-circle: is connectance the cause of observed network properties, or an emergent property of pairwise species interactions? As the later seems far more likely, it now makes sense to focus on why some networks deviate, or not, from the expected degree distribution knowing their connectance. As the density of interaction plays such a central role in May's criteria for stability (May, 1972), clarifying how connectance is shaped by mechanisms regulating pairwise species interactions offers the opportunity of integrating the effects of these mechanisms up to their impact on emergent, community-wide properties. Okuyama & Holland (2008) showed

that in mutualistic systems, resilience (to perturbation) is affected both by degree network size and interaction strength, but also by the degree distribution and connectance of the network; we show here that degree distribution and connectance are tightly linked, and alternative approaches to the question of resilience can focus on the deviation of degree distribution knowing the connectance, rather than the “raw” degree distribution. In keeping with the results we present in the first part of the paper, Rozdilsky & Stone (2001) report that there is a inverted hump-shaped relationship between connectance, and the proportion of “feasible and stable” systems, with the lowest proportion of such systems being found at  $C_0 \approx 0.5$ . This strongly suggests that the same mechanisms that limit the realized network space may affect ecological properties, thus emphasizing the need not to discard connectance in profit of more emerging properties in future ecological network analyses.

**Acknowledgements:** We thank Luis Gilarranz, Miguel Lurgi, and Enrico Rezende for comments, and Amael LeSquin for discussions on algebra.

## 6 References

Bascompte, J., Jordano, P., Melián, C. J., & Olesen, J. M. 2003. *The nested assembly of plant–animal mutualistic networks*. Proceedings of the National Academy of Sciences of the United States of America 100(16):9383–9387.

Berlow, E. L., Dunne, J. a, Martinez, N. D., Stark, P. B., Williams, R. J., & Brose, U. 2009. *Simple prediction of interaction strengths in complex food webs*. Proceedings of the National Academy of Sciences of the United States of America 106(1):187–91.

Dunne, J. A. 2006. *The Network Structure of Food Webs*. In J. A. Dunne & M. Pascual (Eds.), Ecological networks: Linking structure and dynamics (pp. 27–86). Oxford University Press.

Dunne, J. A., Williams, R. J., & Martinez, N. D. 2002. *Food-web structure and network theory: the role of connectance and size*. Proceedings of the National Academy of Sciences of the United States of America 99(20):12917.

245 Dunne, J. A., Williams, R. J., & Martinez, N. D. 2002. *Network structure and biodiversity loss in food*  
 246 *webs: robustness increases with connectance*. Ecology Letters 5(4):558–567.

247 Erdos, P., & Rényi, A. 1959. *On random graphs*. Publicationes Mathematicae Debrecen 6:290–297.

248 Flores, C. O., Meyer, J. R., Valverde, S., Farr, L., & Weitz, J. S. 2011. *Statistical structure of host-phage*  
 249 *interactions*. Proceedings of the National Academy of Sciences of the United States of America  
 250 108(28):E288–297.

251 Fortuna, M. A., & Bascompte, J. 2006. *Habitat loss and the structure of plant-animal mutualistic*  
 252 *networks*. Ecology Letters 9(3):281–286.

253 Fortuna, M. A., Stouffer, D. B., Olesen, J. M., Jordano, P., Mouillot, D., Krasnov, B. R., Poulin, R., &  
 254 Bascompte, J. 2010. *Nestedness versus modularity in ecological networks: two sides of the same coin?*  
 255 *Journal of Animal Ecology* 78:811–817.

256 Gravel, D., Poisot, T., Albouy, C., Velez, L., & Mouillot, D. 2013. *Inferring food web structure from*  
 257 *predator-prey body size relationships*. Methods in Ecology and Evolution.

258 Holt, R. D. 1997. *Community modules*. In A. C. Gange & V. K. Brown (Eds.), *Multitrophic Interactions*  
 259 *in Terrestrial Ecosystems* (pp. 333–349). Blackwell Science.

260 Knuth, D. E. 1997. *Volume 2: Seminumerical Algorithms*. The Art of Computer Programming 192.

261 Martinez, N. D. 1992. *Constant connectance in community food webs*. The American Naturalist  
 262 139(6):1208–1218.

263 May, R. M. 1972. *Will a large complex system be stable?*. Nature 238:413–414.

264 Okuyama, T., & Holland, J. N. 2008. *Network structural properties mediate the stability of mutualistic*  
 265 *communities*. Ecology Letters 11(3):208–216.

266 Pimm, S. L. 1982. *Food Webs*. London: Chapman & Hall.

267 Rozdilsky, I. D., & Stone, L. 2001. *Complexity can enhance stability in competitive systems*. Ecology  
 268 Letters 4(5):397–400.

- 269 Schoener, T. W. 1989. *Food webs from the small to the large*. Ecology 70(6):1559–1589.
- 270 Stouffer, D. B., Camacho, J., Guimera, R., Ng, C. A., & Amaral, L. A. N. 2005. *Quantitative patterns in*  
271 *the structure of model and empirical food webs*. Ecology 86(5):1301–1311.
- 272 Vázquez, D. P. 2005. *Degree distribution in plant-animal mutualistic networks: forbidden links or*  
273 *random interactions?*. Oikos 108(2):421–426.
- 274 Williams, R. J. 2009. *Simple MaxEnt models explain food web degree distributions*. Theoretical Ecology  
275 3(1):45–52.
- 276 Williams, R. J. 2011. *Biology, Methodology or Chance? The Degree Distributions of Bipartite Ecological*  
277 *Networks*. (J. Langowski, Ed.) PLoS One 6(3):e17645.
- 278 Williams, R. J., & Martinez, N. D. 2000. *Simple rules yield complex food webs*. Nature 404(March):180–  
279 183.
- 280 Yodzis, P. 1980. *The connectance of real ecosystems*. Nature 284:544–545.