# Neural and psychological underpinnings of gambling disorder:

2	A review
3	
4	Jon E. Grant*, Department of Psychiatry & Behavioral Neuroscience, University of Chicago,
5	Chicago, IL, USA.
6	Brian L. Odlaug, Department of Public Health, Faculty of Health and Medical Sciences,
7	University of Copenhagen, Copenhagen, Denmark.
8	Samuel R. Chamberlain, Department of Psychiatry, University of Cambridge, UK; & Cambridge
9	and Peterborough NHS Foundation Trust, United Kingdom.
10	
11	
12	Correspondence:
13	Jon E. Grant, JD, MD, MPH
14	Professor, Department of Psychiatry & Behavioral Neuroscience
15	University of Chicago, Pritzker School of Medicine
16	5841 S. Maryland Avenue, MC 3077
17	Chicago, IL 60637; USA
18	jongrant@uchicago.edu
19	
20	Number of Words: 3009
21	Number of Figures: 0
22	

# **Abstract**

Gambling disorder affects 0.4 to 1.6% of adults worldwide, and is highly comorbid with other mental health disorders. This article provides a concise primer on the neural and psychological underpinnings of gambling disorder based on a selective review of the literature. Gambling disorder is associated with dysfunction across multiple cognitive domains which can be considered in terms of impulsivity and compulsivity. Neuroimaging data suggest structural and functional abnormalities of networks involved in reward processing and top-down control. Gambling disorder shows 50-60% heritability and it is likely that various neurochemical systems are implicated in the pathophysiology (including dopaminergic, glutamatergic, serotonergic, noradrenergic, and opioidergic). Elevated rates of certain personality traits (e.g. negative urgency, disinhibition), and personality disorders, are found. More research is required to evaluate whether cognitive dysfunction and personality aspects influence the longitudinal course and treatment outcome for gambling disorder. It is hoped that improved understanding of the biological and psychological components of gambling disorder, and their interactions, may lead to improved treatment approaches and raise the profile of this neglected condition.

**Keywords**: gambling; cognition; personality; genetics; imaging

### 1. Introduction

Gambling disorder is characterized by persistent and recurrent maladaptive patterns of gambling behavior, leading to impaired functioning (1). Although most people who engage in gambling do so responsibly and without consequent functional impairment, some individuals find that they become preoccupied with gambling and cannot control their behavior despite multiple negative consequences (2). Surveys suggest that the prevalence of gambling disorder in the general United States population ranges from 0.42% to 1.9%, and similar rates have been reported worldwide (3-5). As such, recognition of why some individuals cannot control their gambling behavior appears worthy of attention from a global public health perspective (6). In recognition of Gambling disorder representing a prototypical 'behavioral addiction', it has been recently reclassified as a 'Substance-Related and Addictive Disorder) in the Diagnostic and Statistical Manual Version 5 (DSM-5) (1).

There exist several comprehensive reviews of specific aspects of gambling disorder (7-12). The aim of this paper is to provide a concise primer examining the neurobiological and psychological

There exist several comprehensive reviews of specific aspects of gambling disorder (7-12). The aim of this paper is to provide a concise primer examining the neurobiological and psychological underpinnings of gambling disorder, incorporating recent evidence derived from the neurosciences. We highlight implications for new treatment directions, along with limitations of this approach and areas in which research is lacking.

### 2. Pathophysiology of gambling disorder

The behaviors that characterize gambling disorder can be regarded as impulsive, in that they are often poorly thought out (or undertaken without adequate forethought), risky, and result in deleterious long-term outcomes (13). Developmentally, impulsive behavior that underlies

gambling disorder tends to begin during late adolescence or early adulthood (14). While the longitudinal profile of Gambling disorder has received little research attention, for some individuals it is likely that patterns of behavior become ingrained and persist over time, especially in the absence of prompt treatment interventions (3, 9).

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

67

64

65

66

# 2.1. Neurocognition

People with gambling disorder often manifest cognitive deficits consistent with tendencies towards impulsivity. Objective brain-based measurable traits that deconstruct top-level phenotypes into meaningful markers more closely related to the underlying etiology are important in trying to understand the neurobiology of Gambling disorder and its relationship with other conditions (15). Deficits in aspects of inhibition, working memory, planning, cognitive flexibility, and time management/estimation have been reported in individuals with gambling disorder compared to healthy volunteers (12). Individuals with gambling disorder also tend to prefer small immediate rewards rather than larger delayed rewards, to the detriment of long-term task outcomes (i.e. they show abnormally elevated 'delay discounting') (16). Impulsivity is not the only aspect of gambling disorder with other cognitive domains likely present to varying degrees in gambling disorder. Gambling disorder for many individuals, for example, is associated with features of compulsivity (17). People with gambling disorder often describe the behavior in ritualistic terms such as the need for "lucky" numbers or clothing to result in favorable outcome. In addition, the nature of gambling behavior may change over time, with early gambling being driven by reward, and later (more chronic) gambling being triggered by aversive/stressful stimuli (3), or being undertaken in order to avert anxiety (17). As such,

there may be a shift from an initial behavior that is reward-seeking (impulsive) towards one that persists to avoid negative consequences or in a habitual fashion (compulsive). Individuals with gambling disorder often score high on the Padua Inventory, a measure of compulsivity (18) and display marked response perseveration (19,20) and difficulties with cognitive flexibility (21).

Although studies of gambling disorder demonstrate that the behavior is associated with diminished performance on inhibition, time estimation, cognitive flexibility, decision-making, spatial working memory, and planning tasks, a temporal relationship has not been established between cognitive deficits and clinically significant symptoms. Most likely, some cognitive deficits predispose (perhaps running in families and representing candidate 'endophenotypes' or intermediate markers of risk), while others could be a consequence of recurrent engagement in gambling itself. While studies of cognitive functioning in unaffected close relatives of people with gambling disorder are lacking, findings from people 'at-risk' of gambling disorders suggest that deficits in decision-making (dependent on neural circuitry including the orbitofrontal and insular cortices) are evident before the illness, while some other domains may be relatively spared (22). Gambling addiction represents a useful model for exploring the 'cause versus effect' issue in addiction more broadly, since chronic gambling is presumably unlikely to exert toxic effects on the brain, as compared to chronic substance misuse.

# 2.2. Neuroimaging

A sparse amount of research on possible neurobiological correlates of gambling disorder currently exists (for reviews, please see 11-12). Most studies have focused on functional rather than structural neuroimaging abnormalities. One functional magnetic resonance imaging (fMRI)

study of gambling urges in male pathological gamblers suggested that gambling disorder is associated with relatively decreased activation within cortical, basal ganglionic and thalamic brain regions compared to control subjects (23). Recent neuroimaging studies have demonstrated that gamblers also show hyporesponsiveness of the dorsomedial prefrontal cortex compared to healthy controls during successful (as well as failed) response inhibition, along with a hypoactive reward system (24-26). Using a graph theoretical approach (network modeling), there was evidence for abnormalities in distributed brain networks in gambling disorder versus controls, such as reduced local efficiency in the left supplementary motor area, and hyperconnectivity between frontal brain regions including the right inferior frontal gyrus (27). In terms of brain structure, there is some evidence that gambling disorder is associated with excess volume of the ventral striatum and right prefrontal cortex (28).

Another area of neuroimaging research in gambling disorder is the use of radioligand measures in conjunction with positron emission tomography (PET). Using this technique, the status of neurochemical systems in people with gambling disorder, both in the resting state and in response to pharmacological challenge, can be explored. Research so far has focused on the dopamine system, given its established importance in substance addiction and more generally in reward-processing (29). In substance addictions, there is considerable evidence that chronic substance intake is associated with downregulation of striatal D2 receptors (30). Interestingly, radioligand studies so far suggest that gambling disorder is not associated with such dopaminergic D2 downregulation. In a study using raclopride (D2/D3 receptor binding) and propyl-hexahydro-naphtho-oxazin (PHNO; D3 receptor binding), no significant differences in inferred striatal dopamine receptor binding were found between people with gambling disorder

and healthy controls (31). However, PHNO binding in the substantia nigra correlated significantly with gambling symptom severity. In another study, using raclopride (D2/D3 receptor binding), no significant differences were found between gambling disorder subjects and controls in terms of inferred striatal dopamine receptor binding (32); but 'urgency' correlated negatively with raclopride binding in the gambling disorder group. Another study, using raclopride, similarly reported no group differences between gambling disorder and controls; but did find that dopamine receptor binding was associated with sensation-seeking in general (33). In all, these radioligand studies suggest that D2 receptor downregulation is not a general feature of gambling disorder, in contrast to findings in substance use disorders. This is consistent with the view that D2/D3 receptor abnormalities in substance use disorders are a consequence of the effects of chronic drug intake on the reward pathways. Dopamine status is relevant to personality-related factors (e.g. sensation-seeking) implicated in the development of gambling disorder. It may also be that other aspects of the dopamine system, not measured using the above ligands, are abnormal in gambling disorder. For example, one raclopride-PET study found an inverted 'U' relationship between striatal dopamine release and gambling task performance in pathological gamblers but not in controls, suggesting enhanced dopaminergic sensitivity to uncertainty in gamblers (34).

150

151

152

153

154

155

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

Neuroimaging studies to date, do not permit characterization of the temporal relationship between the manifestation of neural abnormalities and the symptoms that comprise gambling disorder. As with the neurocognitive findings, abnormal brain structure and function could occur in people 'at-risk' before symptoms develop, alternatively stem from the disorder itself, or perhaps even reflect a secondary or incidental epiphenomenon.

# 2.3. Genetic predisposition

Studies have found that approximately 20% of the first-degree relatives of individuals with gambling disorder also have gambling disorder (3). Research examining familial aggregation of gambling disorder found that individuals with a problem gambling parent were 3.3 times more likely to have gambling disorder (35). In a study using a control group to examine familial aggregation, lifetime estimates of gambling disorder were significantly higher in family members of gamblers (8.3%) compared to control subjects (2.1%) (36). Data from the Vietnam Era Twin Registry (male adults) have shown that the heritability of gambling disorder is approximately 50-60% (37-38). Further analyses of personality features and their association with the heritability of gambling disorder have found that low self-control is associated with the genetic risk for gambling disorder in women (39). As discussed in the subsequent section, various polymorphisms in genes coding for components of brain neurochemical systems (e.g. dopaminergic and serotonergic systems) have been associated with gambling disorder.

# 2.4. Neurobiological factors

Multiple neurotransmitter systems (e.g., dopaminergic, glutamatergic, serotonergic, noradrenergic, opioidergic) have been implicated in the pathophysiology of gambling disorder (3, 40-41). Dopamine is involved in learning, motivation, and the salience of stimuli, including rewards. As discussed in section 2.3, radioligand PET studies militate against an obvious D2/D3 receptor binding abnormality being evident in gambling disorder in the resting state.

Nonetheless, alterations in dopaminergic pathways have been proposed as underlying the seeking of rewards that trigger the release of dopamine and produce feelings of pleasure. In addition,

neuroimaging studies examining pharmacological challenges using dopamine agonists have reported that during the anticipation of monetary rewards, a dopamine agonist increases the activity of the nucleus accumbens and weakens the interaction between the nucleus accumbens and the prefrontal cortex, leading to an increase in impulsive behaviors (42). Dopamine receptor agonist medication appears to predispose the dopaminergic reward system to mediate an increased appetitive drive leading to changed neural processing of negative consequences and learning of contingencies (43). In terms of molecular genetic studies, the D2A1 allele of the D2 dopamine receptor gene (DRD2) has been reported as increased in frequency in individuals with gambling disorder (for a review see 39). Other research has also implicated allelic variants of the DRD1 and DRD3 genes as having an association with gambling disorder (3).

There is also a persuasive body of preclinical evidence suggesting a critical role for glutamate transmission and glutamate receptors in drug reward, reinforcement, and relapse. Glutamate appears to be implicated in long-lasting neuroadaptations in the corticostriatal circuitry (44). An imbalance in glutamate homeostasis results in changes in neuroplasticity that adversely affects communication between the prefrontal cortex and the nucleus accumbens, thereby resulting in reward-seeking behaviors (45). Glutamate is also involved in associative learning between stimuli and promotes the immediate approach response through its link to the dopamine reward system (41). Data from cerebrospinal fluid studies also suggest a dysfunctional glutamate system in gambling disorder (46).

Animal studies of gambling behavior provide evidence that the serotonergic system also appears to play a role in poor decision-making (47) and impaired performance on a gambling task (48).

Serotonin is known as a modulator of neuroplasticity events. A polymorphism in the serotonin transporter gene has been associated with gambling disorder and is found more frequently in males with gambling disorder (49). More recent research found a significant association of the C/C genotype of the serotonin receptor 2A T102C (rs 6313) polymorphism and the gambling disorder phenotype (50). Other support for dysfunction within the serotonergic system in gambling disorder has been shown with decreased levels of platelet monoamine oxidase B (MAO-B) (a peripheral marker of serotonergic function), low levels of serotonin metabolites (5-HIAA) in the cerebrospinal fluid, and a euphoric response to serotonergic pharmacologic challenge studies (3, 40).

Norepinephrine (noradrenaline) appears to be especially involved in decision-making when contingencies are unexpectedly changed and alternatives are explored (51-52). Selective inhibition of norepinephrine reuptake results in reduced premature responding, especially under circumstances when task performance is suboptimal due to demanding task conditions or inherently high baseline levels of impulsive action (53-54). Studies have found that individuals with gambling disorder have significantly higher cerebrospinal fluid levels of 3-methoxy-4-hydroxy-phenylglycol, the main metabolite of the noradrenergic system (55). In addition, individuals with gambling disorder maintained significantly higher noradrenergic levels throughout an entire gambling session whereas healthy controls exhibited elevated levels only at the onset of the gambling session (56).

Preclinical evidence indicates that opioid receptors are distributed widely in the mesolimbic system, and are implicated in the hedonic aspects of reward processing (57-58).

An fMRI study of the  $\mu$ -opioid antagonist naloxone found attenuated reward-related responses in the ventral striatum and enhanced loss-related activity in the medial prefrontal cortex on a wheel of fortune task in healthy volunteers (59). Specifically, the authors used an fMRI gambling task and found that naloxone reduced pleasure ratings for larger rewards and dampened the associated brain responses in the anterior cingulate cortex. Naloxone was also associated with negative outcomes being rated as being more unpleasant, implicating the opioid system both in reward-and aversive-processing (59). Gambling has been associated with elevated blood levels of the endogenous opioid  $\beta$ -endorphin (60), and modulation of the opioid system through opioid receptor antagonists (61) and partial agonists (62-63) has shown significant promise in the treatment of gambling disorder.

# 3. Psychological aspects of gambling disorder

Relationships between gambling disorder and aspects of personality can be considered from several perspectives, including in relation to personality traits (typically measured using questionnaires such as the Barratt Impulsivity Questionnaire), in relation to formal personality disorders, and in relation to other potentially life-long enduring traits (such as aspects of cognition).

### 3.1. Gambling disorder and personality traits

The assessment of personality traits is an evolving field. While questionnaire-based measures relating to personality have proven useful in exploring aspects of gambling disorder, it can be difficult to relate them to underlying brain function (64-65).

Support for impulsivity as a personality characteristic of individuals with gambling disorder rather than transient impulsive behavior, comes from numerous studies over the years (for a

review, please see 66), including a recent study of 37 individuals which found that trait, rather

than state, questionnaire-based impulsivity is associated with gambling disorder (67).

sensation-seeking compared to women (71).

The relationship between impulsivity and gambling, however, may be impacted by a variety of factors, including socioeconomic status, age of onset, and gender. One study found that self-reported impulsivity was associated with the onset of gambling behavior but only in the case of individuals reporting a low socioeconomic background (68). Similarly, in a sample of 1004 males from low socioeconomic status areas, impulsivity at age 14 was related to gambling problems at age 17 (69). With respect to age of onset, one study found that early onset gambling disorder was associated with a more severe clinical presentation and with higher novelty seeking and lower self-directedness (70). In addition, gender appears to have an influence on impulsivity, as men with gambling problems may be more impulsive and score higher on measures of

Several researchers have attempted to categorize gambling disorder based on dimensions of personality, such as impulsivity, and co-occurring psychopathology. One study identified three subtypes of gambling disorder based on self-report questionnaires measuring impulsivity, depression, and anxiety (72). The first subtype consists of behaviorally conditioned gamblers, who develop gambling disorder through continual exposure to gambling and is the least severe type of gambling disorder. A second type, the emotionally vulnerable individual, has poor coping

271 skills, and gambles to regulate emotions. Third, antisocial impulsivity gamblers gamble to 272 regulate affect, but are also characterized by high rates of psychopathology and impulsivity. 273 274 Another study sought to categorize gamblers into four groups (73): Cluster 1 had high 275 impulsivity, rates of psychopathology, early onset, and severe gambling problems; Cluster 2 had 276 low sensation seeking and high avoidant, controlling, and distant behavior, with high rates of 277 alcohol abuse; Cluster 3 was characterized by high impulsivity and early onset, but also had high 278 rates of sensation seeking without psychopathological impairments; and Cluster 4 was defined by 279 low impulsivity and psychopathology, and a late age of onset. 280 281 In a meta-analysis of studies, significantly higher rates of several personality traits were 282 identified in people with gambling disorder compared to controls (medium-large effect sizes), 283 including negative urgency, low premeditation, unconscientious disinhibition (low 284 conscientiousness), negative affect, and disagreeable disinhibition (low agreeableness) (74). The 285 authors suggested that these findings in gambling disorder were similar to those observed in 286 substance use disorders, suggesting that it may be part of a broader group of conditions 287 characterized by externalizing psychopathology. 288 289 Some personality traits have been found to correlate with dopamine functioning. For example, in 290 healthy males, it was found that striatal dopamine receptor binding (measured using raclopride-291 PET) correlated with sensation-seeking according to an inverted 'U' shaped model (75). As 292 noted in section 2.2, dopamine receptor binding – again with raclopride-PET – was associated 293

with sensation-seeking across gambling disorder and control subjects (33).

Current research has just begun to examine how personality dimensions and disorders influence treatment outcome. One study found that treatment dropout was significantly related to impulsivity (76). Other studies have found that although certain personality aspects such as high novelty seeking have been associated with more severe gambling and a young age of gambling disorder onset, these variables were not associated with treatment outcome (70).

# 3.2. Gambling disorder and personality disorders

Personality disorders appear to be relatively common in people with gambling disorder, and are likely to contribute to chronic symptoms. In one study, 45.5% of individuals with gambling disorder met criteria for at least one personality disorder (76). However, the presence of a personality disorder was not clearly related with the severity of gambling symptoms.

There is evidence that rates of personality disorders in gambling disorder may be influenced by other psychiatric comorbidities. In a sample derived from a national survey, one or more personality disorders was evident in 71.4% of gambling disordered individuals with a comorbid anxiety disorder (versus 40.86% of low frequency gamblers with an anxiety disorder), and in 52.9% of gambling disordered individuals without a comorbid anxiety disorder (versus 11.3% of low frequency gamblers without an anxiety disorder) (77).

### 3.3. Gambling disorder and other potentially enduring traits

It is conceivable that some of the cognitive deficits that occur in Gambling disorder could represent enduring traits that predispose towards the development of symptoms. As such, cognitive measures may be useful as proxy 'personality measures' in that they may be enduring and more readily linked to underlying neurobiology than formal personality disorders or scores from personality questionnaires. In order to examine impulsivity at an endophenotypic level, cognitive research has attempted to delineate the complex construct of impulsivity observed in individuals with gambling disorder. Individuals with gambling disorder demonstrate deficiencies in planning, decision-making, motor inhibition, and cognitive flexibility (3). Perceived inability to stop gambling and positive gambling expectancies have also been associated with high school students, college students, and adults with gambling disorder (78). However, it is not known the extent to which these different deficits are trait in nature. To address this issue would require studies in unaffected first degree relatives and also, ideally, longitudinal studies capturing cognitive function before, during, and after the development of Gambling disorder. There is some evidence that decision-making deficits could represent a trait marker, based on findings in people at risk of gambling disorder but without fully developed pathological symptoms (22).

331

332

333

334

335

336

337

338

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

# 4. Conclusions

The literature suggests that gambling disorder is a heterogeneous condition; however, impulsivity appears to be characteristic of the majority of individuals with gambling disorder.

The relatively paucity of neuroimaging data (especially functional imaging), genetic studies, and translational studies from animals to humans in gambling disorder, however, limits our ability in defining gambling disorder as a deficit of a particular component(s) of the brain although

dysfunction in dopaminergic, glutamatergic, and serotonergic transmission have all been implicated. Further, the evidence of a genetic link between gambling disorder and other addictive behaviors is supported by high rates of familial transmission and the cross-beneficial efficacy of opioid antagonists and partial agonists in gambling and substance addiction. More holistic studies involving a number of research paradigms (genetics, cognition, imaging, etc) that explore the pathology of gambling disorder over time may be useful in furthering our understanding of the onset and course of gambling disorder.

# **Declaration of interest**

The authors declare that there are no competing financial interests in relation to the submitted work. No assistance was provided in the writing of this article. Dr. Grant has received research grant support from NIDA, NCRG, Psyadon Pharmaceuticals, Forest Pharmaceuticals, Roche Pharmaceuticals, and Transcept Pharmaceuticals. He has also received royalties from American Psychiatric Publishing Inc, Oxford University Press, Norton, and McGraw Hill Publishers. Mr. Odlaug has received research grant support from the Trichotillomania Learning Center, has consulted for Lundbeck Pharmaceuticals and has received royalties from Oxford University Press. Dr. Chamberlain has consulted for Cambridge Cognition.

# **Authors and contributors**

JEG drafted the first version of the manuscript. BLO and SRC critically revised the initial version of the manuscript for important intellectual content. All authors approve the final version of the manuscript.

# Acknowledgements

This research was supported by a Center for Excellence in Gambling Research grant by the National Center for Responsible Gaming to Dr. Grant, by and a research grant from the Trichotillomania Learning Center to Mr. Odlaug, and by a research grant from the Academy of Medical Sciences (UK) to Dr. Chamberlain.

#### 367 References

368 1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> 369 edition. Washington, D.C.: American Psychiatric Press (2013).

370

67an 2. Grant JE, Kim SW. Demographic and clinical features of 131 adult pathological gamblers. J 372 Clin Psychiatry (2001) **62**(12): 957-62.

373

374 3. Hodgins DC, Stea JN, Grant JE. Gambling disorders. *Lancet* (2011) **378**(9806): 1874-84.

375

376 4. Petry NM, Stinson FS, Grant BF. Comorbidity of DSM-IV pathological gambling and other 377 psychiatric disorders: results from the National Epidemiologic Survey on Alcohol and Related 378 Conditions. J Clin Psychiatry (2005) **66**(5): 564-74.

379

- 380 5. Shaffer HJ, Hall MN, Vander Bilt J. Estimating the prevalence of disordered gambling 381 behavior in the United States and Canada: a research synthesis. Am J Public Health (1999) **89**(9):
- 382 1369-76.

383

- 384 6. Shaffer, H. J., & Kidman, R. Gambling and the public health. In J. E. Grant & M. N. Potenza
- 385 (Eds.), Pathological gambling: A clinical guide to treatment. Washington, DC: American
- 386 Psychiatric Publishing, Inc. (2004).

387

388 7. Ashley LL, Boehlke KK. Pathological gambling: a general overview. J Psychoact 389 Drugs (2012) 44(1):27-37.

390

391 8. Conversano C, Marazziti D, Carmassi C, Baldini S, Barnabei G, Dell'Osso L. Pathological 392 gambling: a systematic review of biochemical, neuroimaging, and neuropsychological findings. 393 Harv Rev Psychiatry (2012) 20(3): 130-48.

394

395 9. Shaffer HJ, Martin R. Disordered gambling: etiology, trajectory, and clinical considerations. 396 Ann Rev Clin Psychol (2011) 7: 483-510.

397

- 398 10. el-Guebaly N, Mudry T, Zohar J, Tavares H, Potenza MN. Compulsive features in 399 behavioural
- 400 addictions: the case of pathological gambling. Addiction (2012) **107**(10): 1726-34. 401

402 11. van Holst RJ, van den Brink W, Veltman DJ, Goudriaan AE. Brain imaging studies in 403 pathological gambling. Current Psychiatr Rep (2010) 12(5): 418-25.

404

405 12. van Holst RJ, van den Brink W, Veltman DJ, Goudriaan AE. Why gamblers fail to win: a 406 review of cognitive and neuroimaging findings in pathological gambling. Neurosci Biobehav Rev 407 (2010) **34**(1): 87-107.

408

409 13. Chamberlain SR, Sahakian BJ. The neuropsychiatry of impulsivity. Curr Opin Psychiatry 410 (2007) **20**(3): 255-61.

- 412 14. Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in
- 413 adolescence: a critical period of addiction vulnerability. Am J Psychiatry (2003) 160(6): 1041-
- 414

- 416 15. Chamberlain SR, Menzies L, Hampshire A, Suckling J, Fineberg NA, del Campo N, Aitken
- 417 M, Craig K, Owen AM, Bullmore ET, Robbins TW, Sahakian BJ. Orbitofrontal dysfunction in
- 418 patients with obsessive-compulsive disorder and their unaffected relatives. Science (2008)
- 419 **321**(5887): 421-2.

420

421 16. Petry NM. Pathological gamblers, with and without substance use disorders, discount 422 delayed rewards at high rates. J Abnorm Psychol (2001) 110(3): 482-7.

423

424 17. Grant JE, Potenza MN. Compulsive aspects of impulse-control disorders. *Psychiatr Clin* 425 North Am (2006) **29**(2): 539-51

426

- 427 18. Blaszczynski A. Pathological gambling and obsessive compulsive spectrum disorders.
- 428 Psychol Rep (1999) 84: 107-13.

429

430 19. Frost RO, Meagher BM, Riskind JH. Obsessive compulsive features in pathological lottery 431 and stratch-ticket gamblers. J Gambl Stud (2001) 17: 5-19.

432

- 433 20. Goudriaan AE, Oosterlaan J, de Beurs E, van den Brink W. Decision making in pathological 434 gambling: a comparison between pathological gamblers, alcohol dependents, persons with
- 435 Tourette syndrome, and normal controls. Brain Res Cogn Brain Res (2005) 23(1): 137-51.

436

- 437 21. Odlaug BL, Chamberlain SR, Kim SW, Schreiber LR, Grant JE. A neurocognitive
- 438 comparison of cognitive flexibility and response inhibition in gamblers with varying degrees of 439 clinical severity. *Psychol Med* (2011) **41**(10): 2111-9.

440

441 22. Grant JE, Chamberlain SR, Schreiber LR, Odlaug BL, Kim SW. Selective decision-making 442 deficits in at-risk gamblers. Psychiatry Res (2011) 189(1): 115-20.

443

- 444 23. Potenza MN, Leung HC, Blumberg HP, Peterson BS, Fulbright RK, Lacadie CM, Skudlarski
- 445 P, Gore JC. An FMRI Stroop task study of ventromedial prefrontal cortical function in
- 446 pathological gamblers. Am J Psychiatry (2003) 160: 1990-4.

447

- 448 24. de Ruiter MB, Veltman DJ, Goudriaan AE, Oosterlaan J, Sjoerds Z, van den Brink W.
- 449 Response perseveration and ventral prefrontal sensitivity to reward and punishment in male
- 450 problem gamblers and smokers. *Neuropsychopharmacology* (2009) **34**(4): 1027-38.

451

- 452 25. Goudriaan AE, de Ruiter MB, van den Brink W, Oosterlaan J, Veltman DJ. Brain activation
- 453 patterns associated with cue reactivity and craving in abstinent problem gamblers, heavy
- 454 smokers and healthy controls: an fMRI study. Addict Biol (2010) 15(4): 491-503.

- 456 26. Miedl SF, Peters J, Büchel C. Altered neural reward representations in pathological gamblers
- 457 revealed by delay and probability discounting. Arch Gen Psychiatry (2012) 69(2): 177-86.

- 459 27. Tschernegg M, Crone JS, Eigenberger T, Schwartenbeck P, Fauth-Buhler M, Lemenager T,
- et al., Abnormalities of functional brain networks in pathological gambling: a graph-theoretical
- 461 approach. Front Hum Neurosci (2013); 7:625. doi: 10.3389/fnhum.2013.00625.

462

- 28. Koehler S, Hasselmann E, Wüstenberg T, Heinz A, Romanczuk-Seiferth N. Higher volume
- of ventral striatum and right prefrontal cortex in pathological gambling. Brain Struct Funct. 2013
- Nov 16. [Epub ahead of print]

466

29. Robbins TW, Everitt BJ, Nutt DJ. Introduction. The neurobiology of drug addiction: new vistas. *Philos Trans R Soc Lond B Biol Sci.* 2008 Oct 12;363(1507):3109-11.

469

- 470 30. Volkow ND, Baler RD. Addiction science: Uncovering neurobiological complexity.
- 471 *Neuropharmacology*. 2014 Jan;76 Pt B:235-49.

472

- 473 31. Boileau I, Payer D, Chugani B, Lobo D, Behzadi A, Rusjan PM, Houle S, Wilson AA, Warsh
- J, Kish SJ, Zack M. The D2/3 dopamine receptor in pathological gambling: a positron emission
- tomography study with [11C]-(+)-propyl-hexahydro-naphtho-oxazin and [11C]raclopride.
- 476 Addiction. 2013 May;108(5):953-63.

477

- 478 32. Clark L, Stokes PR, Wu K, Michalczuk R, Benecke A, Watson BJ, Egerton A, Piccini P,
- Nutt = DJ, Bowden-Jones H, Lingford-Hughes AR Striatal dopamine  $D_2/D_3$  receptor binding in
- pathological gambling is correlated with mood-related impulsivity. *Neuroimage*. 2012 Oct
- 481 15;63(1):40-6.

482

- 483 33. Peterson E, Møller A, Doudet DJ, Bailey CJ, Hansen KV, Rodell A, Linnet J, Gjedde A.
- 484 Pathological gambling: relation of skin conductance response to dopaminergic neurotransmission
- and sensation-seeking. Eur Neuropsychopharmacol. 2010 Nov;20(11):766-75.

486

- 487 34. Linnet, J., Mouridsen, K., Peterson, E., Møller, A., Doudet, D., & Giedde, A. (2012). Striatal
- 488 Dopamine Release Codes Uncertainty in Pathological Gambling. *Psychiatry Research*:
- 489 *Neuroimaging*, 204, 55-60.

490

35. Leeman RF, Potenza MN. A targeted review of the neurobiology and genetics of behavioural addictions: an emerging area of research. *Can J Psychiatry* (2013) **58**(5): 260-73.

493

- 494 36. Black DW, Monahan PO, Temkit M, Shaw M. A family study of pathological gambling.
- 495 *Psychiatry Res* (2006) **141**(3): 295-303.

496

37. Lobo DS, Kennedy JL. Genetic aspects of pathological gambling: a complex disorder with shared genetic vulnerabilities. *Addiction* (2009) **104**(9): 1454-65.

- 38. Slutske WS, Ellingson JM, Richmond-Rakerd LS, Zhu G, Martin NG. Shared genetic
- vulnerability for disordered gambling and alcohol use disorder in men and women: evidence
- from a national community-based Australian Twin Study. Twin Res Hum Genet (2013) **16**(2):
- 503 525-34.

- 39. Gyollai A, Griffiths MD, Barta C, Vereczkei A, Urbán R, Kun B, Kökönyei G, Székely A,
- Sasvári-Székely M, Blum K, Demetrovics Z. The Genetics of Problem and Pathological
- Gambling: A Systematic Review. *Curr Pharm Des* (2013) [Epub ahead of print].

508

- 509 40. Goudriaan AE, Oosterlaan J, de Beurs E, Van den Brink W. Pathological gambling: a
- 510 comprehensive review of biobehavioral findings. *Neurosci Biobehav Rev* (2004) **28**(2):123-41

511

- 512 41. Nussbaum D, Honarmand K, Govoni R, Kalahani-Bargis M, Bass S, Ni X, Laforge K,
- Burden A, Romero K, Basarke S, Courbasson C, Deamond W. An eight component decision-
- making model for problem gambling: a systems approach to stimulate integrative research. J
- 515 *Gambl Stud* (2011) **27**(4): 523-63.

516

- 517 42. Ye Z, Hammer A, Camara E, Münte TF. Pramipexole modulates the neural network of
- 518 reward anticipation. *Hum Brain Mapp* (2011) **32**(5):800-11.

519

- 43. Abler B, Hahlbrock R, Unrath A, Grön G, Kassubek J. At-risk for pathological gambling:
- imaging neural reward processing under chronic dopamine agonists. *Brain* (2009) **132**(Pt 9):
- 522 2396-402.

523

- 524 44. Kalivas PW. The glutamate homeostasis hypothesis of addiction. *Nat Rev Neurosci* (2009)
- **10**(8): 561-72.

526

- 527 45. Kalivas PW, Volkow ND. New medications for drug addiction hiding in glutamatergic
- 528 neuroplasticity. *Mol Psychiatry* (2011) **16**(10): 974-86.

529

- 530 46. Nordin C, Gupta RC, Sjödin I. Cerebrospinal fluid amino acids in pathological gamblers and
- 531 healthy controls. *Neuropsychobiology* (2007) **56**(2-3): 152-8.

532

- 533 47. Koot S, Zoratto F, Cassano T, Colangeli R, Laviola G, van den Bos R, Adriani W.
- 534 Compromised decision-making and increased gambling proneness following dietary serotonin
- 535 depletion in rats. *Neuropharmacology* (2012) **62**(4): 1640-50.

536

- 48. Zeeb FD, Robbins TW, Winstanley CA. Serotonergic and dopaminergic modulation of
- gambling behavior as assessed using a novel rat gambling task. *Neuropsychopharmacology*
- 539 (2009) **34**(10): 2329-43.

540

- 49. Ibáñez A, Blanco C, Perez de Castro I, Fernandez-Piqueras J, Sáiz-Ruiz J. Genetics of
- 542 pathological gambling. *J Gambl Stud* (2003) **19**(1): 11-22.

543

- 50. Wilson D, da Silva Lobo DS, Tavares H, Gentil V, Vallada H. Family-based association
- analysis of serotonin genes in pathological gambling disorder: evidence of vulnerability risk in
- 546 the 5HT-2A receptor gene. *J Mol Neurosci* (2013) **49**(3): 550-3.

- 51. Aston-Jones G, Cohen JD. Adaptive gain and the role of the locus coeruleus-norepinephrine
- system in optimal performance. J Comp Neurol (2005) 493(1): 99-110.

52. Bouret S, Sara SJ. Network reset: a simplified overarching theory of locus coeruleus noradrenaline function. Trends Neurosci (2005) 28(11): 574-82.

53. Baarendse PJ, Vanderschuren LJ. Dissociable effects of monoamine reuptake inhibitors on

54. Fernando AB, Economidou D, Theobald DE, Zou MF, Newman AH, Spoelder M, Caprioli

distinct forms of impulsive behavior in rats. *Psychopharmacology (Berl)* (2012) **219**(2): 313-26.

D, Moreno M, Hipólito L, Aspinall AT, Robbins TW, Dalley JW. Modulation of high

impulsivity and attentional performance in rats by selective direct and indirect dopaminergic and noradrenergic receptor agonists. *Psychopharmacology (Berl)* 2012 **219**(2): 341-52.

- 55. Roy A, DeJong J, Ferraro T, Adinoff B, Gold P, Rubinow D, Linnoila M. CSF GABA and neuropeptides in pathological gamblers and normal controls. Psychiatry Res (1989) 30(2): 137-
- 44.

- 56. Pallanti S, Bernardi S, Allen A, Chaplin W, Watner D, DeCaria CM, Hollander E.
- Noradrenergic function in pathological gambling: blunted growth hormone response to
- clonidine. J Psychopharmacol (2010) 24(6): 847-53.

- 57. Peciña S, Smith KS, Berridge KC. Hedonic hot spots in the brain. *Neuroscientist* (2006)
- (6): 500-11.

- 58. Barbano MF, Cador M. Opioids for hedonic experience and dopamine to get ready for it.
- Psychopharmacology (Berl) (2007) **191**(3): 497-506.

- 59. Petrovic P, Pleger B, Seymour B, Klöppel S, De Martino B, Critchley H, Dolan RJ. Blocking
- central opiate function modulates hedonic impact and anterior cingulate response to rewards and
- losses. J Neurosci (2008) 28(42): 10509-16.

- 60. Shinohara K, Yanagisawa A, Kagota Y, Gomi A, Nemoto K, Moriya E, Furusawa E, Furuya
- K, Terasawa K. Physiological changes in Pachinko players; beta-endorphin, catecholamines,
- immune system substances and heart rate. Appl Human Sci (1999) 18(2): 37-42.

- 61. Grant JE, Kim SW, Hartman BK. A double-blind, placebo-controlled study of the opiate
- antagonist naltrexone in the treatment of pathological gambling urges. J Clin Psychiatry (2008)
- : 783-9.

- 62. Grant JE, Odlaug BL, Potenza MN, Hollander E, Kim SW. Nalmefene in the treatment of
- pathological gambling: multicentre, double-blind, placebo-controlled study. Br J Psychiatry
- (2010) **197**: 330-1.

- 63. Grant JE, Potenza MN, Hollander E, Cunningham-Williams R, Nurminen T, Smits G, Kallio
- A. Multicenter investigation of the opioid antagonist nalmefene in the treatment of pathological

594 gambling. *Am J Psychiatry* (2006) **163**: 303-12.

595

596 64. Gottesman, II, Gould TD. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry* (2003) **160**(4): 636-45.

598

599 65. Chamberlain SR, Menzies L. Endophenotypes of obsessive-compulsive disorder: rationale, 600 evidence and future potential. *Exp Rev Neurother* (2009) **9**(8): 1133-46.

601

- 602 66. Clark L. "Epidemiology and phenomenology of pathological gambling," in Grant JE,
- Potenza MN, editors. *The Oxford Handbook of Impulse Control Disorders*. New York, Oxford University Press (2012). p. 94-116.

605

606 67. Lai FD, Ip AK, Lee TM. Impulsivity and pathological gambling: is it a state or a trait problem. *BMC Res Notes* (2011) **13**: 492.

608

609 68. Auger N, Lo E, Cantinotti M, O'Loughlin J. Impulsivity and socio-economic status interact to increase the risk of gambling onset among youth. *Addiction* (2010) **105**; 2176-83.

611

69. Dussault F, Brendgen M, Vitaro F, Wanner B, Tremblay RE. Longitudinal links between impulsivity, gambling problems and depressive symptoms: a transactional model from adolescence to early adulthood. *J Child Psychol Psychiatry* (2011) **52**: 130-8.

615

- 70. Jiménez-Murcia S, Alvarez-Moya EM, Stinchfield R, Fernández-Aranda F, Granero R,
   Aymamí N, Gómez-Peña M, Jaurrieta N, Bove F, Menchón JM. Age of onset in pathological
- gambling: clinical, therapeutic and personality correlates. *J Gambl Stud* (2010) **26**: 235-48.

619

71. Echeburúa E, González-Ortega I, de Corral P, Polo-López R. Clinical gender differences among adult pathological gamblers seeking treatment. *J Gambl Stud* (2011) **27**(2): 215-27.

622

72. Ledgerwood DM, Petry NM. Subtyping pathological gamblers based on impulsivity, depression, and anxiety. *Psychol Addict Behav* (2010) **24**: 680-8.

625

- 626 73. Álvarez-Moya EM, Ochoa C, Jiménez-Murcia S, Aymamí MN, Gómez-Peña M, Fernández-
- Aranda F, Santamaría J, Moragas L, Bove F, Menchón JM. Effect of executive functioning,
- decision-making and self-reported impulsivity on the treatment outcome of pathological
- 629 gambling. J Psychiatry Neurosci (2011) **36**: 165-75.

630

74. Maclaren VV, Fugelsang JA, Harrigan KA, Dixon MJ. The personality of pathological gamblers: a meta-analysis. *Clin Psychol Rev* (2011) **31**(6):1057-67.

633

- 75. Gjedde A, Kumakura Y, Cumming P, Linnet J, Møller A. Inverted-U-shaped correlation
- between dopamine receptor availability in striatum and sensation seeking. Proc Natl Acad Sci U
- 636 S A. 2010 Feb 23;107(8):3870-5.
- 76. Odlaug BL, Schreiber LR, Grant JE. Personality disorders and dimensions in pathological
- 638 gambling. <u>J Pers Disord.</u> 2012 Apr 11. [Epub ahead of print]

- 77. Giddens JL, Stefanovics E, Pilver CE, Desai R, Potenza MN. Pathological gambling severity
   and co-occurring psychiatric disorders in individuals with and without anxiety disorders in a
   nationally representative sample. *Psychiatry Res* (2012);199(1):58-64.
- 78. Tang CS, Wu AM. Gambling-related cognitive biases and pathological gambling among youths, young adults, and mature adults in Chinese societies. *J Gambl Stud* (2012) **28**(1):139-54.