

# Local adaptation and dispersal evolution interact to drive population response to climate change

Christopher Weiss-Lehman<sup>1</sup> and Allison Shaw<sup>1</sup>

<sup>1</sup>Ecology, Evolution, and Behavior, University of Minnesota

## 1 Introduction

Climate change is expected to dramatically reshape global biogeographic patterns as some species shift their ranges to track changing environmental conditions (Gonzalez et al., 2010; Peñuelas and Boada, 2003; Hansen et al., 2001; Scholze et al., 2006). These range shifts are generally expected to proceed upwards in latitude, elevation, or both as average global temperatures continue to rise (Loarie et al., 2009). In fact, contemporary range shifts have already been observed in a wide range of taxa (Chen et al., 2011; Walther et al., 2002; Parmesan and Yohe, 2003; Parmesan, 2006; Parmesan et al., 1999; Perry et al., 2005). Such range shifts present significant challenges to current and future conservation efforts as they can result in the extinction of populations failing to track a changing climate (Davis and Shaw, 2001; Parmesan, 2006; Zhu et al., 2012; Sekercioglu et al., 2008), the creation of novel species assemblages as not all species shift their ranges at the same rate or at all (Hobbs et al., 2009; Gilman et al., 2010; Williams and Jackson, 2007), or both. It is therefore crucial to understand the dynamics of such climate induced range shifts to better inform current and future conservation work.

While contemporary range shifts are a relatively new phenomenon, important insights can be gained from the related but distinct process of range expansion. Range expansions have been studied for decades, leading to a robust understanding of both the ecological (Skellam, 1951; Hastings et al., 2005) and evolutionary (Shine et al., 2011; Burton et al., 2010; Excoffier et al., 2009; Kubisch et al., 2014) mechanisms responsible for shaping such expansions. For example, while the speed of a

range expansion can be well approximated by a combination of the species' intrinsic growth rate and dispersal ability (Hastings et al., 2005), recent research demonstrates that evolution in both of these traits can have important implications for both the mean and variance of expansion speed through time (Weiss-Lehman et al., 2017; Ochocki and Miller, 2017; Szűcs et al., 2017; Phillips, 2015). As range shifts also involve a leading edge of population advance, they are likely to be affected by similar ecological and evolutionary mechanisms as have been shown to drive dynamics in range expansions (Hargreaves and Eckert, 2014). However, while such insights from range expansions are valuable for understanding and predicting dynamics of range shifts, it is important to recognize that these two processes have significant differences as well.

In particular, range expansions and range shifts due to climate change involve fundamentally distinct initial conditions. Range expansions typically begin from the successful establishment and subsequent spread of a small, founding population as typically occurs with the range expansions of invasive and reintroduced species (Hastings et al., 2005). Such founding populations represent samples from some larger source population and as such lack any initial spatial population structure. In contrast, range shifts involve entire populations with existing spatial structure within the initially stable range (Hargreaves and Eckert, 2014). Such spatial structure can take the form of local adaptation within the range, degree of transition in population size from range core to edge, or some combination of the two (Hargreaves and Eckert, 2014; Hargreaves et al., 2015; Henry et al., 2013).

These different aspects of spatial population structure have been shown to dramatically impact the response of populations to climate change. For example, the nature of the gradient forming the range edge (e.g. declines in birth rates vs. increases in extinction risk) can directly change the probability of extinction a species faces during a climate driven range shift (Henry et al., 2013). A population's risk of extinction during a range shift has also been related to local adaptation within the range. Specifically, a broad environmental niche (i.e. little local adaptation) can decrease a population's ability to track a changing climate if dispersal occurs in a stepping stone manner, allowing some individuals to block dispersal of better adapted phenotypes (Atkins and Travis, 2010). Local adaptation also has the potential to interact with dispersal evolution during climate change,

driving increased dispersal evolution in an asexual species (Hargreaves et al., 2015), though it is unclear if this result will hold for sexually reproducing species in which locally adapted genotypes interact via gene flow. As the gradient forming the range edge can influence the distribution of dispersal phenotypes within the range (Henry et al., 2013; Hargreaves and Eckert, 2014), it is likely that this would also result in an altered interaction between dispersal evolution and local adaptation in sexually reproducing species. In fact, pollen dispersal in flowering plants has been shown to dramatically reduce the likelihood of a range shift in response to climate change (Aguilée et al., 2016). Evolution during range shifts due to local adaptation and dispersal evolution are both potentially important drivers of range shift dynamics (Van Petegem et al., 2016) and it is therefore important to consider both when predicting the dynamics of populations responding to climate change.

Here, we use a complex, individual-based model incorporating evolution in both fitness and dispersal distance to determine the interaction between local adaptation and the starkness of the gradient at the range edge in driving population dynamics during climate change for a sexually reproducing species. Using this model, we simultaneously vary the strength of local adaptation, the starkness of the range boundary, and the speed of climate change to ascertain the relationship among them and test how dispersal evolution interacts with local adaptation to help or hinder a population’s ability to track a changing climate. We additionally compare the dynamics of successful versus doomed simulated populations to understand the factors contributing to population extinction and identify warning signs for populations in danger of extinction as a result of a range shift.

## 2 Methods

### 2.1 Overview

#### 2.1.1 Purpose

This model tests an evolving population’s ability to track a changing climate under a variety of conditions. Specifically, populations are simulated with (1) stark or gradual range boundaries, (2)

strong or weak local selection pressures, and (3) varying speeds of climate change. In all simulations, individual dispersal and relative fitness are defined by an explicit set of quantitative diploid loci subject to mutation, thus allowing both traits to evolve over time. All simulations begin with stable climate conditions for 2000 generations to allow the populations to reach a spatial equilibrium before the onset of climate change. Climate change is then modeled as a constant, directional shift in the location of environmentally suitable habitat (see the *Submodels* section below). Finally, simulations end with another short period of climate stability to assess the population’s ability to persist and recover after shifting its range.

### 2.1.2 State variables and scales

The model simulates a population of male and female individuals characterized by spatial coordinates for their location and diploid loci for both fitness and dispersal. Space is modeled as a lattice of discrete patches overlaying a continuous Cartesian coordinate system with a fixed width along the y axis and without bounds on the x axis. To avoid edge effects along the y axis, the model employs wrapping boundaries such that if an individual disperses out of the landscape on one side, it appears at the opposite end of the same row of the landscape. Patches are defined by the location of the patch center in x and y coordinates and a patch width parameter defining the relationship between continuous Cartesian space and the discrete patches used for population dynamics (see the *Submodels* section below).

The abiotic environment is defined by a parameter denoting the center of the range, the slope of the range boundary, and the width of the range (See Figure 1). The range can then shift in space by altering the parameter defining the range center, which is how climate change is implemented in the model. Further, a gradient of phenotypic optimum values for fitness is imposed within the range to allow for local adaptation within the range boundaries. The severity of this gradient is defined by a separate parameter independent of the other environmental parameters (greater values of this parameter will lead to steeper gradients in phenotypic optima within the range).




Figure 1: Environmental quality, as defined by  $f(x, t)$ , over a spatial gradient. The parameters of  $f(x, t)$  are shown on the figure at significant points along the  $x$  axis. Additionally, the lattice of discrete patches in which population dynamics occur is shown beneath. This lattice of  $\eta \times \eta$  patches are mapped to environmental conditions along the  $x$  axis using  $f(x, t)$  and are environmentally constant along the  $y$  axis. Dispersal is unbounded in the  $x$  direction and implemented with wrapping boundaries in the  $y$  direction.

### 2.1.3 Process overview and scheduling

Time is also modeled in discrete intervals defining single generations of the population. Within each generation, individuals first reproduce within their natal patch. After reproduction, all adults die and the offspring disperse according to their dispersal kernels (*Submodels*), resulting in discrete, non-overlapping generations (See Figure 2). Reproduction occurs according to a stochastic implementation of the classic Ricker model (Ricker, 1954) taking into account the mean fitness of individuals within the patch (*Submodels*). Parental pairs form via random sampling of the local population (with replacement) weighted by individual fitness such that individuals with high fitness are likely to produce multiple offspring while individuals with no fitness may not produce any. Individuals inherit one allele from each parent at each loci, assuming independent segregation and a mutation process (*Submodels*). After reproduction, all individuals in the parental generation perish, forming discrete non-overlapping generations. The offspring then disperse according to their dispersal traits to start the next generation in their post dispersal locations.

## 2.2 Design concepts

### 2.2.1 Emergence

Emergent phenomena in this model include the spatial equilibrium of population abundances and trait values within the stable range, the demographic dynamics of the shifting population during climate change, and the evolutionary trajectories of both fitness and dispersal traits during climate change. These are all examined in the context of their impact on the population’s ability to keep pace with the changing climate.

### 2.2.2 Stochasticity

All biological processes in this model are stochastic including realized population growth in each patch, dispersal distances of each individual, and inheritance of loci. Environmental parameters are fixed, however, and the process of climate change (i.e. the movement of environmentally suitable habitat through time) is deterministic. Thus, the model removes the confounding influence of environmental stochasticity to focus on demographic and evolutionary dynamics of range shifts.




Figure 2: The life cycle of simulated populations is shown divided between events contributing to reproduction and dispersal. Each generation begins with new offspring dispersing according to their phenotype, after which reproduction occurs in local populations defined by the discrete lattice. After reproduction, all parental individuals perish, resulting in discrete, non-overlapping generations.

### 2.2.3 Interactions

Individuals in the model interact via mating and density-dependent competition within patches. Other important interactions are the relationship between dispersal evolution and local adaption, particularly in edge populations, and how this relationship impacts a population's ability to avoid extinction and track a changing climate. These interactions are examined in the context of different configurations of the initial, stable range and different speeds of climate change.

### 2.2.4 Desired output

After each model run, full details of all surviving individuals at the last time point are recorded (spatial coordinates and loci values for both traits). If a population went extinct during the model run, the time of extinction is recorded. Throughout the simulation, certain aggregated values are calculated and recorded for each occupied patch including the population size and the mean and variance of allele values for each trait.

## 2.3 Details

### 2.3.1 Initialization

The following parameters are set at the beginning of each simulation and form the initial conditions of the model: the mean and variance for allele values of each trait, population size, location of environmentally suitable habitat, number of generations for the pre-, post-, and during climate change periods of the simulation, and all other necessary parameters for the submodels defined below. Simulated populations are initialized in the center of the range and allowed to spread and equilibrate throughout the range during the period of stable climate conditions. This ensures that the populations reacting to a changing climate truly represent the expected spatial distribution for a given range, rather than the initial parameter values used in the simulation. See Table 1 for a full list of all parameter values used in the simulations described here.



### 2.3.2 Submodels

**Environmentally suitable habitat** Environmentally suitable habitat is determined by the population's carrying capacity as it ranges in space ( $K_x$ ). The maximum achievable carrying capacity ( $K_{max}$ ) occurs in the center of the species' range and declines with increasing distance from the center. Specifically, the carrying capacity at a location  $x$  is defined as a combination of  $K_{max}$  and a function  $f(x, t)$ , where  $f(x, t)$  ranges from 1 in the range center to 0 far away from the center and is defined as follows

$$f(x, t) = \begin{cases} \frac{e^{\gamma(x-\beta_t+\tau)}}{1+e^{\gamma(x-\beta_t+\tau)}} & x \leq \beta_t \\ \frac{e^{-\gamma(x-\beta_t-\tau)}}{1+e^{-\gamma(x-\beta_t-\tau)}} & x > \beta_t \end{cases} \quad (1)$$

where  $\beta_t$  defines the center of the area of suitable habitat at time  $t$ ,  $\tau$  sets the width of the range, and  $\gamma$  affects the slope of the function at the range boundaries (See Figure 1). To understand the relationship between  $\gamma$  and the slope of  $f(x, t)$  at the range boundary, the partial derivative of  $f(x, t)$  over the spatial dimension can be shown to be

$$f(x, t) = \begin{cases} \frac{e^{\gamma(x-\beta_t+\tau)}}{1+e^{\gamma(x-\beta_t+\tau)}} & x \leq \beta_t \\ \frac{e^{-\gamma(x-\beta_t-\tau)}}{1+e^{-\gamma(x-\beta_t-\tau)}} & x > \beta_t \end{cases} \quad (2)$$

yielding a derivative of  $\pm \frac{\gamma}{4}$  at the inflection points on either side of the range center ( $x = \beta_t \pm \tau$ ).

Population dynamics occur within discrete patches, so to calculate a  $K_x$  value for a discrete patch from the continuous function  $f(x, t)$ , we use another parameter defining the spatial scale of each patch ( $\eta$ ; See Figure 1). The local carrying capacity of a patch centered on  $x$  ( $K_x$ ) is then calculated as the mean of  $f(x, t)$  over the interval of the patch multiplied by  $K_{max}$ .

$$K_x = \frac{K_{max}}{\eta} \int_{x-\frac{\eta}{2}}^{x+\frac{\eta}{2}} f(x, t) dx \quad (3)$$

By varying the parameters defining  $f(x, t)$ , we can change both the total achievable carrying capacity of the population throughout the range (by altering both  $\tau$  and  $\gamma$ ) and the slope at which  $K_x$  declines to 0 (by altering  $\gamma$ ). We are primarily interested in the effect of altering the slope at

167 which  $K_x$  declines at the range boundaries, however as doing so can also alter the total achievable  
 168 carrying capacity of the population, we control for this confounding factor by fixing the total area  
 169 under the curve  $f(x, t)$ . The indefinite integral of  $f(x, t)$  can be shown to be

$$\int_{-\infty}^{\infty} f(x, t) dx = \frac{2 \ln(e^{\gamma \tau} + 1)}{\gamma} \quad (4)$$

170 which can be solved for  $\tau$ . Thus, if the total area under the curve is fixed, an appropriate value of  
 171  $\tau$  can be calculated for each value of  $\gamma$ .

172 Thus,  $\gamma$  and  $\tau$  are both fixed within a given simulation and  $\beta_t$  (the location of the center of  
 173 suitable habitat) is used to simulate climate change. During the periods before and after climate  
 174 change  $\beta_t$  is constant, but to simulate climate change it varies with time as follows

$$\beta_t = \nu \eta (t - \hat{t}) \quad (5)$$

175 where  $\nu$  is the velocity of climate change per generation in terms of discrete patches,  $\eta$  is the  
 176 spatial scale of each patch,  $t$  is the current generation, and  $\hat{t}$  is the last generation of stable climatic  
 177 conditions before the onset of climate change.

178 **Local adaptation** To allow an arbitrary degree of local adaptation within the range, the local  
 179 phenotypic optima ( $z_{opt,x}$ ) is set as follows

$$z_{opt,x} = \lambda (x - \beta_t) \quad (6)$$

180 where  $\lambda$  defines the strength of local selection with values close to 0 resulting in little to no change in  
 181 phenotypic optimum across the range and values of greater magnitude resulting in large differences  
 182 in phenotypic optima across the range. Individual fitness ( $w_{i,x}$ ) values are then calculated according  
 183 to the following equation assuming stabilizing selection

$$w_{i,x} = e^{\frac{-(z_i - z_{opt,x})^2}{2\omega^2}} \quad (7)$$

where  $\omega$  defines the strength of stabilizing selection and  $z_i$  is an individual's fitness phenotype (Lande, 1976). All loci are assumed to contribute additively to the phenotype with no dominance or epistasis, meaning an individual's phenotype is simply the sum of the individual's allele values for the fitness trait.

**Population dynamics** Population growth within each patch is modeled with a stochastic implementation of the classic Ricker model (Ricker, 1954; Melbourne and Hastings, 2008). To account for fitness effects on population growth, expected population growth is scaled by the mean relative fitness of individuals within the patch ( $\bar{w}_x$ ). The expected number of new offspring in patch  $x$  at time  $t + 1$  is then given by

$$\hat{N}_{t+1,x} = \bar{w}_x F_{t,x} \frac{R}{\psi} e^{\frac{-RN_{t,x}}{K_x}} \quad (8)$$

where  $F_{t,x}$  is the number of females in patch  $x$  at time  $t$ ,  $R$  is the intrinsic growth rate for the population,  $\psi$  is the expected sex ratio of the population,  $N_{t,x}$  is the number of individuals (males and females) in patch  $x$  at time  $t$ , and  $K_x$  is the local carrying capacity based on the environmental conditions. To incorporate demographic stochasticity, the realized number of offspring for each patch is then drawn from a Poisson distribution.

$$N_{t+1,x} \sim \text{Poisson}(\hat{N}_{t+1,x}) \quad (9)$$

Parentage of the offspring is then assigned by random sampling of the local male and female population. The sampling is weighted by individual fitness and occurs with replacement so highly fit individuals are likely to have multiple offspring while low fitness individuals may not have any. Each offspring inherits one allele per locus from each parent, assuming no linkage among loci. After reproduction, all members of the previous generation die and the offspring disperse to begin the next generation.

**Mutation** Inherited alleles are subject to mutation such that some offspring might not inherit identical copies of certain alleles from their parents. The mutation process is defined by two parameters for each trait  $T$ : the diploid mutation rate ( $U^T$ ) and the mutational variance ( $V_m^T$ ).

207 Using these parameters along with the number of loci defining trait  $T$  ( $L^T$ ), the per locus probability  
 208 of a mutation is

$$\frac{U^T}{2L^T} \quad (10)$$

209 Mutational effects are drawn from a normal distribution with mean 0 and a standard deviation of

$$\sqrt{V_m^T U^T} \quad (11)$$

210 By defining the mutation process in this manner rather than setting a probability of mutation and  
 211 mutational effect directly, similar mutational dynamics can be imposed regardless of the number  
 212 of loci used in the simulation.

213 **Dispersal** Finally, individuals disperse according to an exponential dispersal kernel defined by  
 214 each individual's dispersal phenotype. An individual's dispersal phenotype is the expected dispersal  
 215 distance and is given by

$$d_i = \frac{D\eta e^{\rho \Sigma L^D}}{1 + e^{\rho \Sigma L^D}} \quad (12)$$

216 where  $D$  is the maximum expected dispersal distance in terms of discrete patches,  $\eta$  is the spatial  
 217 scale of discrete patches,  $\rho$  is a constant determining the slope of the transition between 0 and  $D$ ,  
 218 and the summation is the sum of all alleles contributing to dispersal. Thus, as with fitness, loci are  
 219 assumed to contribute additively with no dominance or epistasis. The expected dispersal distance,  
 220  $d_i$  is then used to draw a realized distance from an exponential dispersal kernel. The direction  
 221 of dispersal is drawn from a uniform distribution bounded by 0 and  $2\pi$ . If a dispersal trajectory  
 222 takes an individual outside the bounds of the landscape in the  $y$  axis, the individual reappears  
 223 at the same  $x$  coordinate but the opposite end of the  $y$  axis, thus wrapping the top and bottom  
 224 edges of the landscape to avoid edge effects. Dispersal occurs from the center of each patch and  
 225 the individual's new patch is then determined according to its location in the overlaid grid of  $\eta$  x  
 226  $\eta$  patches (see Figure 1).

Parameter	Description	Value
$N_1$	Initial population size (seeded across multiple patches) when beginning the simulations	2500
$\beta_1$	Center of environmentally suitable habitat before climate change	0
$\eta$	Spatial dimensions of habitat patches in continuous space	50
$y_{max}$	Number of patches the discrete lattice extends in the $y$ direction	10
$\hat{t}$	Last time point of stable climate conditions	2000
$t_\Delta$	Duration of climate change	100
$t_{max}$	Total number of time points in the simulation	2150
$R$	Intrinsic growth rate of the population	2
$K_{max}$	Maximum achievable carrying capacity in the range	100
$\psi$	Expected sex ratio in the population	0.5
$D$	Maximum achievable dispersal phenotype	1000
$\rho$	Determines the slope of the transition in dispersal phenotypes from 0 to $D$	0.5
$\omega$	Defines the strength of stabilizing selection on fitness traits	3
$U^T$	Diploid mutation rate for each trait	0.02 for each trait
$V_m^T$	Mutational variance for each trait	0.0004 for each trait
$L^T$	The number of diploid loci defining each trait	5 for each trait
$\mu_1^f$	The initial mean allele value for the fitness trait	0
$\mu_1^d$	The initial mean allele value for the dispersal trait	-1
$\sigma_1^f$	The initial standard deviation of allele values for fitness	0.025
$\sigma_1^d$	The initial standard deviation of allele values for dispersal	1

Table 1: Simulation parameters held constant across all scenarios presented here.

Habitat gradient at range edge	Strength of local adaptation	$\gamma$	$\tau$	$\lambda$
Shallow	None	0.0025	250	0
	Moderate	0.0025	250	0.004
	Strong	0.0025	250	0.008
Moderate	None	0.025	421.479	0
	Moderate	0.025	421.479	0.004
	Strong	0.025	421.479	0.008
Stark	None	0.25	421.48	0
	Moderate	0.25	421.48	0.004
	Strong	0.25	421.48	0.008

Table 2: Descriptions and parameter values for the 9 different scenarios examined here. For each scenario,  $\nu$  (the speed of climate change) was varied among 0.5, 1, and 2.

## 2.4 Simulation descriptions

Most of the parameters described above were held constant over all simulations we performed (Table 1). We varied certain parameter values to explore the interacting roles of local adaptation, the habitat gradient at the range edge, and the speed of climate change in the dynamics of range shifts. Specifically, we considered a factorial combination of three severities of habitat quality decline at the range edge, three degrees of local adaptation within the range, and three speeds of climate change (Table 2) for a total of 27 different model scenarios. For each scenario, we performed 100 simulations to assess the interacting ecological and evolutionary dynamics of climate driven range shifts. All simulations were performed in R version 3.4.4 (Team, 2000) and the code is available at <https://github.com/tpweiss06/ShiftingSlopes>.

## References

Aguilée, R., G. Raoul, F. Rousset, and O. Ronce. 2016. Pollen dispersal slows geographical range shift and accelerates ecological niche shift under climate change. *Proceedings of the National Academy of Sciences* 113:E5741–E5748.

- Atkins, K., and J. Travis. 2010. Local adaptation and the evolution of species? ranges under climate change. *Journal of Theoretical Biology* 266:449–457.
- Burton, O. J., B. L. Phillips, and J. M. Travis. 2010. Trade-offs and the evolution of life-histories during range expansion. *Ecology Letters* 13:1210–1220.
- Chen, I.-C., J. K. Hill, R. Ohlemüller, D. B. Roy, and C. D. Thomas. 2011. Rapid range shifts of species associated with high levels of climate warming. *Science* 333:1024–1026.
- Davis, M. B., and R. G. Shaw. 2001. Range shifts and adaptive responses to quaternary climate change. *Science* 292:673–679.
- Excoffier, L., M. Foll, and R. J. Petit. 2009. Genetic consequences of range expansions. *Annual Review of Ecology, Evolution, and Systematics* 40:481–501.
- Gilman, S. E., M. C. Urban, J. Tewksbury, G. W. Gilchrist, and R. D. Holt. 2010. A framework for community interactions under climate change. *Trends in ecology & evolution* 25:325–331.
- Gonzalez, P., R. P. Neilson, J. M. Lenihan, and R. J. Drapek. 2010. Global patterns in the vulnerability of ecosystems to vegetation shifts due to climate change. *Global Ecology and Biogeography* 19:755–768.
- Hansen, A. J., R. P. Neilson, V. H. Dale, C. H. Flather, L. R. Iverson, D. J. Currie, S. Shafer, R. Cook, and P. J. Bartlein. 2001. Global change in forests: Responses of species, communities, and biomes: Interactions between climate change and land use are projected to cause large shifts in biodiversity. *AIBS Bulletin* 51:765–779.
- Hargreaves, A., S. Bailey, and R. A. Laird. 2015. Fitness declines towards range limits and local adaptation to climate affect dispersal evolution during climate-induced range shifts. *Journal of evolutionary biology* 28:1489–1501.
- Hargreaves, A. L., and C. G. Eckert. 2014. Evolution of dispersal and mating systems along geographic gradients: implications for shifting ranges. *Functional Ecology* 28:5–21.

265 Hastings, A., K. Cuddington, K. F. Davies, C. J. Dugaw, S. Elmendorf, A. Freestone, S. Harrison,  
266 M. Holland, J. Lambrinos, U. Malvadkar, et al. 2005. The spatial spread of invasions: new  
267 developments in theory and evidence. *Ecology Letters* 8:91–101.

268 Henry, R. C., G. Bocedi, and J. M. Travis. 2013. Eco-evolutionary dynamics of range shifts: elastic  
269 margins and critical thresholds. *Journal of Theoretical Biology* 321:1–7.

270 Hobbs, R. J., E. Higgs, and J. A. Harris. 2009. Novel ecosystems: implications for conservation  
271 and restoration. *Trends in ecology & evolution* 24:599–605.

272 Kubisch, A., R. D. Holt, H.-J. Poethke, and E. A. Fronhofer. 2014. Where am i and why? synthe-  
273 sizing range biology and the eco-evolutionary dynamics of dispersal. *Oikos* 123:5–22.

274 Lande, R. 1976. Natural selection and random genetic drift in phenotypic evolution. *Evolution*  
275 30:314–334.

276 Loarie, S. R., P. B. Duffy, H. Hamilton, G. P. Asner, C. B. Field, and D. D. Ackerly. 2009. The  
277 velocity of climate change. *Nature* 462:1052.

278 Melbourne, B. A., and A. Hastings. 2008. Extinction risk depends strongly on factors contributing  
279 to stochasticity. *Nature* 454:100.

280 Ochocki, B. M., and T. E. Miller. 2017. Rapid evolution of dispersal ability makes biological  
281 invasions faster and more variable. *Nature communications* 8:14315.

282 Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annu. Rev.*  
283 *Ecol. Evol. Syst.* 37:637–669.

284 Parmesan, C., N. Ryrholm, C. Stefanescu, J. K. Hill, C. D. Thomas, H. Descimon, B. Huntley,  
285 L. Kaila, J. Kullberg, T. Tammaru, et al. 1999. Poleward shifts in geographical ranges of butterfly  
286 species associated with regional warming. *Nature* 399:579.

287 Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across  
288 natural systems. *Nature* 421:37.



- Peñuelas, J., and M. Boada. 2003. A global change-induced biome shift in the montseny mountains (ne spain). *Global change biology* 9:131–140.
- Perry, A. L., P. J. Low, J. R. Ellis, and J. D. Reynolds. 2005. Climate change and distribution shifts in marine fishes. *science* 308:1912–1915.
- Phillips, B. L. 2015. Evolutionary processes make invasion speed difficult to predict. *Biological Invasions* 17:1949–1960.
- Ricker, W. E. 1954. Stock and recruitment. *Journal of the Fisheries Board of Canada* 11:559–623.
- Scholze, M., W. Knorr, N. W. Arnell, and I. C. Prentice. 2006. A climate-change risk analysis for world ecosystems. *Proceedings of the National Academy of Sciences* 103:13116–13120.
- Sekercioglu, C. H., S. H. Schneider, J. P. Fay, and S. R. Loarie. 2008. Climate change, elevational range shifts, and bird extinctions. *Conservation biology* 22:140–150.
- Shine, R., G. P. Brown, and B. L. Phillips. 2011. An evolutionary process that assembles phenotypes through space rather than through time. *Proceedings of the National Academy of Sciences* 108:5708–5711.
- Skellam, J. G. 1951. Random dispersal in theoretical populations. *Biometrika* 38:196–218.
- Szűcs, M., M. Vahsen, B. Melbourne, C. Hoover, C. Weiss-Lehman, and R. Hufbauer. 2017. Rapid adaptive evolution in novel environments acts as an architect of population range expansion. *Proceedings of the National Academy of Sciences* 114:13501–13506.
- Team, R. C. 2000. R language definition. Vienna, Austria: R foundation for statistical computing.
- Van Petegem, K. H., J. Boeye, R. Stoks, and D. Bonte. 2016. Spatial selection and local adaptation jointly shape life-history evolution during range expansion. *The American Naturalist* 188:485–498.

- 312 Walther, G.-R., E. Post, P. Convey, A. Menzel, C. Parmesan, T. J. Beebee, J.-M. Fromentin,  
313 O. Hoegh-Guldberg, and F. Bairlein. 2002. Ecological responses to recent climate change. *Nature*  
314 416:389.
- 315 Weiss-Lehman, C., R. A. Hufbauer, and B. A. Melbourne. 2017. Rapid trait evolution drives  
316 increased speed and variance in experimental range expansions. *Nature communications* 8:14303.
- 317 Williams, J. W., and S. T. Jackson. 2007. Novel climates, no-analog communities, and ecological  
318 surprises. *Frontiers in Ecology and the Environment* 5:475–482.
- 319 Zhu, K., C. W. Woodall, and J. S. Clark. 2012. Failure to migrate: lack of tree range expansion in  
320 response to climate change. *Global Change Biology* 18:1042–1052.