

Important declarations

Please remove this info from manuscript text if it is also present there.

Associated Data

Data supplied by the author:

1. Github: 2. 385737018 3. <https://git.io/JRmfe>

Required Statements

Competing Interest statement:

The authors declare that they have no competing interests.

Funding statement:

Research start-up funds grant No. 322388 were assigned to Miguel Uyaguari-Diaz at the University of Manitoba. Collaborative grant No. 52622 (Drs. Uyaguari and Yuan) was awarded by the Faculty of Science, University of Manitoba.

Quantitation of human enteric viruses as alternative indicators of fecal pollution to evaluate wastewater treatment processes

Audrey Garcia^{Equal first author, 1}, Tri Le^{Equal first author, 1}, Paul Jankowski¹, Kadir Yanaç², Qiuyan Yuan², Miguel I Uyaguari-Diaz^{Corresp. 1}

¹ Department of Microbiology, University of Manitoba, Winnipeg, Manitoba, Canada

² Department of Civil Engineering, University of Manitoba, Winnipeg, Manitoba, Canada

Corresponding Author: Miguel I Uyaguari-Diaz
Email address: Miguel.Uyaguari@umanitoba.ca

We investigated the potential use and quantitation of human enteric viruses in municipal wastewater samples of Winnipeg (Manitoba, Canada) as alternative indicators of contamination and evaluated the processing stages of the wastewater treatment plant. During the fall 2019 and winter 2020 seasons, samples of raw sewage, activated sludge, effluents, and biosolids (sludge cake) were collected from the North End Sewage Treatment Plant (NESTP), which is the largest wastewater treatment plant in the City of Winnipeg. DNA and RNA enteric viruses, as well as the *uidA* gene found in *Escherichia coli* were targeted in the samples collected from the NESTP. Total nucleic acids from each wastewater treatment sample were extracted using a commercial spin-column kit. Enteric viruses were quantitated in the extracted samples via quantitative PCR using TaqMan assays. The average gene copies assessed in the raw sewage were not significantly different (p-values ranged between 0.0547 and 0.7986) than the average gene copies assessed in the effluents for Adenovirus and crAssphage (DNA viruses), Pepper Mild Mottle Virus (RNA virus), and *uidA* in terms of both volume and biomass. A significant reduction of these enteric viruses was observed consistently in activated sludge samples compared with those for raw sewage. Corresponding reductions in gene copies per volume and gene copies per biomass were also seen for *uidA* but were not statistically significant (p-value = 0.8769 and p-value = 0.6353, respectively). The higher gene copy numbers of enteric viruses and *E. coli* observed in the effluents may be associated with the 12-hour hydraulic retention time in the facility. Enteric viruses found in gene copy numbers were at least one order of magnitude higher than the *E. coli* marker *uidA*. This indicate that enteric viruses may survive the wastewater treatment process and viral-like particles are being released into the aquatic environment. Our results suggest that Adenovirus, crAssphage, and Pepper mild mottle virus can be used as complementary viral indicators of human fecal

pollution.

Quantitation of human enteric viruses as alternative indicators of fecal pollution to evaluate wastewater treatment processes

Audrey Garcia^{1*}, Tri Le^{1*}, Paul Jankowski^{1#a}, Kadir Yanaç², Qiuyan Yuan² and Miguel Uyaguari-Díaz^{1^}

¹ Department of Microbiology, Faculty of Science, University of Manitoba, Winnipeg, MB, Canada R3T 2N2

² Department of Civil Engineering, Price Faculty of Engineering, University of Manitoba, Winnipeg, MB, Canada R3T 2N2

* These authors contributed equally to this work.

^{#a} Current address: Department of Medical Microbiology and Infectious Diseases, University of Manitoba, Winnipeg, Manitoba, Canada

[^] Corresponding Author:

Miguel Uyaguari-Díaz

45 Chancellors Circle, Winnipeg, Manitoba, R3T 2N2, Canada

Email address: Miguel.Uyaguari@umanitoba.ca

ABSTRACT

We investigated the potential use and quantitation of human enteric viruses in municipal wastewater samples of Winnipeg (Manitoba, Canada) as alternative indicators of contamination and evaluated the processing stages of the wastewater treatment plant. During the fall 2019 and winter 2020 seasons, samples of raw sewage, activated sludge, effluents, and biosolids (sludge cake) were collected from the North End Sewage Treatment Plant (NESTP), which is the largest wastewater treatment plant in the City of Winnipeg. DNA and RNA enteric viruses, as well as the *uidA* gene found in *Escherichia coli* were targeted in the samples collected from the NESTP. Total nucleic acids from each wastewater treatment sample were extracted using a commercial spin-column kit. Enteric viruses were quantitated in the extracted samples via quantitative PCR using TaqMan assays.

The average gene copies assessed in the raw sewage were not significantly different (p-values ranged between 0.0547 and 0.7986) than the average gene copies assessed in the effluents for Adenovirus and crAssphage (DNA viruses), Pepper Mild Mottle Virus (RNA virus), and *uidA* in terms of both volume and biomass. A significant reduction of these enteric viruses was observed consistently in activated sludge samples compared with those for raw sewage. Corresponding reductions in gene copies per volume and gene copies per biomass were also seen for *uidA* but were not statistically significant (p-value = 0.8769 and p-value = 0.6353, respectively). The higher gene copy numbers of enteric viruses and *E. coli* observed in the effluents may be associated with the 12-hour hydraulic retention time in the facility. Enteric viruses found in gene copy numbers were at least one order of magnitude higher than the *E. coli* marker *uidA*. This indicate that enteric viruses may survive the wastewater treatment process and viral-like particles are being released into the aquatic environment. Our results suggest that Adenovirus,

crAssphage, and Pepper mild mottle virus can be used as complementary viral indicators of human fecal pollution.

INTRODUCTION

The human fecal waste present in raw sewage (RS) contains pathogens that can cause numerous diseases. This can have a huge negative impact to public, aquatic health, and the economy (Stachler, et al., 2017). Wastewater treatment plants (WWTPs) serve as protective barriers between communities and the environment by reducing the organic matter present in wastewater. Water quality is currently assessed using traditional markers such as coliforms and *Escherichia coli*, leaving other microbes such as viruses largely unexplored. The North End Sewage Treatment Plant (NESTP) in Winnipeg, Manitoba handles approximately 70% of the city's wastewater treatment, serving over 400,000 people (City of Winnipeg, Water and Waste Department, 2020). The treatment process at the NESTP first involves RS undergoing primary treatment to remove solids. During the next treatment cycle, activated sludge (AS), a heterotrophic cocktail of bacteria and protozoa, degrades organic matter present in solid waste. The activated sludge (also known as biological treatment or secondary treatment) is the most widely used process around the world to treat municipal wastewater (Racz et al., 2010; Scholz, 2016), and its use will likely continue for centuries as it is a cheap and efficient treatment process. After the biological treatment, wastewater is UV-disinfected and discharged as effluents (EF) into the river (City of Winnipeg, Water and Waste Department, 2020). Approximately 200 million liters of EF are discharged per day (City of Winnipeg, Water and Waste Department, 2020).

The main indicator of biological contamination used in wastewater treatment screening is *E. coli*, a fecal coliform bacterium (Hood et al., 1983). It is present in the gut of humans and warm-blooded animals, and widely used as the main indicator of fecal pollution during the wastewater treatment process. Nevertheless, the use of only fecal bacteria indicator in wastewater excludes other possible pathogen groups present, such as human enteric viruses. Targeting these viruses in EF could be an alternative method to monitor the wastewater treatment process. Within this context, Dutilh et al. (2014) targeted the DNA crAssphage genome in a human fecal sample. With further bioinformatics testing, it was predicted that the crAssphage genome is highly abundant, and it was identified in 73% of human fecal metagenomes surveyed (Dutilh, et al., 2014). In a study conducted by Zhang et al. (2006), the most abundant fecal virus found in dry weight fecal matter was the plant RNA virus, Pepper mild mottle virus (PMMV). In the present study, samples of RS, AS, EF, and biosolids/sludge cake (SC) from the NESTP were collected (during fall 2019 and winter 2020) to investigate the potential of quantitating human enteric viruses in wastewater samples as complementary indicators of contamination to evaluate the processing stages of wastewater treatment. DNA enteric viruses in this study include human Adenovirus (AdV) and cross assembled phage (crAssphage), while RNA enteric viruses include PMMV, Noroviruses (NoV) of the genogroups GI and GII, Astrovirus (AstV), Sapovirus (SaV), and Rotavirus (RoV). We also studied the presence of a molecular marker for *E. coli*, the *uidA* gene, in the samples collected from the NESTP. An overview of the workflow is illustrated in *Fig. 1*.

MATERIALS AND METHODS

Sample Collection. A liter of RS, AS, EF, and 1 kg of SC were collected from the NESTP during each of the sampling events in fall 2019 and winter 2020. Each sample was sealed in a 1-L sterile polyethylene container lined with a sterile plastic bag. Samples were collected on October 22nd, 2019 (Event 1) and November 28th, 2019 (Event 2) in the fall season. In the winter season, samples were collected on December 18th, 2019 (Event 3) and February 6th, 2020 (Event 4). SC samples were collected earlier in the day during Events 3 and 4. All samples were kept at 4°C and processed within 24 hours of collection.

Assessment of Ultrafiltration for Viral Recovery Efficiency. Armored RNA (Asuragen, Inc., Austin, TX, USA), an artificial virus, was used to assess recovery efficiency of the ultrafiltration method employed herein. We spiked in 40,000 copies of Armored RNA into 120 mL raw sewage samples collected in duplicates from the NESTP, but this was not included as part of this study. Primers (381F: 5'- AGCCTGTCAATACCTGCACC-3' and 475R: 5'- CACGCTTAGATCTCCGTGCT-3'), and probe (420P: 5' Cy5- AGAGTATGAGAGGTCGACGA-TAO 3') were designed using Primer design tool of Geneious Prime version 2021.1.1 (<https://www.geneious.com>) and targeted a 95-bp region within the Armored RNA genome. This targeted 95-bp fragment was sent to Integrated DNA Technologies (IDT, Inc., Coralville, Iowa) to synthesize a gBlock construct. Serial dilutions of this synthetic fragment were used to generate standards and quantify gene copy numbers (GCNs) of Armored RNA via quantitative reverse transcription PCR (RT-qPCR). Standard and raw sewage samples were run in triplicates.

Thermal cycling reactions were performed at 50°C for 5 minutes, followed by 45 cycles at 95°C for 10 seconds and 60°C for 30 seconds on a QuantStudio 5 Real-Time PCR System (Life Technologies, Carlsbad, CA, USA). Each 10- μ l RT-qPCR mixture consisted of 2.5 μ L 4X TaqMan Fast Virus 1-Step Master Mix (Life Technologies, Carlsbad, CA, USA), 400 nM each primer, 200 nM probe, and 2.5 μ l of template and ultrapure DNase/RNase free distilled water (Promega Corporation, Fitchburg, WI, USA).

Ultrafiltration of Wastewater Samples. Each wastewater treatment sample (RS, AS, and EF), including Millipore Milli-Q water as a negative control, was first filtered via a funnel and cheesecloth to remove any solid waste or debris. Next, 140 mL of each wastewater sample was concentrated using an ultrafiltration method with Centricon Plus-70 filter units (Millipore Corporation, Billerica, MA, USA). The ultrafiltration process used a sterile glass pipette, where 70 mL of each wastewater sample was added into their correspondingly labeled sample filter cup pre-assembled with the filtrate collection cup. Each assembly was then sealed with a cap. The Centricon Plus-70 assemblies were placed into a swinging bucket rotor and centrifuged at 3000 x g for 30 minutes at 20°C. Subsequently, the filtrate was discarded, and the remaining 70 mL of the samples was added into their correspondingly labeled sample filter cup pre-assembled with the filtrate collection cup. Samples were spun at the same speed and temperature for 45 minutes. After centrifugation, the sample filter cup was separated from the filtrate collection cup. The concentration collection cup was then turned upside down and placed on top of the sample filter cup. The device was carefully inverted and placed into the centrifuge. Centricon Plus-70 filter units were centrifuged at 800 x g for 2 minutes at 20°C. After this step, the concentrated sample was collected from the concentration cup via a micropipette. The final volume was measured for

each wastewater sample. If needed, 10 mM Tris-HCl, pH 8.5 buffer (Qiagen Sciences, Maryland, MD) was added to the concentrate to make up a total volume of 250 μ L. If the final volume of the concentrate was over 250 μ L, Tris buffer was not added. Aliquots containing 250 μ L were made and stored at 4°C and processed within 24h.

Sludge Cake Preparation for Ultrafiltration. To remove cells from the SC samples, a 1X phosphate-buffered solution (PBS) with 0.15M NaCl, 0.05% Tween-20, and pH 7.5 was used. Approximately 30 g of SC sample per sampling event (Events 3 and 4) was collected and divided into six Falcon tubes for each event (~5-6 g per tube). Approximately 30 mL of PBS was added to each tube. The Falcon tubes filled with SC samples were homogenized at constant agitation for 15 minutes at 2500 rpm in a vortex mixer. These tubes were then centrifuged at a speed of 4500 x g for 50 minutes. The supernatant from each tube was subsequently recovered and transferred to a new sterile Falcon tube. For each sample event, 140 mL of supernatant was used for ultrafiltration as described previously.

Nucleic Acid (DNA/RNA) Extraction and Fluorometric Assessment. Once the final volume of concentrate was collected from each wastewater sample, the sample was pretreated with InhibitEX buffer (Qiagen Sciences, Maryland, MD) as indicated by the manufacturer. Then, QIAamp MinElute virus spin kit (Qiagen Sciences, Maryland, MD) was used to extract total nucleic acids from each wastewater sample as per the manufacturer's instructions, which included the use of Qiagen Protease and carrier RNA (Qiagen Sciences, Maryland, MD). Samples were eluted in 75 μ L of Buffer AVE (Qiagen Sciences, Maryland, MD), quantified, and stored at -80°C for downstream processes. The nucleic acid concentration and purity were

assessed using Qubit dsDNA high sensitivity and RNA assay kits in a Qubit 4 fluorometer (Invitrogen, Carlsbad, CA, USA), respectively. Qubit results can be found in Supplementary Materials (*Table S1*).

qPCR Primers, Probes, and gBlocks Gene Fragments. *Table 1* summarizes the primers and probes used in this study. Forward and reverse primers listed in *Table 1* were used in the Primer-BLAST tool to extract gene target regions (Ye, et al., 2012). Extracted regions were then uploaded to the Geneious software to verify oligonucleotide sequences associated to the flanking regions and probe. The generated sequences were sent to Integrated DNA Technologies (IDT, Inc., Coralville, Iowa, USA) to generate the desired gBlocks constructs. IDT manufactured all the primers used for qPCR, as well as the probes Ast-P, Ring1a.2, and Ring 2.2 (*Table 1*). However, probes Sav124TP, Sav5TP, NSP3-P, AdV-P, PMMV-P, and CrAss-P were manufactured by Life Technologies (Carlsbad, CA, USA).

Quantitative PCR Assays. Taqman Environmental Master Mix 2.0 (Life Technologies, Carlsbad, CA, USA) was used for assays involving DNA enteric viruses and *uidA*, while 4x Taqman Fast Virus 1-Step Master Mix (Life Technologies, Carlsbad, CA, USA) was used for RNA enteric viruses. Each 10 µl qPCR reaction contained 500 nM of each of the forward primer and the reverse primer and 250 nM of its designated probe when targeting both DNA and RNA viruses. Five µl of Environmental Master Mix was utilized in each qPCR reaction for targeting DNA viruses, while 2.5 µl of 4x Fast Virus Master Mix was used in each qPCR reaction for targeting RNA viruses. The *uidA* qPCR reaction consisted of 5 µl of Environmental Master Mix, 400 nM of each primer, and 100 nM of probe. All qPCR reactions used 2 µl of template.

Each qPCR reaction was performed in triplicates on an ABI QuantStudio 5 PCR system (Applied Biosystems, Foster City, CA, USA). The DNA enteric viruses (AdV and crAssphage) and *uidA* were subjected to the following conditions: 50.0°C for 2 minutes and 95.0°C for 10 minutes followed by 40 cycles of 95.0°C for 15 seconds and 60.0°C for 1 minute. The RNA enteric viruses (SaV, RoV, AstV, GI and GII NoV, PMMV) were subjected to the following conditions: 50.0°C for 5 minutes and 95.0°C for 20 seconds followed by 40 cycles of 95.0°C for 3 seconds and 60.0°C for 30 seconds. Raw qPCR output files can be found on GitHub (<https://git.io/J8VJ6>).

Assessment of Gene Copy Numbers by Volume and Biomass. Gene copy numbers (GCNs) were expressed in terms of sample (per mL or g of sample) and biomass (per ng of DNA or RNA). GCNs per mL of sample were calculated as previously described by Ritalahti et al. (2006). When calculating GCNs per mL of sample, the final volume recovered after filtering 140 mL of wastewater sample was used in the formula. For the SC samples, the mass of SC collected was used in the formula to produce results in GCNs per g of sample.

Collection of Metadata for Sampling Events. To perform Principal Component Analysis (PCA) and Spearman's rank correlation analysis for EF samples, metadata pertinent to the sampling events was retrieved. Water quality parameters obtained from the NESTP were combined with their October 2019 monitoring data (City of Winnipeg, Water and Waste Department, 2019) to complete some of the missing fields. For a value not found in either document, data interpolation was performed by taking an average of the corresponding values for the days before and after the sampling event. In addition, the Government of Canada's historical

weather database was utilized to obtain the mean temperature on the sampling dates and the total precipitation over three days before each sampling event (hereafter referred to as “precipitation”) (Environment and Climate Change Canada, 2021). The values for all parameters were transformed using \log_{10} , except for precipitation due to the presence of zero values. These variables were used with \log_{10} -transformed GCNs per mL sample for AdV, crAssphage, PMMV, and *uidA* (targets with quantifiable qPCR readings for all replicates across all events) as input for downstream analyses (PCA and Spearman’s rank correlation analysis).

Data Handling, Statistical Analysis, and Data Visualization. Various applications were employed to process data at different steps of the pipeline. Input data, such as output from the qPCR instrument, was subjected to rudimentary formatting and cleaning in Microsoft Excel, which was also used to calculate GCNs per mL or g sample and per ng nucleic acid. R (R Core Team, 2021) and its integrated development environment RStudio (RStudio Team, 2021) were utilized to further process the data and perform statistical analyses and output visualizations. These operations included general linear models (and estimated pairwise differences) using the package *sasLM* version 0.6.0 (Bae, 2021), PCA (corresponding biplots were created using the package *ggbiplot* version 0.55 (Vu, 2011)), and Spearman’s correlation matrix using the package *Hmisc* version 4.5-0 (Harrell Jr., 2021). The package *reshape2* version 1.4.4 (Wickham, 2020) was used to reformat these correlation matrices to enhance compatibility with other data-handling tools. Information about other packages is provided in Supplementary Materials (Table S2). The R script used for analysis can be found on GitHub (<https://git.io/J8VUI>).

Another software involved in data visualization was Tableau. Specifically, it was used to generate boxplots for GCNs per mL or g sample and per ng nucleic acid, as well as the heatmap representing the above-mentioned Spearman correlation matrix.

For all tests, a p-value of 0.05 was assumed to be the minimum level of significance.

RESULTS

From our assessment of the ultrafiltration method used in this study, the recovery efficiency of Armored RNA as measured by RT-qPCR was estimated to be between 7.14% and 8.64% for both raw sewage samples.

The GCN values for the DNA and RNA viruses and *uidA* were transformed into log₁₀ form. These values were run through a general linear model Tukey-Kramer analysis, and the means of each wastewater processing stage for each target were analyzed. The GCNs were expressed in terms of volume (mL) or weight (g) of sample and biomass (ng of nucleic acids). The presence of DNA and RNA viral gene copies and *uidA* in the Milli-Q water (negative control) samples across all Events 1-4 were negative. The orange-dotted lines in *Figs. 2-6* indicate the mean of the number of gene copies of each wastewater treatment sample across all events.

The average GCNs assessed in the RS were not significantly different (p-values ranged between 0.0547 and 0.7986) compared to the average GCNs assessed in the EF for the DNA enteric viruses (AdV and crAssphage) (*Fig. 2*), PMMV (*Fig. 3*), and *uidA* (*E. coli*) (*Fig. 4*) in terms of both volume and biomass. However, the average GCNs of the DNA enteric viruses assessed in AS were significantly and consistently lower compared to RS. Corresponding reductions in gene copies per volume and gene copies per biomass were also seen for *uidA*, although these reductions were not statistically significant, with p-values being 0.8769 and 0.6353, respectively.

244 For all the aforementioned targets, there was a relatively higher number of gene copies observed
 245 in the EF across all events compared to AS samples.

246 NoV GI and GII were also targets for our study. Boxplots of their GCNs across the different
 247 wastewater stages throughout Events 1-4 can be found in Supplementary Materials (*Fig. S1*). In
 248 Events 1 and 2 (Fall season), NoV GI was below qPCR detection limits for all samples (RS, AS,
 249 and EF). In addition, NoV GII GCNs for all samples collected in Event 2 and AS samples in
 250 Events 3 and 4 (Winter season) were also below the detection limits. Among the quantifiable
 251 samples, statistically significant GCN differences in terms of volume/mass and biomass were
 252 calculated for the pairs of AS-EF (p-values were 4.483×10^{-6} and 3.226×10^{-7} , respectively), AS-
 253 RS (1.658×10^{-6} , 1.091×10^{-5}), and AS-SC (1.481×10^{-9} , 4.083×10^{-7}). No significant
 254 differences were detected among treatments for GCNs of NoV GI. There was no significant
 255 difference between the mean gene copies of NoV GII in the RS and EF samples in terms of
 256 volume (p-value = 0.7377), but the difference was significant in terms of biomass (p-value =
 257 0.04905). The corresponding quantities of all the other sample pairs were statistically significant
 258 when looking at both the volume/mass and biomass perspectives, with p-values ranging from
 259 1.304×10^{-8} to 0.0046, except for AS-RS GCN difference in terms of biomass (p-value =
 260 0.0637).

261 RoV gene copies across the various wastewater treatment stages from Event 1 to 4 were also
 262 examined. The boxplots illustrating these results in terms of both sample and biomass can be
 263 found in the Supplementary Materials (*Fig. S2*). RoV GCNs were below detection limit for all
 264 samples collected in Events 1 and 2. Looking at the EF-SC pair, the mean GCNs differed
 265 significantly in terms of volume/mass (p-value = 2.649×10^{-7}) but not biomass (p-value =
 266 0.4298). No significant GCN differences could be detected between RS and AS samples in terms

of both volume (p-value = 0.4155) and biomass (p-value = 0.6662). The equivalent magnitudes for the remaining pairs per volume/mass and per biomass were statistically significant, with p-values being between 7.907×10^{-10} and 0.02433, respectively.

In the present study, there was no detection of gene copies for AstV and SaV (Sav1, Sav124, and Sav5) in any of the wastewater samples across all events. In addition, to eliminate the possibility of inhibitors or contaminants such as humic acids, additional qPCR tests using bovine serum albumin (data not shown) were conducted with environmental samples (including AS). No significant differences were observed between samples with and without the enzyme.

To investigate any potential relationship between collected data for EF samples, PCA was performed with \log_{10} -transformed variables. We found that three components (PC1, PC2, and PC3) explained 99.14% of the variance between variables. A summary of the weight of components is included in the Supplementary Materials (*Table S3*). PC1 and PC2 were used to create the biplot in *Fig. 5*. Biplots for PC1 versus PC3 (*Fig. S3*) and PC2 versus PC3 (*Fig. S4*) are included in the Supplementary Materials.

Overall, based on the biplot of PC1 and PC2, samples from the four events were distinct from one another, as point clusters of the four events can be seen occupying different quadrants. PC1, explaining 54.9% of the observed variance, received a notable and positive contribution from COD, cBOD, BOD, and TOC. Strongly negative contributors to PC1 were mean temperature, grab filtered UVT, $\text{NO}_x\text{-N}$, and TS. These observations were supported by subsequent Spearman's rank correlation analysis (*Fig. 6*), as COD, cBOD, BOD, and TOC demonstrated strongly positive correlations with one another (rho ranging between 0.8000 and 0.9487) (p-value < 0.005) and strongly negative correlations with mean temperature, grab filtered UVT, $\text{NO}_x\text{-N}$, and TS (rho ranging between -1.000 and -0.8000) (p-value < 0.005). PC2 explained

31.9% of the variance between sampling events and showed a strong contribution from crAssphage, *uidA*, and grab temperature. This observation was also supported by the Spearman's rank correlation analysis showing these variables having strongly positive correlation with one another (rho ranging between 0.7169 and 0.9218) (p-value < 0.0100). Additionally, in the biplot, the axes representing *E. coli* and fecal coliform specifically pointed towards the same quadrant, which was reflected in their moderately positive Spearman's coefficient (0.6325) (p-value = 0.0273). However, it is worth noting that *uidA* and *E. coli* exhibited a moderately weak negative correlation (rho = -0.3073), although it was not statistically significant (p-value = 0.3313). The two parameters with the strongest contribution against PC2 were grab pH and turbidity, which was illustrated by the strongly positive Spearman's coefficient heatmap (rho = 0.8000) (p-value = 0.0018).

DISCUSSION

The ultrafiltration method used in this study was assessed and the recovery efficiency was estimated to be between 7.14% and 8.64%. This range was lower compared to other methods to concentrate viral particles such as JumboSep (13.38% ± 9.11%) or skimmed milk flocculation (15.27% ± 3.32%), spiked-in wastewater samples, and using Armored RNA as internal control (Yanaç and Uyaguari, unpublished results). Viral particles may have been sorbed to biosolids present in wastewater samples. In this context, matrix has a significant effect for recovery of viral particles. When compared to other environmental matrices such as surface water samples, recovery efficiency is higher using ultrafiltration (tangential flow filtration) (32.6%±11.81%) and skimmed milk flocculation (42.64%± 15.12%) (Francis and Uyaguari, unpublished results). Samples with high concentration of particles or suspended solids tend to saturate filters and

impact the recovery of viral particles (Aslan et al., 2011; Karim et al., 2009; Uyaguari-Diaz, et al., 2016).

The GCNs were expressed in terms of biomass and volume (except for SC, which was expressed in g s of sample). The higher abundance and more stable signal over time of GCNs of AdV and crAssphage (Fig. 2) as well as PMMV (Fig. 3) relative to the results of other assays make these target more representative for conducting comparisons with *E. coli*. This persistent presence is consistent with various longitudinal studies previously performed (Ballesté, et al., 2019; Farkas, et al., 2018; Farkas, et al., 2019; Hamza et al., 2019; Nour, et al., 2021; Schmitz et al., 2016; Tandukar et al., 2020; Worley-Morse et al., 2019).

A reduction of AdV, crAssphage, PMMV, and *uidA* GCNs was observed consistently in AS samples (Figs. 2-4). This could be a result of viral particles being sorbed to larger fractions of organic matter that had been filtered by cheesecloth early in the sample-handling process or retained in the filtration devices as previously described. It is important to mention that samples were collected within a 2-hour period from RS→AS→EF consecutively within each sampling event. The higher GCNs of viruses and *E. coli* observed in the EF may be associated with the hydraulic retention time (12 hours) in the facility and may not reflect wastewater treatment profiles at the time of collection. Other variables to consider are the overflow of sewage from rainy events and fluctuations in mixed liquor-suspended solids (Pérez et al., 2019). In our study, there were 4.6 mm of precipitation for Event 1, which may have affected the results. In the PCA analysis (Fig. 5), the vector for precipitation sharply denotes data points representing Event 1, indicating a possible relationship. Precipitation was also found to have positive correlations with grab flow ($\rho = 0.7746$) and raw flow ($\rho = 0.7746$) (Fig. 6). Nonetheless, further studies are needed to corroborate the potential link between precipitation and microbial counts.

Moreover, the duration of anaerobic sludge digestion is 15 days (City of Winnipeg, Water and Waste Department, 2020). In this context, GCNs of *uidA* in the SC were significantly reduced by anaerobic digestion (Fig. 4). This may explain why the gene copies of *uidA* in terms of biomass were lower in SC compared to all treatments (p-value < 0.02). The average gene copies across all wastewater stages (RS, AS, and EF) for *uidA* were not significantly different in terms of both volume and biomass. When compared to *uidA*, enteric viruses were found to be at least one order of magnitude more abundant than the *E. coli* marker.

GCNs of crAssphage in terms of biomass in SC were significantly higher than RS (p-value = 0.0040) and AS (p-value = 5.877×10^{-5}) (Fig. 2). For PMMV, SC samples had significantly more GCNs in terms of biomass than samples from other parts of the wastewater treatment process (p-values ranged from 1.487×10^{-5} to 0.03788) (Fig. 3). Since SC is the by-product of RS and AS using anaerobic digestion, this may indicate that the presence of crAssphage and PMMV was lower in the wastewater being treated in the AS, but higher in the solids. On the other hand, GCNs of AdV in terms of biomass were not significantly different between the AS and SC samples (Fig. 2). Meanwhile, plant viruses such as PMMV remain more stable (in terms of biomass) during these digestion processes (Jumat, et al., 2017).

The higher presence of RoV gene copies in the EF (p-value = 0.0006592 in terms of sample and p-value = 0.001550 in terms of biomass) during the winter season (Fig. S2) may indicate a higher risk of transmission during cold seasons (Atabakhsh et al. 2020), since a greater presence of RoV in EF has been previously found during the winter season (Li, et al., 2011).

The negative results of SaV (Sav1, Sav124, and Sav5) across all wastewater treatment stages during the fall and winter season are consistent with Varela et al. (2018) where samples were

retrieved from a wastewater treatment plant in Tunisia. Their results did not support the general belief that the peak of detection of SaV occurs during the cold and rainy months of the year. However, quantitative detection of SaV in wastewater and river water in Japan showed an increased concentration of SaV in influents between winter and spring (December to May), but a decrease in SaV concentration during the summer and autumn months (July to October) (Haramoto et al., 2008). Yet another pattern of SaV presence was reported in France, as Sima et al. (2011) found the virus to be readily detected in influents but had no clear variations in numbers over the 9-month (October to June) duration of the study. Similarly, seasonal differences in SaV concentrations were not statistically significant in a 3-year study conducted by Song et al. (2021) in China between 2017 and 2019. As a result, there are other factors that can influence wastewater SaV concentrations. For example, it has been hypothesized that isoelectric point could affect how viruses and their different strains behave in bioreactors (Miura et al., 2015). The NESTP

The gene copies of NoV GI and GII were below the detection limit in many of the AS samples (in terms of both volume and biomass), but still relatively abundant in the EF (*Fig. S1*). A possible explanation for the greatly reduced viral GCNs in AS samples is the high efficiency with which NoV GI and GII are removed, a notion supported by literature (Ibrahim et al., 2020; Kitajima et al., 2014; Schmitz et al., 2016). Furthermore, considering the observation that these viruses were found in abundance in SC samples, another contributing factor could be limitations in the sample collection process, which might not have adequately retrieved the slurry part of the sludge where the viruses are found in greater numbers as they might have sorbed to the larger fractions of the sludge solids. The relative abundance of NoV GI and GII gene copies in RS and EF during the winter months (December and February) and the absence of NoV GII in RS in

October may be due to seasonal variability (Pérez, Guerrero, Orellana, Figuerola, & Erijman, 2019). However, the presence of NoV GI and GII gene copies in RS during Event 4 (February) is consistent with a study conducted by Flannery et al. (2012), in which the concentration of NoV GI and GII gene copies in the influents of a wastewater treatment plant were significantly higher during the winter months (January to March). This seasonal trend is also reflected colloquially through the virus's sobriquet, the winter vomiting bug (Farkas, et al., 2021).

In a study conducted by El-Senousy et al. (2007), high numbers of AstV gene copies (per liter) in sewage water samples (from the Greater Cairo area in Egypt) were observed at the end of autumn and during the winter months, but the AstV concentrations tended to decrease as temperatures increased. These results are different from our findings where there was no detection of AstV in any of the wastewater treatment stages across all events. These results may be due to seasonal variability (Pérez et al., 2019) and/or reflect the pattern of infection (Corpuz et al., 2020) within the community under study.

Additionally, PCA (*Fig. 5*) and Spearman's rank correlation analysis (*Fig. 6*) were conducted for EF samples to investigate potential connections between various physical, chemical, and biological parameters. PCA revealed that samples collected during different events from October to February were distinct from one another. This could indicate a seasonal variation in wastewater, at least in effluents. This outcome is consistent with previous literature (Comber et al., 2020). Organic chemical parameters such as COD, cBOD, BOD, and TOC were notable positive contributors to PC1, while mean temperature, grab filtered UVT, NO_x-N, and TS most negatively contributed to PC1. These observations were validated by subsequent Spearman's rank correlation analysis showing statistically significant coefficients. Grab filtered UVT being inversely correlated with COD, cBOD, BOD, and TOC is consistent with the widespread use of

UV radiation to regulate microbial growth in a variety of medium, including water (Raeiszadeh & Adeli, 2020). Furthermore, it had been suggested that UV is an important influence to the survival of pathogens in wastewater environments, especially in cold weather conditions, such as those found in Manitoba during the surveying period (Murphy, 2017). The NESTP uses UV disinfection. Further studies are needed to evaluate the survival of enteric viruses in these reservoirs by using modification to the biological treatment and/or the disinfection process. Some of these modifications include fixed bed reactors (Sizirici & Yildiz, 2020), biofilm systems such as membrane bioreactors, biofilters, biofiltration, and carriers (Zhao et al., 2019). Other disinfection processes include the use of chlorine (liquid sodium hypochlorite solution, solid calcium hypochlorite) or newer methods such as ozone (Mezzanotte et al., 2007; Abou-Elela et al., 2012; Collivignarelli et al., 2018).

There is a possibility that viral GCNs quantified in the EF may represent an overestimation of the actual number of infectious viral particles since qPCR detects both infective and non-infective agents and UV treatment influences viral viability (Lizasoain et al., 2017). Thus, the interpretation of these results must be performed with cautiousness. On the other hand, it is also possible that the non-enveloped enteric viruses (Corpuz, et al., 2020) studied here survived the wastewater treatment process. Non-enveloped viruses are more resilient than their enveloped counterparts in numerous environmental conditions and water treatment processes (La Rosa et al., 2020). This is because of the latter group's envelope, which contains receptors needed for infection; if the envelope is lysed, infection is not possible (La Rosa et al., 2020). Various publications have noted the resilience of non-enveloped viruses after wastewater treatment (Adefisoye et al., 2016; Campos & Lees, 2014; Farkas, et al., 2019; Fitzgerald, 2015; Fong et al., 2010; Li, et al., 2021; Prevost, et al., 2015; Ruggeri, et al., 2015; Varela, et al., 2018). In this

context, we have detected GCNs of AdV, crAssphage, and PMMV in environmental surface waters receiving discharges from the NESTP, two other WWTPs, and other areas radiating away from the WWTPs within the city of Winnipeg (Francis and Uyaguari, unpublished data). Therefore, despite potential factors affecting interpretation, our results still reflect the presence of several non-enveloped enteric viruses in EF samples with reasonable quantitative accuracy.

CONCLUSION

Our study's primary goal was to identify human enteric viruses with the potential to become alternative indicators of fecal pollution. Towards that end, we propose AdV, crAssphage, and PMMV as more stable viral indicators of water quality due to their quantifiability illustrated in this investigation and the literature. Regular monitoring of these organisms can be useful complements to current methods for assessing wastewater treatment processes. Such vigilance could be a helpful tool to assist public health efforts in the event of a viral outbreak. Additionally, our study indicated that enteric viruses may have survived the wastewater treatment process and viral-like particles are possibly being released into the aquatic environment. Therefore, in addition to such methods as UV radiation (which is currently used in the NESTP and was shown in our study to be inversely correlated with biological parameters), we also suggest that WWTPs consider implementing modifications and/or additions (disinfection processes) to their workflow to reduce the number of viral particles across different stages of the wastewater treatment process.

448 **ABBREVIATIONS**

449 **AdV:** Adenovirus

450 **AS:** activated sludge

451 **AstV:** Astrovirus

452 **BOD:** biochemical oxygen demand

453 **cBOD:** carbonaceous biochemical oxygen demand

454 **COD:** chemical oxygen demand

455 **EF:** effluents

456 **GCN:** gene copy number

457 **NESTP:** North End Sewage Treatment Plant

458 **NH₄-N:** ammonium-nitrogen

459 **NoV:** Norovirus

460 **NO_x-N:** nitrogen oxides - nitrogen

461 **PCA:** Principal Component Analysis

462 **PMMV:** Pepper mild mottle virus

463 **PO₄-P:** orthophosphate as phosphorus

464 **RoV:** Rotavirus

465 **RS:** raw sewage

466 **RT-qPCR:** quantitative reverse transcription PCR

467 **SaV:** Sapovirus

468 **SC:** sludge cake

469 **sCOD:** soluble chemical oxygen demand

470 **TN:** total nitrogen

471 **TOC:** total organic carbon

472 **TP:** total phosphorus

473 **TS:** total solids

474 **TSS:** total suspended solids

475 ***uidA*:** β -d-glucuronidase gene

476 **WWTP:** wastewater treatment plant

477 **AUTHOR CONTRIBUTIONS**

478 AG performed the experiments, analyzed the data, prepared the figures and tables, and wrote and
479 reviewed the drafts of the manuscript.

480 TL analyzed the data, prepared the figures and tables, and wrote and reviewed the drafts of the
481 manuscript.

482 PJ performed the experiments and reviewed the drafts of the manuscript.

483 KY performed the validation experiments here described and reviewed the drafts of the
484 manuscript.

485 QY contributed the analysis tools and reviewed the drafts of the manuscript.

486 MUD designed the experiments, provided lead guidance during the experiments and analyses,
487 contributed the analysis tools, and reviewed the drafts of the manuscript.

488 All authors read and approved of the final manuscript.

489 **ACKNOWLEDGEMENTS**

490 Special thanks to the City of Winnipeg and Palwinder Singh, graduate student, Department of
491 Civil Engineering at the University of Manitoba (UoM) for sample collection. Research start-up
492 funds grant No. 322388 were assigned to Miguel Uyaguari-Diaz at the UoM. We acknowledge
493 The Faculty of Science, UoM, collaborative grant No. 52622 (Drs. Uyaguari and Yuan).

494 This research was conducted at the University of Manitoba. "*The University of Manitoba*
 495 *campuses are located on original lands of Anishinaabeg, Cree, Oji-Cree, Dakota, and Dene*
 496 *peoples, and on the homeland of the Métis Nation*".

497

REFERENCES

- Abou-Elela, S. I., El-Sayed, M. M. H., El-Gendy, A. S., & Abou-Taleb, E. M. (October 2012). Comparative study of disinfection of secondary treated wastewater using chlorine, UV and ozone. *Journal of Applied Sciences Research*, pp.5190-5197 ref.12
- Adefisoye, M. A., Nwodo, U. U., Green, E., & Okoh, A. I. (2016). Quantitative PCR Detection and Characterisation of Human Adenovirus, Rotavirus and Hepatitis A Virus in Discharged Effluents of Two Wastewater Treatment Facilities in the Eastern Cape, South Africa. *Food and Environmental Virology*, 8, 262-274. doi:10.1007/s12560-016-9246-4
- Aslan, A., Xagorarakis, I., Simmons, F., Rose, J., & Dorevitch, S. (2011, August 19). Occurrence of adenovirus and other enteric viruses in limited-contact freshwater recreational areas and bathing waters. *Journal of Applied Microbiology*, 111(5), 1250-1261. doi:10.1111/j.1365-2672.2011.05130.x
- Bae, K.-S. (2021). sasLM: 'SAS' Linear Model. Retrieved from <https://CRAN.R-project.org/package=sasLM>
- Ballesté, E., Pascual-Benito, M., Martín-Díaz, J., Blanch, A. R., Lucena, F., Muniesa, M., . . . García-Aljaro, C. (2019, May 15). Dynamics of crAssphage as a human source tracking marker in potentially faecally polluted environments. *Water Research*, 155, 233-244. doi:10.1016/j.watres.2019.02.042
- Campos, C. J., & Lees, D. N. (2014, June). Environmental Transmission of Human Noroviruses in Shellfish Waters. *Applied and Environmental Microbiology*, 80(12), 3552-3561. doi:10.1128/AEM.04188-13
- City of Winnipeg, Water and Waste Department. (2019, October). North End Water Pollution Control Centre Monitoring Data. Winnipeg, Manitoba, Canada. Retrieved July 21, 2021,

from

[https://www.winnipeg.ca/waterandwaste/pdfs/sewage/ComplianceReporting/2019/oct/ne
wpcc.pdf](https://www.winnipeg.ca/waterandwaste/pdfs/sewage/ComplianceReporting/2019/oct/ne
wpcc.pdf)

City of Winnipeg, Water and Waste Department. (2020, October 8). *Sewage Treatment Plants*.

Retrieved July 21, 2021, from City of Winnipeg:

[https://www.winnipeg.ca/waterandwaste/sewage/treatmentPlant/default.stm#tab-north-
end-sewage-treatment-plant](https://www.winnipeg.ca/waterandwaste/sewage/treatmentPlant/default.stm#tab-north-
end-sewage-treatment-plant).

Collivignarelli, M.C., Abbà, A., Benigna, I., Sorlini, S., & Torretta, V. (2018). Overview of the
Main Disinfection Processes for Wastewater and Drinking Water Treatment
Plants. *Sustainability*, 10, 86. <https://doi.org/10.3390/su10010086>.

Comber, S. D., Gardner, M. J., & Ellor, B. (2020, September). Seasonal variation of contaminant
concentrations in wastewater treatment works effluents and river waters. *Environmental
Technology*, 41(21), 2716-2730. doi:10.1080/09593330.2019.1579872

Corpuz, M. V., Buonerba, A., Vigliotta, G., Zarra, T., Ballesteros Jr, F., Campiglia, P., . . .
Naddeo, V. (2020, November 25). Viruses in wastewater: occurrence, abundance and
detection methods. *Science of the Total Environment*, 745.
doi:10.1016/j.scitotenv.2020.140910

Dutilh, B. E., Cassman, N., McNair, K., Sanchez, S. E., Silva, G. G., Boling, L., . . . Edwards, R.
A. (2014). A highly abundant bacteriophage discovered in the unknown sequences of
human faecal metagenomes. *Nature Communications*, 5(4498), 1-11.
doi:10.1038/ncomms5498

El-Senousy, W. M., Guix, S., Abid, I., Pintó, R. M., & Bosch, A. (2007, January). Removal of
astrovirus from water and sewage treatment plants, evaluated by a competitive reverse

transcription-PCR. *Applied and Environmental Microbiology*, 73(1), 164-7.
doi:10.1128/AEM.01748-06

Environment and Climate Change Canada. (2021). *Historical Data*. Retrieved July 21, 2021, from Government of Canada:
https://climate.weather.gc.ca/historical_data/search_historic_data_e.html

Farkas, K., Adriaenssens, E. M., Walker, D. I., McDonald, J. E., Malham, S. K., & Jones, D. L. (2019, June). Critical Evaluation of CrAssphage as a Molecular Marker for Human-Derived Wastewater Contamination in the Aquatic Environment. *Food and Environmental Virology*, 11(2), 113-119. doi:10.1007/s12560-019-09369-1

Farkas, K., Green, E., Rigby, D., Cross, P., Tyrrel, S., Malham, S. K., & Jones, D. L. (2021, May 27). Investigating awareness, fear and control associated with norovirus and other pathogens and pollutants using best–worst scaling. *Scientific Reports*, 11.

Farkas, K., Marshall, M., Cooper, D., McDonald, J. E., Malham, S. K., Peters, D. E., . . . Jones, D. L. (2018, November). Seasonal and diurnal surveillance of treated and untreated wastewater for human enteric viruses. *Environmental Science and Pollution Research*, 25(33), 33391-33401. doi:10.1007/s11356-018-3261-y

Fitzgerald, A. (2015). *Review of Approaches for Establishing Exclusion Zones for Shellfish Harvesting around Sewage Discharge Points - Desk Study to Inform Consideration of the Possible Introduction of Exclusion Zones as a Control for Norovirus in Oysters*. Technical Report, Aquatic Water Services Ltd. Retrieved July 26, 2021, from <https://webarchive.nationalarchives.gov.uk/20150418173120/http://www.food.gov.uk/sites/default/files/Exclusion%20Zones%20Project%20FS513404%20-%20Technical%20Report%20FINAL.pdf>

- 567 Fong, T.-T., Phanikumar, M. S., Xagorarakis, I., & Rose, J. B. (2010, February). Quantitative
568 detection of human adenoviruses in wastewater and combined sewer overflows
569 influencing a Michigan river. *Applied and Environmental Microbiology*, 76(3), 715-23.
570 doi:10.1128/AEM.01316-09
- 571 Frahm, E., & Obst, U. (2003). Application of the fluorogenic probe technique (TaqMan PCR) to
572 the detection of *Enterococcus* spp. and *Escherichia coli* in water samples. *Journal of*
573 *Microbiological Methods*, 52(1), 123-31. doi:10.1016/s0167-7012(02)00150-1
- 574 Genz, A., Bretz, F., Miwa, T., Mi, X., Leisch, F., Scheipl, F., & Hothorn, T. (2021). mvtnorm:
575 Multivariate Normal and t Distributions. Retrieved from [http://CRAN.R-](http://CRAN.R-project.org/package=mvtnorm)
576 [project.org/package=mvtnorm](http://CRAN.R-project.org/package=mvtnorm)
- 577 Hamza, H., Rizk, N. M., Gad, M. A., & Hamza, I. A. (2019, November). Pepper mild mottle
578 virus in wastewater in Egypt: a potential indicator of wastewater pollution and the
579 efficiency of the treatment process. *Archives of Virology*, 164(11), 2707-2713.
580 doi:10.1007/s00705-019-04383-x
- 581 Haramoto, E., Katayama, H., Phanuwat, C., & Ohgaki, S. (2008, March). Quantitative detection
582 of sapoviruses in wastewater and river water in Japan. *Letters in Applied Microbiology*,
583 46(3), 408-13. doi:10.1111/j.1472-765X.2008.02330.x
- 584 Harrell Jr., F. E. (2021). Hmisc: Harrell Miscellaneous. Retrieved from [https://CRAN.R-](https://CRAN.R-project.org/package=Hmisc)
585 [project.org/package=Hmisc](https://CRAN.R-project.org/package=Hmisc)
- 586 Hood, M. A., Ness, G. E., & Blake, N. J. (1983). Relationship among fecal coliforms,
587 *Escherichia coli*, and *Salmonella* spp. in shellfish. *Applied and environmental*
588 *microbiology*, 45(1), 122-126. doi:10.1128/aem.45.1.122-126.1983

589 Ibrahim, C., Hammami, S., Khelifi, N., Pothier, P., & Hassen, A. (2020). The Effectiveness of
590 Activated Sludge Procedure and UV-C 254 in Norovirus Inactivation in a Tunisian
591 Industrial Wastewater Treatment Plant. *Food and Environmental Virology*, 12, 250-259.
592 doi:10.1007/s12560-020-09434-0

593 Jumat, M. R., Hasan, N. A., Subramanian, P., Heberling, C., Colwell, R. R., & Hong, P.-Y.
594 (2017). Membrane Bioreactor-Based Wastewater Treatment Plant in Saudi Arabia:
595 Reduction of Viral Diversity, Load, and Infectious Capacity. *Water*, 9(7).
596 doi:10.3390/w9070534

597 Kageyama, T., Kojima, S., Shinohara, M., Uchida, K., Fukushi, S., Hoshino, F. B., . . .
598 Katayama, K. (2003, April). Broadly Reactive and Highly Sensitive Assay for Norwalk-
599 Like Viruses Based on Real-Time Quantitative Reverse Transcription-PCR. *Journal of*
600 *Clinical Microbiology*, 41(4), 1548-1557. doi:10.1128/JCM.41.4.1548-1557.2003

601 Karim, M. R., Rhodes, E. R., Brinkman, N., Wymer, L., & Fout, G. S. (2009, April). New
602 Electropositive Filter for Concentrating Enteroviruses and Noroviruses from Large
603 Volumes of Water. *Applied and Environmental Microbiology*, 75(8), 2393-2399.
604 doi:10.1128/AEM.00922-08

605 Kitajima, M., Iker, B. C., Pepper, I. L., & Gerba, C. P. (2014, August 1). Relative abundance and
606 treatment reduction of viruses during wastewater treatment processes--identification of
607 potential viral indicators. *Science of the Total Environment*, 488-489, 290-296.
608 doi:10.1016/j.scitotenv.2014.04.087

609 La Rosa, G., Bonadonna, L., Lucentini, L., Kenmoe, S., & Suffredini, E. (2020, July 15).
610 Coronavirus in water environments: Occurrence, persistence and concentration methods -
611 A scoping review. *Water Research*, 179. doi:10.1016/j.watres.2020.115899

- 612 Li, D., Gu, A. Z., Zeng, S.-Y., Yang, W., He, M., & Shi, H.-C. (2011, May). Monitoring and
613 evaluation of infectious rotaviruses in various wastewater effluents and receiving waters
614 revealed correlation and seasonal pattern of occurrences. *Journal of Applied*
615 *Microbiology*, 110(5), 1129-37. doi:10.1111/j.1365-2672.2011.04954.x
- 616 Li, X., Cheng, Z., Dang, C., Zhang, M., Zheng, Y., & Xia, Y. (2021, July). Metagenomic and
617 viromic data mining reveals viral threats in biologically treated domestic wastewater.
618 *Environmental Science and Ecotechnology*, 7. doi:10.1016/j.ese.2021.100105
- 619 Lizasoain, A., Tort, L., García, M., Gillman, L., Alberti, A., Leite, J., . . . Colina, R. (2017).
620 Human enteric viruses in a wastewater treatment plant: evaluation of activated sludge
621 combined with UV disinfection process reveals different removal performances for
622 viruses with different features. *Letters in Applied Microbiology*, 66(3), 215-221. doi:DOI:
623 10.1111/lam.12839
- 624 Makowski, D., Ben-Shachar, M., Patil, I., & Lüdecke, D. (2020). Automated Results Reporting
625 as a Practical Tool to Improve Reproducibility and Methodological Best Practices
626 Adoption. Retrieved from <https://github.com/easystats/report>.
- 627 Mezzanotte, V., Antonelli, M., Citterio, S., & Nurizzo, C. (2007). Wastewater disinfection
628 alternatives: chlorine, ozone, peracetic acid, and UV light. *Water Environment Research*,
629 79(12), 2373-2379. doi:10.2175/106143007x183763.
- 630 Miura, T., Okabe, S., Nakahara, Y., & Sano, D. (2015, May 15). Removal properties of human
631 enteric viruses in a pilot-scale membrane bioreactor (MBR) process. *Water Research*, 75,
632 282-291. doi:10.1016/j.watres.2015.02.046
- 633 Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control.
634 (2017). Detecting Norovirus by Fast Real-Time RT-PCR. British Columbia, Canada.

635 Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control.
 636 (2017). Performing the GI Virus Panel by Real-Time PCR Procedure. British Columbia,
 637 Canada.

638 Murphy, H. (2017). Persistence of Pathogens in Sewage and Other Water Types. In J. Rose, & B.
 639 Jiménez-Cisneros (Eds.), *Global Water Pathogen Project* (Vol. 4). E. Lansing, MI,
 640 UNESCO. doi:10.14321/waterpathogens.51

641 Nour, I., Hanif, A., Zakri, A. M., Al-Ashkar, I., Alhetheel, A., & Eifan, S. (2021, April 29).
 642 Human Adenovirus Molecular Characterization in Various Water Environments and
 643 Seasonal Impacts in Riyadh, Saudi Arabia. *International Journal of Environmental*
 644 *Research and Public Health*, 18(9), 4773. doi:10.3390/ijerph18094773

645 Oka, T., Katayama, K., Hansman, G. S., Kageyama, T., Ogawa, S., Wu, F.-T., . . . Takeda, N.
 646 (2006). Detection of human sapovirus by real-time reverse transcription-polymerase
 647 chain reaction. *Journal of Medical Virology*, 78(10), 1347-1353. doi:10.1002/jmv.20699

648 Pérez, M. V., Guerrero, L. D., Orellana, E., Figuerola, E. L., & Erijman, L. (2019, July 2). Time
 649 Series Genome-Centric Analysis Unveils Bacterial Response to Operational Disturbance
 650 in Activated Sludge. *mSystems*, 4(4). doi:10.1128/mSystems.00169-19

651 Prevost, B., Lucas, F. S., Ambert-Balay, K., Pothier, P., Moulin, L., & Wurtzer, S. (2015,
 652 October). Deciphering the Diversities of Astroviruses and Noroviruses in Wastewater
 653 Treatment Plant Effluents by a High-Throughput Sequencing Method. *Applied and*
 654 *Environmental Microbiology*, 81(20), 7215-7222. doi:10.1128/AEM.02076-15.

655 R Core Team. (2021). R: A language and environment for statistical computing. R Foundation
 656 for Statistical Computing, Vienna, Austria. Retrieved from <https://www.R-project.org/>

657 Racz, L., T. Datta, and R. Goel. 2010. Effect of organic carbon on ammonia oxidizing bacteria in
658 a mixed culture. *Bioresource Technology*, 101 (16), 6454-60.

659 Raeiszadeh, M., & Adeli, B. (2020, October 14). A Critical Review on Ultraviolet Disinfection
660 Systems against COVID-19 Outbreak: Applicability, Validation, and Safety
661 Considerations. *ACS Photonics*. doi:10.1021/acsp Photonics.0c01245

662 Ritalahti, K. M., Amos, B. K., Sung, Y., Wu, Q., Koenigsberg, S. S., & Löffler, F. E. (2006).
663 Quantitative PCR targeting 16S rRNA and reductive dehalogenase genes simultaneously
664 monitors multiple Dehalococcoides strains. *Applied and Environmental Microbiology*,
665 72(4), 2765-74. doi:10.1128/AEM.72.4.2765-2774.2006

666 Rosario, K., Symonds, E. M., Sinigalliano, C., Stewart, J., & Breitbart, M. (2009). Pepper Mild
667 Mottle Virus as an Indicator of Fecal Pollution. *Applied and Environmental*
668 *Microbiology*, 75(22), 7261-7267. doi:10.1128/AEM.00410-09

669 Rosman, N. H., Anuar, A. N., Chelliapan, S., Din, M. F., & Ujang, Z. (2014, June).
670 Characteristics and performance of aerobic granular sludge treating rubber wastewater at
671 different hydraulic retention time. *Bioresource Technology*, 161, 155-61.
672 doi:10.1016/j.biortech.2014.03.047

673 RStudio Team. (2021). RStudio: Integrated Development Environment for R. Boston, MA:
674 RStudio, PBC. Retrieved from <http://www.rstudio.com/>

675 Ruggeri, F. M., Bonomo, P., Ianiro, G., Battistone, A., Delogu, R., Germinario, C., . . . Fiore, L.
676 (2015, January). Rotavirus Genotypes in Sewage Treatment Plants and in Children
677 Hospitalized with Acute Diarrhea in Italy in 2010 and 2011. *Applied and Environmental*
678 *Microbiology*, 81(1), 241-249. doi:10.1128/AEM.02695-14

679 Sarkar, D. (2008). *Lattice: Multivariate Data Visualization with R*. New York: Springer.

Scholz, M. Chapter 15-Activated Sludge Processes, Editor(s): Miklas Scholz, Wetlands for Water Pollution Control (Second Edition), Elsevier, Pages 91-105.

Schmitz, B. W., Kitajima, M., Campillo, M. E., Gerba, C. P., & Pepper, I. L. (2016). Virus Reduction during Advanced Bardenpho and Conventional Wastewater Treatment Processes. *Environmental Science & Technology*, 50(17), 9524-9532. doi:10.1021/acs.est.6b01384.

Sima, L. C., Schaeffer, J., Saux, J.-C. L., Parnaudeau, S., Elimelech, M., & Guyader, F. S. (2011, August). Calicivirus Removal in a Membrane Bioreactor Wastewater Treatment Plant. *Applied and Environmental Microbiology*, 77(15), 5170-5177. doi:10.1128/AEM.00583-11.

Sizirici, B., & Yildiz, I. (2020). Organic matter removal via activated sludge immobilized gravel in fixed bed reactor. *E3S Web of Conferences*, 191, 03006, 1-5.

Song, K., Lin, X., Liu, Y., Ji, F., Zhang, L., Chen, P., . . . Xu, A. (2021). Detection of Human Sapoviruses in Sewage in China by Next Generation Sequencing. *Food and Environmental Virology*, 13, 270-280. doi:10.1007/s12560-021-09469-x

Stachler, E., Akyon, B., Carvalho, N. A., Ference, C., & Bibby, K. (2018). Correlation of crAssphage qPCR Markers with Culturable and Molecular Indicators of Human Fecal Pollution in an Impacted Urban Watershed. *Environmental Science & Technology*, 52(13), 7505-7512. doi:10.1021/acs.est.8b00638

Stachler, E., Kelty, C., Sivaganesan, M., Li, X., Bibby, K., & Shanks, O. C. (2017). Quantitative CrAssphage PCR Assays for Human Fecal Pollution Measurement. *Environmental Science & Technology*, 51(16), 9146-9154. doi:10.1021/acs.est.7b02703

702 Tandukar, S., Sherchan, S. P., & Haramoto, E. (2020). Applicability of crAssphage, pepper mild
 703 mottle virus, and tobacco mosaic virus as indicators of reduction of enteric viruses during
 704 wastewater treatment. *Scientific Reports*, 10(3616). doi:10.1038/s41598-020-60547-9

705 Therneau, T. (2021). A Package for Survival Analysis in R. Retrieved from [https://CRAN.R-](https://CRAN.R-project.org/package=survival)
 706 [project.org/package=survival](https://CRAN.R-project.org/package=survival)

707 Ushey, K., Allaire, J., Wickham, H., & Ritchie, G. (2020). rstudioapi: Safely Access the RStudio
 708 API. Retrieved from <https://CRAN.R-project.org/package=rstudioapi>

709 Uyaguari-Diaz, M. I., Chan, M., Chaban, B. L., Croxen, M. A., Finke, J. F., Hill, J. E., . . . Tang,
 710 P. (2016). A comprehensive method for amplicon-based and metagenomic
 711 characterization of viruses, bacteria, and eukaryotes in freshwater samples. *Microbiome*,
 712 4(20). doi:10.1186/s40168-016-0166-1

713 Varela, M. F., Ouadani, I., Kato, T., Kadoya, S., Aouni, M., Sano, D., & Romalde, J. L. (2018,
 714 March 1). Sapovirus in Wastewater Treatment Plants in Tunisia: Prevalence, Removal,
 715 and Genetic Characterization. *Applied and Environmental Microbiology*, 84(6).
 716 doi:10.1128/AEM.02093-17

717 Vu, V. Q. (2011). ggbiplot: A ggplot2 based biplot. Retrieved from
 718 <http://github.com/vqv/ggbiplot>

719 Wang, X. M., Wei, Z. M., Guo, J. M., Cai, J. M., Chang, H. M., Ge, Y. M., & Zeng, M. M.
 720 (2019, November). Norovirus Activity and Genotypes in Sporadic Acute Diarrhea in
 721 Children in Shanghai During 2014–2018. *The Pediatric Infectious Disease Journal*,
 722 38(11), 1085-1089. doi:10.1097/INF.0000000000002456

723 Wickham, H. (2011). The Split-Apply-Combine Strategy for Data Analysis. *Journal of*
 724 *Statistical Software*, 40(1), 1-29. doi:10.18637/jss.v040.i01

725 Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag.

726 Wickham, H. (2020, April 9). reshape2: Flexibly Reshape Data: A Reboot of the Reshape
727 Package. Retrieved from <https://cran.r-project.org/package=reshape2>

728 Wickham, H., & Bryan, J. (2019). readxl: Read Excel Files. Retrieved from [https://CRAN.R-](https://CRAN.R-project.org/package=readxl)
729 [project.org/package=readxl](https://CRAN.R-project.org/package=readxl)

730 Wickham, H., & Bryan, J. (2021). usethis: Automate Package and Project Setup. Retrieved from
731 <https://CRAN.R-project.org/package=usethis>

732 Wickham, H., & Seidel, D. (2020). scales: Scale Functions for Visualization. Retrieved from
733 <https://CRAN.R-project.org/package=scales>

734 Wickham, H., François, R., Henry, L., & Müller, K. (2021). dplyr: A Grammar of Data
735 Manipulation. Retrieved from <https://CRAN.R-project.org/package=dplyr>

736 Wickham, H., Hester, J., & Chang, W. (2021). devtools: Tools to Make Developing R Packages
737 Easier. Retrieved from <https://CRAN.R-project.org/package=devtools>

738 Worley-Morse, T., Mann, M., Khunjar, W., Olabode, L., & Gonzalez, R. (2019, September).
739 Evaluating the fate of bacterial indicators, viral indicators, and viruses in water resource
740 recovery facilities. *Water Environment Research*, 91(9), 830-842. doi:10.1002/wer.1096

741 Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., & Madden, T. L. (2012, June 18).
742 Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction.
743 *BMC Bioinformatics*, 13(134). doi:10.1186/1471-2105-13-134

744 Zeileis, A., & Croissant, Y. (2010). Extended Model Formulas in R: Multiple Parts and Multiple
745 Responses. *Journal of Statistical Software*, 34(1), 1-13. doi:10.18637/jss.v034.i01

746 Zeng, S. Q., Halkosalo, A., Salminen, M., Szakal, E. D., Puustinen, L., & Vesikari, T. (2008).
 747 One-step quantitative RT-PCR for the detection of rotavirus in acute gastroenteritis.
 748 *Journal of Virological Methods*, 153(2), 238-40. doi:10.1016/j.jviromet.2008.08.004
 749 Zhang, T., Breitbart, M., Lee, W. H., Run, J.-Q., Wei, C. L., Soh, S. W., Hibberd, M. L., Liu, E.
 750 T., Rohwer, F., & Ruan, Y. (2006, January). RNA Viral Community in Human Feces:
 751 Prevalence of Plant Pathogenic Viruses. *PLOS Biology*, 4(1).
 752 doi:10.1371/journal.pbio.0040003
 753 Zhao, Y., Liu, D., Huang, W., Yang, Y., Ji, M., Nghiem, L. D., Trinh, Q. T., Tran, N.
 754 H. (2018). Insights into biofilm carriers for biological wastewater treatment
 755 processes: Current state-of-the-art, challenges, and opportunities. *Bioresource*
 756 *Technology*, 288, 121619. doi.org/10.1016/j.biortech.2019.121619.
 757

758 **Table 1. Primers and probes used in the present study.**
 759 **Table S1. Qubit results of extracted nucleic acid samples.**
 760 **Table S2. R packages used that were not mentioned in the manuscript.**
 761 **Table S3. Summary of weight of components of PCA for EF samples and related metadata.**
 762

Figure 1. Graphical abstract of workflow.

Figure 2. Box plots of the number of gene copies of DNA enteric viruses across each wastewater stage throughout Events 1-4.

The unit for the SC in Figures 2A and 2C is gene copies per g of sample.

Figure 3. Box plots of the number of genes copies of PMMV across each wastewater stage throughout Events 1-4.

The unit for the SC in Figure 3A is gene copies per g of sample.

Figure 4. Box plots of the number of gene copies of *uidA* across each wastewater stage throughout Events 1-4.

The unit for the SC in Figure 4A is gene copies per g of sample.

Figure 5. Principal Component Analysis of \log_{10} -transformed EF parameters, PC1 versus PC2.

The only variable not \log_{10} -transformed was precipitation due to presence of zero values.

Figure 6. Heatmap showing Spearman's rank correlation analysis between parameters collected for EF sampling events.

Figure S1. Box plots of the number of gene copies of Noroviruses GI and GII across each wastewater stage throughout Events 1-4.

786 The unit for the SC in Figures S1A and S1C is gene copies per g of sample.

787

788 **Figure S2. Box plots of the number of gene copies of Rotavirus across each wastewater**
789 **stage throughout Events 1-4.**

790 The unit for the SC in Figure S2A is gene copies per g of sample.

791

792 **Figure S3. Principal Component Analysis of \log_{10} -transformed EF parameters, PC1 versus**
793 **PC3.**

794 The only variable not \log_{10} -transformed was precipitation due to presence of zero values.

795

796 **Figure S4. Principal Component Analysis of \log_{10} -transformed EF parameters, PC2 versus**
797 **PC3.**

798 The only variable not \log_{10} -transformed was precipitation due to presence of zero values.

799

Table 1 (on next page)

Table 1. Primers and probes used in the present study.

1

Target	DN A or RN A	Primer/ Probe	Sequence (5'-3')	Genomic Target	References
Adenovirus 40/41	DN A	AdV-F	GCC TGG GGA ACA AGT TCA G	Hexon	Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control, 2017a
		AdV-R	ACG GCC AGC GTA AAG CG		
		AdV-P (Probe)	NED-ACC CAC GAT GTA ACC AC- MGB-NFQ		
Astrovirus	RN A	Ast-F	AAG CAG CTT CGT GAR TCT GG	Junction of polymerase and capsid	Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control, 2017a
		Ast-R	GCC ATC RCA CTT CTT TGG TCC		
		Ast-P (Probe)	Cy5-CAC AGA AGA GCA ACT CCA TCG CAT TTG-Tao-IBDRQ		
crAssphage	DN A	CrAss-F	CAG AAG TAC AAA CTC CTA AAA AAC GTA GAG	Genomic base pair region: 14731 bp- 14856 bp	Stachler, Akyon, Carvalho, Ference, & Bibby, 2018
		CrAss-R	GAT GAC CAA TAA ACA AGC CAT TAG C		
		CrAss-P (Probe)	FAM-AAT AAC GAT TTA CGT GAT GTA AC		
<i>Escherichia coli</i>	DN A	uidA_784F	GTG TGA TAT CTA CCC GCT TCG C	uidA	Frahm and Obst, 2003
		uidA_866R	AGA ACG GTT TGT GGT TAA TCA GGA		
		EC807 probe	FAM-TCGGCATCCGGTCAGTGGCAGT- BHQ1		
Norovirus GI	RN A	Cog1F-flap	AATAAATCATAACGYTGATGCGNTT YCATGA	Genotype I	Kageyama, et al., 2003; Wang, et al., 2019 Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control, 2017b
		Cog1R-flap	AATAAATCATAACTAGACGCCATCA TCATTYAC		
		Ring1a.2 (Probe)	6-FAM- AGATYGCGR/ZEN/ TCYCCTGTCCA -IBFQ		
Norovirus GII	RN A	Cog2F-flap	AATAAATCATAACARGARBCNATGTT YAGRTGGAT GAG	Genotype II	Kageyama, et al., 2003; Wang, et al., 2019 Molecular Microbiology & Genomics Team, British Columbia Centre for Disease Control, 2017b
		Cog2R-flap	AATAAATCATAATCGACGCCATCTTC ATTCA		
		Ring 2.2 (Probe)	JOE - TGGGAGGGY/ZEN/ GATCGCAATCT - IBFQ		
Pepper Mild Mottle Virus	RN A	PMMV- FP1-rev	GAG TGG TTT GAC CTT AAC GTT TGA	1878 bp- 1901 bp ^a and 1945 bp-1926 bp ^a	Rosario, Symonds, Sinigalliano, Stewart, & Breitbart, 2009
		PMMV- RP1	TTG TCG GTT GCA ATG CAA GT		
		PMMV-P (Probe)	FAM-CCT ACC GAA GCA AAT G-MGB- NFQ		
Rotavirus Type A	RN A	NSP3-F	ACC ATC TWC ACR TRA CCC TCT ATG AG	Non- structural Protein 3	Zeng, et al., 2008
		NSP3-R	GGT CAC ATA ACG CCC CTA TAG C		
		NSP3-P (Probe)	VIC-AGT TAA AAG CTA ACA CTG TCA AA-MGB-NFQ		
Sapovirus	RN A	Sav1F	TTG GCC CTC GCC ACC TAC	Junction of polymerase and capsid	Oka, et al., 2006
		Sav5F	TTT GAA CAA GCT GTG GCA TGC TAC		
		Sav124F	GAY CAS GCT CTC GCY ACC TAC		
		Sav124R	CCC TCC ATY TCA AAC ACT A		
		Sav124TP (Probe)	FAM-CCR CCT ATR AAC CA-MGB-NFQ		
		Sav5TP (Probe)	FAM-TGC CAC CAA TGT ACC A-MGB- NFQ		

^aCorresponding nucleotide position of GenBank accession number M81413 (PMMoV strain S)

2

Figure 1

Figure 1. Graphical abstract of workflow.




Figure 2

Figure 2. Box plots of the number of gene copies of DNA enteric viruses across each wastewater stage throughout Events 1-4.

The unit for the SC in Figures 2A and 2C is gene copies per g of sample.





Figure 3

Figure 3. Box plots of the number of genes copies of PMMV across each wastewater stage throughout Events 1-4.

The unit for the SC in Figure 3A is gene copies per g of sample.

3A.



3B.




Figure 4

Figure 4. Box plots of the number of gene copies of *uidA* across each wastewater stage throughout Events 1-4.

The unit for the SC in Figure 4A is gene copies per g of sample.




Figure 5

Figure 5. Principal Component Analysis of \log_{10} -transformed EF parameters, PC1 versus PC2.

The only variable not \log_{10} -transformed was precipitation due to presence of zero values.




Figure 6

Figure 6. Heatmap showing Spearman's rank correlation analysis between parameters collected for EF sampling events.

