

Global Exercise - 14

Tuan Vo

25th January 2022

1 Conservation form - Finite Volume Method

Example 1. Determine consistent numerical flux function for Lax-Friedrichs scheme.

Recall the conservation form used for FVM reads

$$u_j^{n+1} = u_j^n + \frac{\Delta t}{\Delta x} \left(\tilde{f}_{j-1/2}^n - \tilde{f}_{j+1/2}^n \right). \quad (1)$$

Meanwhile, the Lax-Friedrichs scheme reads

$$u_j^{n+1} = \frac{1}{2} (u_{j-1}^n + u_{j+1}^n) - \frac{\Delta t}{2\Delta x} (f(u_{j+1}^n) - f(u_{j-1}^n)). \quad (2)$$

Next, we would like to write (2) in terms of (1). The derivation is done by some algebraic manipulations, as follows

$$\begin{aligned} u_j^{n+1} &= \frac{1}{2} (u_{j-1}^n + u_{j+1}^n) - \frac{\Delta t}{2\Delta x} (f(u_{j+1}^n) - f(u_{j-1}^n)) \\ &= u_j^n + \frac{1}{2}u_{j-1}^n - \frac{1}{2}u_j^n + \frac{1}{2}u_{j+1}^n - \frac{1}{2}u_j^n - \frac{\Delta t}{2\Delta x} f(u_{j+1}^n) + \frac{\Delta t}{2\Delta x} f(u_{j-1}^n) \\ &= u_j^n + \frac{\Delta t}{\Delta x} \left(\frac{\Delta x}{2\Delta t} u_{j-1}^n - \frac{\Delta x}{2\Delta t} u_j^n + \frac{\Delta x}{2\Delta t} u_{j+1}^n - \frac{\Delta x}{2\Delta t} u_j^n \right. \\ &\quad \left. - \frac{1}{2} f(u_{j+1}^n) + \frac{1}{2} f(u_{j-1}^n) \right) \\ &= u_j^n + \frac{\Delta t}{\Delta x} \left(\underbrace{\left(\frac{\Delta x}{2\Delta t} u_{j-1}^n - \frac{\Delta x}{2\Delta t} u_j^n + \frac{1}{2} f(u_{j-1}^n) \right)}_{=:\tilde{f}_{j-1/2}^n} \right. \\ &\quad \left. - \underbrace{\left(\frac{\Delta x}{2\Delta t} u_j^n - \frac{\Delta x}{2\Delta t} u_{j+1}^n + \frac{1}{2} f(u_{j+1}^n) \right)}_{=:\tilde{f}_{j+1/2}^n} \right) \\ &= u_j^n + \frac{\Delta t}{\Delta x} \left(\tilde{f}_{j-1/2}^n - \tilde{f}_{j+1/2}^n \right), \end{aligned} \quad (3)$$

which confirms that the *Lax-Friedrichs* scheme given at (2) is able to be written in the *conservation form* with the numerical flux function recognized as follows

$$\tilde{f}_{j-1/2}^n = \tilde{f}_{j-1/2}^n(u_{j-1}^n, u_j^n) = \frac{\Delta x}{2\Delta t} (u_{j-1}^n - u_j^n) + \frac{1}{2} f(u_{j-1}^n) \quad (4)$$

$$\tilde{f}_{j+1/2}^n = \tilde{f}_{j+1/2}^n(u_j^n, u_{j+1}^n) = \frac{\Delta x}{2\Delta t} (u_j^n - u_{j+1}^n) + \frac{1}{2} f(u_{j+1}^n) \quad (5)$$

However, these numerical flux function $\tilde{f}(\cdot, \cdot)$ is not consistent with the original flux function $f(\cdot)$, which can be checked $\forall \beta \in \mathbb{R}$ as follows

$$\tilde{f}_{j-1/2}^n(\beta, \beta) = \frac{\Delta x}{2\Delta t} (\beta - \beta) + \frac{1}{2} f(\beta) = \frac{1}{2} f(\beta) \quad \nless (6)$$

$$\tilde{f}_{j+1/2}^n(\beta, \beta) = \frac{\Delta x}{2\Delta t} (\beta - \beta) + \frac{1}{2} f(\beta) = \frac{1}{2} f(\beta) \quad \nless (7)$$

2 Godunov's method

Example 2. *Summary*

→ One-sided method cannot be used for system, i.e. mixed sign of eigenvalue causes difficulty.

→

3 Approximate Riemann Solvers

Example 3. *Linearized Riemann solvers - Roe Solver.*

Example 4. *Local Lax-Friedrichs flux.*

4 High resolution methods

Example 5. *Linearized Riemann solvers - Roe Solver.*