

Chrono-Energetics: An Emergent Approach to Spacetime Metric via Local Dissipation and Vacuum Stiffness

Gregory Lombardo

Independent Researcher, Alumnus of Classes Préparatoires (Maths Sup / Spé) & Engineering School, Theoretical Physics Enthusiast, Antibes, France

Abstract

We propose a novel conceptual framework, termed Chrono-Energetics, wherein time is not treated as a fundamental, passive coordinate of a geometric background, but as a macroscopic emergent property arising from local thermodynamic and quantum phenomena. We formalize this hypothesis through a local differential toy-model equation where the generation of the local temporal flow (dt) is directly proportional to the localized dissipated energy (dE_{diss}) and inversely proportional to the intrinsic structural stiffness of the quantum vacuum ($\mathcal{R}_v$), derived from Einstein's gravitational constant. By applying dimensional analysis, we introduce the fundamental unit of emergent time, the Chronom (χ), which naturally maps to the dimension of length ($1 \chi = 1 \text{ meter}$). This unified approach provides elegant alternative explanations for major cosmological anomalies: Cosmological Expansion, Galactic Rotation Curves (replicating MOND), and the Bullet Cluster Enigma. Finally, we propose three distinct tabletop experimental setups to test the validity of the Chronom in laboratory environments.

Keywords: Emergent Time, Quantum Vacuum Stiffness, Gravity as an Entropic Force, Dark Matter Alternatives, Chrono-Energetics, Thermodynamics of Spacetime

1. Introduction

The nature of time remains one of the most persistent enigmas in modern physics, sitting precisely at the irreconcilable fracture between General Relativity (GR) and Quantum Mechanics (QM). In Einstein's field equations, time is seamlessly woven with space into a smooth, continuous four-dimensional Lorentzian manifold. Conversely, in the standard formulation of quantum field theory, time recovers its Newtonian role: an external, rigid parameter ticking uniformly in the background, untouched by the quantum states it governs.

To bridge this conceptual chasm, several contemporary programs—most notably loop quantum gravity, thermal time hypotheses (Rovelli & Connes), and entropic gravity (Verlinde)—have suggested that spacetime metrics might be purely emergent properties. Following this trajectory,

this paper introduces the **Theory of Chrono-Energetics**. Our core postulate is that the local flow of time is a dynamic consequence of the macroscopic struggle between the arrow of time and the elastic resistance of the quantum vacuum.

2. Mathematical Framework and Dimensional Analysis of the Chronom

Let a physical system undergo an irreversible thermodynamic process or a quantum decoherence event, resulting in a localized dissipation of energy denoted by dE_{diss} . We formalize this relationship through the fundamental differential toy-model equation:

$$dt = \alpha \cdot \frac{dE_{\text{diss}}}{\mathcal{R}_v} \quad (1)$$

Where $\mathcal{R}_v$ represents the local effective stiffness of the vacuum, governed by the inverse of Einstein's gravitational constant κ :

$$\mathcal{R}_0 = \frac{c^4}{8\pi G} \approx 1.21 \times 10^{43} \text{ N} \quad (2)$$

Email address:

gregory.lombardo.research@email.com (Gregory Lombardo)

A profound property of the Lombardo equation appears when performing a rigorous dimensional analysis within the SI system:

$$\left[\frac{dE_{\text{diss}}}{\mathcal{R}_v} \right] = \frac{\text{Joules (J)}}{\text{Newtons (N)}} = \text{meter (m)} \quad (3)$$

This mathematical identity demonstrates that the fundamental unit of emergent time, which we name the **Chronom** (symbol: χ), possesses the intrinsic dimension of a **length** ($1 \chi = 1 \text{ meter}$).

To map this interval to the macroscopic second (s), we set $\alpha = \frac{1}{c}$, yielding:

$$dt_{\text{seconds}} = \frac{8\pi G}{c^5} \cdot dE_{\text{diss}} \quad (4)$$

3. Galactic Rotation Curves and MOND Equivalence

We hypothesize that the vacuum stiffness $\mathcal{R}_v$ is not a global constant, but a local field variable $\mathcal{R}_v(r)$ that depends on the surrounding baryon density:

$$\mathcal{R}_v(r) = \mathcal{R}_0 \cdot \left[\frac{1 + \left(\frac{r}{r_c}\right)^2}{1 + \left(\frac{r}{r_c}\right)^2 \cdot \frac{a_0}{g_N(r)}} \right] \quad (5)$$

When a star orbits at the outer edge of a galaxy ($r \gg r_c$), the Newtonian acceleration drops ($g_N(r) \ll a_0$). In this low-acceleration regime, the effective resistance of space sags, modifying the effective gravitational acceleration $g_{\text{eff}}(r)$:

$$g_{\text{eff}}(r) = \sqrt{g_N(r) \cdot a_0} = \frac{\sqrt{GM_{\text{bar}}a_0}}{r} \quad (6)$$

By equating $g_{\text{eff}}(r)$ to the centripetal acceleration $\frac{v^2}{r}$, the spatial dependence r cancels out perfectly, deriving the **Baryonic Tully-Fisher relation**:

$$v^4 = GM_{\text{bar}}a_0 \quad (7)$$

4. The Bullet Cluster and Vacuum Hysteresis

The Lombardo framework resolves the Bullet Cluster paradox by introducing a temporal retardation parameter into the field equation of vacuum stiffness. We postulate that the quantum vacuum behaves like a non-ideal elastic medium exhibiting **geometric hysteresis**. The stiffness field $\mathcal{R}_v(\mathbf{x}, t)$

relaxes according to a characteristic finite time constant τ_v :

$$\frac{\partial \mathcal{R}_v(\mathbf{x}, t)}{\partial t} = -\frac{1}{\tau_v} \left[\mathcal{R}_v(\mathbf{x}, t) - \mathcal{R}_{v,\text{static}}(\rho(\mathbf{x}, t)) \right] \quad (8)$$

During the ultra-rapid collision, the baryonic plasma is strongly decelerated by hydrodynamic friction. However, because $\tau_v > 0$, the structural profile of softened vacuum continues to propagate along the original inertial trajectories, decoupling the weak lensing signal from the physical location of the gas without requiring dark matter particles.

5. Conclusion & Experimental Perspectives

The Theory of Chrono-Energetics reveals that time generation is fundamentally equivalent to a spatial metric extension regulated by energy dissipation. To test the validity of the Chronom, we propose three experimental setups:

- **Experiment A:** High-Entropy Laser Interferometry to measure optical phase shifts induced by localized gas ionization.
- **Experiment B:** Casimir-Constrained Qubit Decoherence to observe changes in temporal flow near conductive plates.
- **Experiment C:** Cryogenic Optical Lattice Clocks to detect micro-decelerations in internal rhythm near absolute zero.