CONTENTS

# Final

## Yuechen Liu, Mufeng Xu, Yanhao Li

# Contents

Introduction	2
Load, clean, and tidy data	2
Exploratory analysis/ visualization	3

```
library(tidyverse)
library(caret)
library(glmnet)
library(ISLR)
library(pls)
library(AppliedPredictiveModeling)
library(MASS)
library(mHASS)
library(mlbench)
library(pROC)
library(arsenal)
```

#### Introduction

Stroke is a serious life-threatening medical condition. According to the World Health Organization, stroke is the second leading cause of death globally. To better understand which factors correlate to the stroke event, our group find a stroke prediction dataset. This dataset contains twelve columns. The first column labels the unique identifier of the patient. The last column records the occurrence of stroke by 1 (Yes) or 0 (No). The other ten columns contain the observations of possible predictors.

### Load, clean, and tidy data

```
stroke = read_csv("./healthcare-dataset-stroke-data.csv") %>%
 mutate(
    bmi = as.numeric(bmi)
 )
stroke1 = stroke %>%
  janitor::clean_names() %>%
  filter(
    bmi != "N/A"
  ) %>%
  mutate(
    gender = recode(
      gender,
      "Male" = 0,
      "Female" = 1,
      "0ther" = 2
    ),
    ever_married = recode(
      ever_married,
      "No" = 0,
      "Yes" = 1
    ),
    work_type = recode(
      work_type,
      "children" = 0,
      "Govt job" = 1,
      "Never_worked" = 2,
```

```
"Private" = 3,
      "Self-employed" = 4
   residence_type = recode(
     residence_type,
      "Rural" = 0,
      "Urban" = 1
   ),
   smoking_status = recode(
      smoking_status,
      "formerly smoked" = 0,
      "never smoked" = 1,
      "smokes" = 2,
      "Unknown" = 3
   ),
   stroke = as.factor(stroke)
  ) %>%
 relocate(
   age, avg_glucose_level, bmi
stroke2 = stroke1 %>%
 mutate(
   gender = as.factor(gender),
   hypertension = as.factor(hypertension),
   heart_disease = as.factor(heart_disease),
   ever_married = as.factor(ever_married),
   work_type = as.factor(work_type),
   residence_type = as.factor(residence_type),
   smoking_status = as.factor(smoking_status)
```

## Exploratory analysis/ visualization

```
stats = tableby(stroke ~ gender + age + hypertension + heart_disease + ever_married + work_type + Resid
summary(stats, text = TRUE) %>% knitr::kable()
```

	0 (N=4861)	1 (N=249)	Total ( $N=5110$ )	p value
gender				0.790
- Female	2853 (58.7%)	141~(56.6%)	2994 (58.6%)	
- Male	2007 (41.3%)	108 (43.4%)	2115 (41.4%)	
- Other	1(0.0%)	0 (0.0%)	1(0.0%)	
age				< 0.001
- Mean (SD)	41.972 (22.292)	67.728 (12.727)	43.227 (22.613)	
- Range	0.080 - 82.000	1.320 - 82.000	0.080 - 82.000	
hypertension				< 0.001
- Mean (SD)	0.089 (0.285)	0.265 (0.442)	0.097(0.297)	
- Range	0.000 - 1.000	0.000 - 1.000	0.000 - 1.000	
heart_disease				< 0.001

	0  (N=4861)	1 (N=249)	Total (N=5110)	p value
- Mean (SD)	0.047(0.212)	0.189(0.392)	0.054 (0.226)	
- Range	0.000 - 1.000	0.000 - 1.000	0.000 - 1.000	
ever_married				< 0.001
- No	$1728 \ (35.5\%)$	29~(11.6%)	1757 (34.4%)	
- Yes	3133~(64.5%)	220~(88.4%)	3353~(65.6%)	
work_type				< 0.001
- children	685 (14.1%)	2(0.8%)	687 (13.4%)	
- Govt_job	$624\ (12.8\%)$	$33\ (13.3\%)$	$657\ (12.9\%)$	
- Never_worked	$22 \ (0.5\%)$	0 (0.0%)	22 (0.4%)	
- Private	2776 (57.1%)	149 (59.8%)	2925~(57.2%)	
- Self-employed	$754 \ (15.5\%)$	65~(26.1%)	$819\ (16.0\%)$	
$Residence\_type$				0.269
- Rural	$2400 \ (49.4\%)$	114 (45.8%)	2514 (49.2%)	
- Urban	2461~(50.6%)	135 (54.2%)	2596 (50.8%)	
$avg\_glucose\_level$				< 0.001
- Mean (SD)	$104.796 \ (43.846)$	132.545 (61.921)	$106.148 \ (45.284)$	
- Range	55.120 - 267.760	56.110 - 271.740	55.120 - 271.740	
bmi				0.003
- N-Miss	161	40	201	
- Mean (SD)	$28.823 \ (7.908)$	$30.471 \ (6.329)$	28.893 (7.854)	
- Range	10.300 - 97.600	16.900 - 56.600	10.300 - 97.600	
$smoking\_status$				< 0.001
- formerly smoked	815 (16.8%)	70 (28.1%)	885 (17.3%)	
- never smoked	1802 (37.1%)	90 (36.1%)	1892 (37.0%)	
- smokes	$747 \ (15.4\%)$	42 (16.9%)	$789 \ (15.4\%)$	
- Unknown	1497 (30.8%)	47 (18.9%)	$1544 \ (30.2\%)$	

