

Instructions for Administration:

1 Month Old

Note Ensure to wash your hands before handling the child and to handle the child gently. For neonates and young infants (<3 months) offer a little bit of time after testing each item, to allow the infant to settle.

Definitions

Plane: In reference to items requiring holding a body part on the same plane as another, this refers to being level to.

Prone Position: The prone position refers to lying horizontally with the face and torso facing down. The infant lies with knees drawn up under the abdomen with pelvis high. Head is turned to one side.

Supine Position: The supine position refers to lying horizontally with the face and torso facing up.

Ventral Suspension: The baby is placed in prone position and suspended in the air by a hand placed under the chest. The head hangs down with some flexion of elbows and knees.


1.1 Gross Motor

Notes If the child does not perform the gross motor item for their age, but performs a more advanced gross motor item, then the item specific for their age is considered normal.

The first step towards the development of locomotion is the development of head control, that is, the ability to support the head in all positions of the body. The steps in this are observed in three situations: in ventral suspension, in prone position, and in supine position.

1.1.6	Writhing movement of four limbs in supine position	Age: 1 - 2 months
Materials	Padded Surface	
Instructions	Observe the infant in supine position. Note spontaneous movements for a couple of minutes, preferably while the infant is awake and quiet. Observe both the quality and quantity of movements. Evaluate the patterns and the sequence of movements of the whole body, rather than concentrating on single segments, such as arms and legs.	
Response	The movements should be fluent and alternating.	
Grading	Present Sluggish No movement OR Asymmetric OR Exaggerated	

1.1.7	Lifts chin up momentarily when in prone position	Age: 1 month
Materials	Padded Surface	
Overview	The first step towards the development of locomotion is the development of head control – that is, the ability to support the head in all positions of the body. The steps in this are observed in three situations: in ventral suspension, in prone position, and in supine position.	
Instructions	The infant is placed in prone position with the head in the midline, and the movements of the head are noted for about 30 seconds. 	
Response	Lifts chin up momentarily when lying in prone position.	
Grading	Present Rolls head over, chin not raised Feeble movement of head No response OR persistently raised head	

1.1.8	Lifts head up momentarily in ventral suspension	Age: 1 month
Materials	Padded Surface	
Overview	The first step towards the development of locomotion is the development of head control – that is, the ability to support the head in all positions of the body. The steps in this are observed in three situations: in ventral suspension, in prone position, and in supine position.	
Instructions	From prone position, lift the infant horizontally with one or both hands under the chest so they are in ventral suspension, about 20-25cm from a surface. Note the infant's attempt to lift the head for about 30 seconds. 	
Response	Lifts head up momentarily, within 30 seconds, but not in line with the body.	
Grading	Present Moves head but does not raise it Feeble movement of head No response OR persistently raised head	

1.1.9	Lifts head forward on pull to sitting	Age: 1 month
Materials	Padded Surface	
Instructions	With the infant in supine position, grasp the shoulders and pull the infant towards a sitting position. Posture of the head is noted, when the shoulders are elevated to about 45 degrees. 	
Response	Lifts head slightly on pull to sitting.	
Grading	Present Shows effort, but drops back Feeble effort No Effort	

1.2 Fine Motor Item

Note If the child does not perform the fine motor item for their age, but performs a more advanced fine motor item, then the item specific for their age is considered normal.

1.2.1	Hand posture		Age: Birth - 1 month
Materials	None		
Instructions	Observe the infant when awake and alert, and at rest.		
Response	Intermittent fisting, with thumbs in or out of the palms.		
Grading	Present Atypical Hand Posture, either: a) continuous fisting OR thumb adduction b) index finger flexion, thumb opposition		
Reference	Typical Hand Posture 	Atypical Hand Posture A) <i>Thumb Adduction</i> 	Atypical Hand Posture B) <i>Opposition of index finger and thumb (other three fingers extended)</i> 

1.3 Vision Items

1.3.4	Regards face	Age: 1 month
Materials	If no response to face: Compact Mirror, Torch	
Instructions	Ensure the room is well-lit. The infant may be held on the examiner's lap or tested in a propped up supine position. With the examiner's face 20-25cm in front of the infant's face, move slowly in a horizontal and vertical arc. If no response, then test with a reflecting object at a distance of 15-25 cm. If still no response, test with a light source (e.g., torch). Avoid talking to the infant during this item.	
Response	Regards face.	
Grading	Present Face regard questionable Regards reflecting object Regards light source or less	



1.3.5	Regards woolly ball	Age: 1 month
Materials	Red Woolly Ball If no response to Red Woolly Ball: Compact Mirror, Torch	
Instructions	Ensure the room is well-lit. The infant is tested in a propped up supine position and the stimulus is presented at a distance of 15 - 25cm, starting in the midline and moving laterally in either direction, then vertically. If no response, then test with a reflecting object at a distance of 15 - 25 cm. If still no response, then test with a light source (e.g., torch).	
Response	Regards the woolly ball.	
Grading	Present Questionable Regards reflecting object Regards light source or less	



1.4 Hearing Items

Note Soft whisper - 10 to 25 db; conversational speech - 25 to 55 db; crying baby - 55 to 65 db; rattle - 60 to 80db.

1.4.4	Response to clapper bell	Age: 1 - 3 months
1.4.5	Response to rattle	
1.4.6	Response to voice	
Materials	Clapper Bell, Rattle, Assistant, Padded Surface	
Instructions	The infant may be held in a propped up supine position on a padded surface, or on the caregiver's lap. If the infant is able to hold the neck up, they may be positioned sitting in the caregiver's lap. The head should be supported in the midline if in supine position. T The auditory stimulus is: (a) presented at the ear level, (b) about 15 cm from the ear, (c) out of the line of vision. Test the right ear, then the left ear. A trained assistant provides the stimulus, and the tester observes the response. 	
Response	Infant will brighten/startle/eyes will widen/become still/show respiratory changes/blink/shift of eyes to sound. The best response to any one of the three sound sources is taken as normal.	
Grading	Present Questionable Response to Louder Ring/Louder Rattle/Voice No response	

1.5 Expressive Language Item

Note This domain relies primarily on observation of infant vocalisations during the session:

- The examiner has to be vigilant about recording all infant vocalisations during the entire assessment so that none are missed.
- Interact and speak with the child to encourage vocalisations.
- If no vocalisations are made throughout the assessment, ask the caregiver if this is typical as it is not uncommon for young children to behave differently in new and unfamiliar situations. Document specific examples of vocalisations reported by the caregiver on the record form.


1.5.1	Loudness of cry	Age: Birth – 1 month
Materials	None	
Instructions	Crying in the newborn usually falls within a typical range for pitch and loudness.	
Response	Note the typicality of the loudness of the crying	
Grading	Typical/Within normal limit Poor Crying Sound No Crying Sound	

1.6 Cognition Items

1.6.4	Brightens to face	Age: 1 month
Materials	None	
Instructions	The infant may be held on the examiner's lap or tested in a propped up supine position. With the examiner's face 30-40cm in front of the infant's face, move slowly in a horizontal and vertical arc. Avoid talking to the infant during this item.	
Response	Observe if the infant alerts to the face, which may be expressed as a brightening of the expression and/or widening of the eyes.	
Grading	Present Weak Questionable None	



1.6.5	Brightens to voice	Age: 1 month
Materials	Assistant	
Instructions	The infant is tested in a propped up supine position or held in the caregiver's lap. The assistant speaks softly into one of the infant's ears outside of the infant's line of vision, about 15-30cm away from their face. The sounds of the voice may be varied in pitch and intensity to obtain the infant's attention and avoid habituation. Usually, a soft high-pitched voice is the most potent stimulus.	
Response	Observe if the infant alerts to the voice, which may be expressed as brightening of expression and/or widening of the eyes.	
Grading	Present Weak Questionable None	



1.6.6	Brightens to touch	Age: 1 month
Materials	None	
Instructions	Touch and stroke the infant gently over the arms and face. 	
Response	Observe if the infant alerts to the touch, which may be expressed as brightening of expression and/or widening of the eyes.	
Grading	Present Weak Questionable None	

1.7 Behaviour Items

Note It is essential the examiner scores the Behaviour items based on their observations and caregiver report regarding sleep. Observe and record the child's behaviour according to the activities listed under this domain.

1.7.5	Normal and consolable patterns of crying	Age: 1 - 2 months
Materials	None	
Instructions	This measures the infant's ability to be consoled once they have reached a crying state, either spontaneously or by various manoeuvres (e.g., cooing, rocking, feeding, changing nappy) carried out by the caregiver.	
Response	Mild, moderate, and severe grades of response are: sometimes consolable, achieved with difficulty, and unconsolable.	
Grading	Present Sometimes consolable Achieved with difficulty Unconsolable	

1.7.6	Sleep/awake rhythm	Age: 1 - 11 months
Materials	None	
Instructions	Ask the caregiver about the infant's sleep patterns.	
Response	Score according to caregiver's recall. Over the first 3 months of life the total amount of sleep declines little, but the cycles are longer so that, by the age of 3 months, 70% of infants are sleeping right through the night. By 6 months, 80% of infants sleep through the night, but there are still 10% waking every night by the age of 1 year. By this age, the great majority of infants have established a stable pattern of sleep and wakefulness with a long sleep at night and a nap in the morning and afternoon. Mild, moderate and severe grades of response are: occasional disturbance, considerable disturbance, and constant problem. Either the infant is sleeping too much or is awake for longer periods of time.	
Grading	Normal sleep patterns Occasional disturbance Considerable disturbance Constant problems	

1.7.7	Displays positive affect most of the time	Age: 1 - 11 months
Materials	None	
Instructions	Note the infant's affect.	
Response	The grading is done according to the degree of negative affect displayed during the assessment. Positive affect comprises of smiling, laughing, sound making that is perceived as the expression of excitement, happiness or pleasure. Negative affect includes fussing, pouting, whining, crying, vocal or physical expressions of anger.	
Grading	Displays positive affect Sometimes negative Mostly negative Constantly negative	

1.8 Seizure Item

1.8.2	Seizures (provoked or unprovoked)	Age: 1 - 2 months
Materials	None	
Instructions	Ask caregiver or check medical records to determine whether infant has ever experienced a seizure. Seizures are manifestations of neuronal injury due to excessive electrical activity in the brain. Any disturbance in the brain may cause seizures, which may be single or repetitive, and involves a sudden change in behaviour.	
Response	Consider typical if the infant has not had a provoked or unprovoked seizure.	
Grading	No seizures (provoked or unprovoked) Once, but none in the past two weeks Once in the past two weeks OR > once, but none in past two weeks > Once, also in past two weeks	