



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Chronic Lumbar Disc Herniation with Sciatica

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #22

Clinical Details

Doctor: Dr. PSP

Date: 2026-04-28

Description: Patient presents with persistent radiating pain from the lower back down to the right calf. Diagnosed with L4-L5 disc protrusion. Pain is aggravated by prolonged sitting and forward bending.

Physical Measurements

Height: 178 cm

Weight: 85 kg

Blood Group: APositive

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Moderate

Smart Clinical Analysis

Clinical Explanation:

The patient presents with persistent radiating pain from the lower back to the right calf, attributed to an L4-L5 disc protrusion. This condition is characterized by nerve root irritation, leading to radicular symptoms. The pain exacerbates with prolonged sitting and forward bending, indicating potential mechanical stress on the lumbar spine. The patient's moderate activity level and absence of chronic diseases suggest a favorable prognosis for rehabilitation.

Immediate Physiotherapy Recommendations:

A tailored physiotherapy program focusing on core stabilization, lumbar mobilization, and flexibility exercises is recommended. Emphasis should be placed on educating the patient about proper body mechanics and posture to alleviate strain on the lumbar region. Gradual progression of exercises should be encouraged based on tolerance.

Precautions:

The patient should avoid activities that involve heavy lifting, prolonged sitting, and forward bending, as these may exacerbate symptoms. Care should be taken to monitor pain levels during rehabilitation to prevent aggravation of the condition.

Estimated Recovery Timeline:

The estimated recovery timeline for this condition typically ranges from 6 to 12 weeks, depending on adherence to the rehabilitation program and individual response to treatment.

Risk Assessment:

Potential risks include worsening of symptoms, development of chronic pain, or progression of neurological deficits if the condition is not managed appropriately. Regular follow-up assessments are essential to monitor progress.

Red Flags:

Emergency warning signs include sudden onset of severe pain, loss of bowel or bladder control, significant weakness in the lower extremities, or any new neurological symptoms. Immediate medical evaluation is warranted if these occur.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:
Normal knee flexion range for adults is typically between 0 to 135 degrees.

Clinical Interpretation:
The recorded values indicate a progressive improvement in knee flexion on the right side, moving from 95 degrees to 115 degrees over the assessment period. This suggests an increase in range of motion, likely due to effective rehabilitation strategies.

Progress Evaluation:
The trend shows a consistent increase in knee flexion, indicating positive recovery and improved functional mobility. Continued monitoring is necessary to ensure this trend persists.

Physiotherapy Focus:
Focus on maintaining and enhancing knee flexion through targeted stretching and strengthening exercises, while ensuring lumbar stability is prioritized to prevent exacerbation of lower back symptoms.


HIGHEST
115.0

LOWEST
95.0

AVERAGE
106.0

OVERALL PROGRESS

Increased by 21.1% ↑ Since first test (2026-04-14)



Date	Result
2026-04-14	95
2026-04-18	100
2026-04-22	110
2026-04-26	110
2026-04-28	115