



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Post-ACL Reconstruction Rehabilitation

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #24

Clinical Details

Doctor: Dr. PSP

Date: 2026-03-28

Description: A 28-year-old male athlete, 4 weeks post-op for ACL reconstruction using a hamstring graft. Currently experiencing mild swelling and limited range of motion (ROM) in the right knee.

Physical Measurements

Height: 182 cm

Weight: 78 kg

Blood Group: OPositive

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Active

Additional Patient Info

InjuryHistory: Complete ACL tear occurred during a football match on January 15, 2026, due to a sudden pivoting motion. Surgery was performed on March 1, 2026.

Medications: Ibuprofen 400mg as needed for pain, Vitamin D3 supplements, and Omega-3 fish oil.

FunctionalAbility: Partial weight-bearing with one crutch. Knee flexion is currently at 95 degrees. Struggles with stairs and long-distance walking.

PersonalGoals: Full return to competitive football within 9 months, restoration of full quad strength, and achieving 135-degree knee flexion.

Smart Clinical Analysis

Clinical Explanation:

The patient is a 28-year-old male athlete who is currently four weeks post-operative following an ACL reconstruction using a hamstring graft. He presents with mild swelling and limited range of motion in the right knee, which is common during the early stages of rehabilitation. The patient has a functional ability of partial weight-bearing with the assistance of one crutch and has demonstrated a gradual improvement in knee flexion. His personal goals include a full return to competitive football within nine months, restoration of full quadriceps strength, and achieving 135 degrees of knee flexion.

Immediate Physiotherapy Recommendations:

The patient should continue with a structured rehabilitation program focusing on range of motion exercises, strengthening of the quadriceps and hamstrings, and proprioceptive training. Modalities such as ice therapy may be utilized to manage swelling. Gradual progression to weight-bearing activities is advised, with an emphasis on maintaining proper biomechanics during movements.

Precautions:

The patient should avoid high-impact activities, pivoting motions, and any exercises that cause pain or discomfort in the knee. Care should be taken to monitor swelling and pain levels during rehabilitation sessions.

Estimated Recovery Timeline:

Given the patient's current status and goals, a full recovery with the ability to return to competitive football may be anticipated within 6 to 9 months post-surgery, contingent upon adherence to rehabilitation protocols and absence of complications.

Risk Assessment:

Potential risks include re-injury of the ACL, development of stiffness or scar tissue, and delayed healing due to inadequate rehabilitation. Close monitoring of knee function and symptoms is essential.

Red Flags:

Emergency warning signs include severe swelling, increased pain not alleviated by medication, inability to bear weight on the affected leg, signs of infection (such as fever or redness), and any sudden changes in knee stability.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:

Knee flexion range for a healthy adult typically ranges from 0 to 135 degrees.

Clinical Interpretation:

The patient has shown a gradual increase in knee flexion from 95 degrees to 115 degrees over the assessment period, indicating positive progress in rehabilitation. However, the current measurement of 115 degrees is still below the normal reference range.

Progress Evaluation:

The trend shows consistent improvement in knee flexion, with the most recent measurements indicating a steady increase. This suggests effective rehabilitation strategies are being implemented.

Physiotherapy Focus:

Continue to emphasize range of motion exercises, particularly focusing on flexion. Incorporate gentle stretching and mobilization techniques to facilitate further improvement in knee flexion.

HIGHEST

115.0

LOWEST

95.0


AVERAGE

106.0

OVERALL PROGRESS

Increased by 21.1% ↑

Since first test (2026-04-01)



Date	Result
2026-04-01	95
2026-04-05	100
2026-04-10	110
2026-04-20	110
2026-04-26	115