



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Post-ACL Reconstruction Rehabilitation

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #24

Clinical Details

Doctor: Dr. PSP

Date: 2026-03-28

Description: A 28-year-old male athlete, 4 weeks post-op for ACL reconstruction using a hamstring graft. Currently experiencing mild swelling and limited range of motion (ROM) in the right knee.

Physical Measurements

Height: 182 cm

Weight: 78 kg

Blood Group: OPositive

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Active

Additional Patient Info

InjuryHistory: Complete ACL tear occurred during a football match on January 15, 2026, due to a sudden pivoting motion. Surgery was performed on March 1, 2026.

Medications: Ibuprofen 400mg as needed for pain, Vitamin D3 supplements, and Omega-3 fish oil.

FunctionalAbility: Partial weight-bearing with one crutch. Knee flexion is currently at 95 degrees. Struggles with stairs and long-distance walking.

PersonalGoals: Full return to competitive football within 9 months, restoration of full quad strength, and achieving 135-degree knee flexion.

Smart Clinical Analysis

Clinical Explanation:

The patient is a 28-year-old male athlete who is four weeks post-operative following an anterior cruciate ligament (ACL) reconstruction using a hamstring graft. He presents with mild swelling and limited range of motion (ROM) in the right knee, which is common during the early stages of rehabilitation. The patient has a history of a complete ACL tear sustained during a football match, and his current rehabilitation goals include a full return to competitive football within nine months, restoration of full quadriceps strength, and achieving 135 degrees of knee flexion.

Immediate Physiotherapy Recommendations:

The focus of physiotherapy should include gentle range of motion exercises to improve knee flexion, as well as isometric quadriceps strengthening exercises. Gradual progression to weight-bearing activities should be encouraged, while monitoring for pain and swelling. Incorporating modalities such as ice and elevation may assist in managing swelling.

Precautions:

The patient should avoid high-impact activities, sudden pivoting motions, and any exercises that cause significant pain or discomfort. Close monitoring of swelling and pain levels during rehabilitation is essential.

Estimated Recovery Timeline:

Considering the current progress and rehabilitation goals, a full recovery to competitive football may be anticipated within 6 to 9 months post-surgery, contingent upon adherence to rehabilitation protocols and absence of complications.

Risk Assessment:

Potential risks include re-injury to the knee, development of stiffness, or complications related to the surgical site. It is crucial to ensure that the patient does not rush the rehabilitation process.

Red Flags:

Emergency warning signs include severe swelling, increased pain not alleviated by rest or medication, inability to bear weight on the affected leg, signs of infection (such as fever or redness), and any unusual popping or locking sensations in the knee.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:
Normal knee flexion for a healthy adult is typically between 135 to 150 degrees.

Clinical Interpretation:
The patient currently demonstrates knee flexion measurements ranging from 95 to 115 degrees, indicating a gradual improvement in range of motion since surgery.

Progress Evaluation:
The trend shows a positive progression in knee flexion, with measurements increasing from 95 degrees to 115 degrees over the assessment period. This suggests effective rehabilitation and adherence to physiotherapy protocols.


Physiotherapy Focus:
Continue to emphasize range of motion exercises and consider incorporating more dynamic activities as tolerated, while ensuring that the patient maintains proper form and technique to prevent further injury.

HIGHEST
115.0

LOWEST
95.0

AVERAGE
106.0

OVERALL PROGRESS
Increased by 21.1% ↑ Since first test (2026-04-01)



Date	Result
2026-04-01	95
2026-04-05	100
2026-04-10	110
2026-04-20	110
2026-04-26	115