



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Post-ACL Reconstruction Rehabilitation

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #24

Clinical Details

Doctor: Dr. PSP

Date: 2026-03-28

Description: A 28-year-old male athlete, 4 weeks post-op for ACL reconstruction using a hamstring graft. Currently experiencing mild swelling and limited range of motion (ROM) in the right knee.

Physical Measurements

Height: 182 cm

Weight: 78 kg

Blood Group: OPositive

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Active

Additional Patient Info

InjuryHistory: Complete ACL tear occurred during a football match on January 15, 2026, due to a sudden pivoting motion. Surgery was performed on March

1, 2026.

Medications: Ibuprofen 400mg as needed for pain, Vitamin D3 supplements, and Omega-3 fish oil.

FunctionalAbility: Partial weight-bearing with one crutch. Knee flexion is currently at 95 degrees. Struggles with stairs and long-distance walking.

PersonalGoals: Full return to competitive football within 9 months, restoration of full quad strength, and achieving 135-degree knee flexion.

Smart Clinical Analysis

Clinical Explanation:

The patient is a 28-year-old male athlete who is currently 4 weeks post-operative following an anterior cruciate ligament (ACL) reconstruction using a hamstring graft. He presents with mild swelling and limited range of motion (ROM) in the right knee, which is common during the early stages of rehabilitation. The patient has a functional ability of partial weight-bearing with the assistance of one crutch and is experiencing challenges with activities such as stair climbing and long-distance walking. His rehabilitation goals include a full return to competitive football within 9 months, restoration of full quadriceps strength, and achieving 135 degrees of knee flexion.

Immediate Physiotherapy Recommendations:

The focus of physiotherapy should include progressive range of motion exercises, gentle strengthening of the quadriceps and hamstrings, and modalities to manage swelling. Emphasis should be placed on functional activities to improve mobility and confidence in weight-bearing.

Precautions:

The patient should avoid high-impact activities, sudden pivoting motions, and excessive loading of the knee joint until adequate strength and stability are achieved. Care should be taken to monitor for any signs of increased swelling or pain during rehabilitation exercises.

Estimated Recovery Timeline:

Given the current progress and goals, a full recovery to competitive football is anticipated within 9 months, contingent upon adherence to rehabilitation protocols and absence of complications.

Risk Assessment:

Potential risks include re-injury of the ACL, development of stiffness in the knee joint, and muscle atrophy if rehabilitation is not appropriately managed. Close monitoring of swelling and pain levels is essential.

Red Flags:

Emergency warning signs include severe swelling, increased pain not alleviated by rest or medication, inability to bear weight on the affected leg, and any signs of infection such as fever or redness around the surgical site.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:
0-135 degrees

Clinical Interpretation:
The patient currently demonstrates knee flexion ranging from 95 to 115 degrees, indicating a gradual improvement in range of motion since surgery. The current measurements suggest that the patient is progressing towards the normal range but has not yet achieved full flexion.

Progress Evaluation:
The trend shows a positive increase in knee flexion from 95 degrees to 115 degrees over the assessment period, indicating effective rehabilitation efforts and a favorable response to therapy.


Physiotherapy Focus:
Continue to emphasize range of motion exercises, including passive and active-assisted flexion, while monitoring for any signs of discomfort or swelling. Incorporate strengthening exercises for the quadriceps and hamstrings as tolerated.

HIGHEST
115.0

LOWEST
95.0

AVERAGE
106.0

OVERALL PROGRESS
Increased by 21.1% ↑ Since first test (2026-04-01)



Date	Result
2026-04-01	95
2026-04-05	100
2026-04-10	110
2026-04-20	110
2026-04-26	115