



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Chronic Lumbar Disc Herniation with Sciatica

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #22

Clinical Details

Doctor: Dr. PSP

Date: 2026-04-28

Description: Patient presents with persistent radiating pain from the lower back down to the right calf. Diagnosed with L4-L5 disc protrusion. Pain is aggravated by prolonged sitting and forward bending.

Physical Measurements

Height: 178 cm

Weight: 85 kg

Blood Group: APositive

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Moderate

Smart Clinical Analysis

Clinical Explanation:

The patient presents with persistent radiating pain from the lower back to the right calf, consistent with an L4-L5 disc protrusion. The pain is exacerbated by prolonged sitting and forward bending, indicating a mechanical component to the pain. The patient's moderate activity level and absence of chronic diseases suggest a potential for effective rehabilitation. The focus will be on pain management, improving range of motion, and strengthening the lumbar and pelvic regions to support spinal stability.

Immediate Physiotherapy Recommendations:

A structured physiotherapy program should include:

- Core stabilization exercises to enhance lumbar support.
- Flexibility exercises targeting the hamstrings and hip flexors to alleviate tension on the lumbar spine.
- Gradual introduction of aerobic conditioning to improve overall fitness without exacerbating symptoms.

Precautions:

- Avoid prolonged sitting and forward bending activities that may aggravate symptoms.
- Monitor for any signs of increased pain or neurological symptoms during exercises.
- Ensure proper body mechanics during daily activities to prevent strain.

Estimated Recovery Timeline:

With adherence to the rehabilitation program, a significant improvement may be observed within 6 to 12 weeks, although full recovery may take longer depending on individual response to therapy.

Risk Assessment:

Potential risks include exacerbation of symptoms due to improper exercise execution or overexertion. Close monitoring of pain levels and functional abilities is essential.

Red Flags:

- Sudden onset of severe pain.
- Loss of bowel or bladder control.
- Progressive weakness in the lower extremities.
- Numbness or tingling in the saddle area.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:

Normal knee flexion range for adults is typically between 0 to 135 degrees.

Clinical Interpretation:

The recorded values indicate a progressive improvement in knee flexion from 95 degrees to 115 degrees over the assessment period, suggesting enhanced mobility and function in the right knee.

Progress Evaluation:

The trend shows a consistent increase in knee flexion, indicating positive recovery and adaptation to the rehabilitation program.

Physiotherapy Focus:

Continue to emphasize knee flexion exercises and incorporate functional movements that promote strength and stability in the lower extremity.

HIGHEST

115.0

LOWEST

95.0


AVERAGE

106.0

OVERALL PROGRESS

Increased by 21.1% ↑

Since first test (2026-04-14)



Date	Result
2026-04-14	95
2026-04-18	100
2026-04-22	110
2026-04-26	110
2026-04-28	115