



Physiotherapy & Rehabilitation Platform

Medical Report: Smeer Alarkany

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #25

Clinical Details

Doctor: Dr. HAKIMALI

Date: 2026-07-13

Description: Work Tracking System

Physical Measurements

Height: 175 cm

Weight: 70 kg

Blood Group: AB+

Health History

Smoker: Yes

Chronic Diseases: Yes

Activity Level: Sedentary

Additional patient information

Injury History: test

Medications: test

Functional Ability: test

Personal Goals: tes

Smart Clinical Analysis

Clinical Explanation:

The patient presents with a sedentary lifestyle and a history of chronic diseases, which necessitates a cautious approach to rehabilitation. The primary focus will be on improving functional ability and mobility while considering the patient's overall health status. The rehabilitation strategy should incorporate gradual progression in physical activity to enhance strength and range of motion while minimizing the risk of exacerbating any chronic conditions.

Immediate Physiotherapy Recommendations:

A tailored physiotherapy program should be initiated, focusing on low-impact exercises to improve joint mobility and strength. Emphasis should be placed on flexibility and stability exercises, particularly for the lower extremities, to support functional movements.

Precautions:

Due to the patient's chronic diseases and sedentary lifestyle, it is crucial to monitor vital signs during exercise. Avoid high-impact activities and ensure that exercises are performed within a pain-free range. Regular assessments should be conducted to adjust the rehabilitation plan as needed.

Estimated Recovery Timeline:

The estimated recovery timeline may vary based on the patient's adherence to the rehabilitation program and response to treatment. A general timeline of 6 to 12 weeks is anticipated for noticeable improvements in mobility and strength, contingent upon consistent engagement in therapy.

Risk Assessment:

Potential risks include exacerbation of chronic conditions, joint instability, and the possibility of falls due to reduced physical activity. Close monitoring is essential to mitigate these risks.

Red Flags:

Emergency warning signs include severe pain during movement, sudden swelling or redness in the joints, shortness of breath, dizziness, or any other unusual symptoms that may indicate a serious underlying condition.

Test Group: Kurtoğlu Test

Test: big problem sir

HIGHEST
95.0

LOWEST
95.0

AVERAGE
95.0

Latest Result
95
2026-07-16

Test Group: Range of Motion (ROM)

Test: Knee Extension

Clinical AI Analysis

Doctor Standard/Target:
No doctor target was set

Clinical Interpretation:
The results of 179 and 156 degrees indicate varying degrees of knee extension capability, suggesting potential improvement or fluctuation in mobility.

Progress Evaluation:
Without a defined target, it is challenging to assess progress accurately. However, the values suggest that there may be some improvement in knee extension, which should be monitored closely.

Physiotherapy Focus:
Incorporate specific exercises aimed at enhancing knee extension range, focusing on flexibility and strength training for the quadriceps and hamstrings.

HIGHEST
179.0

LOWEST
156.0

AVERAGE
167.5

OVERALL PROGRESS

Decreased by 12.8% ↓ Since first test (2026-07-16)



Date	Result	Standard
2026-07-16	179	—
2026-07-16	156	—

Important Notice

This report is AI-assisted and intended for informational support only. It is not a medical diagnosis and does not replace professional clinical judgment. Please consult your doctor or therapist for any medical decisions.