



Physiotherapy & Rehabilitation Platform

Medical Report: Chronic Lumbar Disc Herniation with Sciatica

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #22

Clinical Details

Doctor: Dr. Alarkany

Date: 2026-04-28

Description: Patient presents with persistent radiating pain from the lower back down to the right calf. Diagnosed with L4-L5 disc protrusion. Pain is aggravated by prolonged sitting and forward bending.

Physical Measurements

Height: 178 cm

Weight: 85 kg

Blood Group: A+

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Moderate

Smart Clinical Analysis

Clinical Explanation:

The patient presents with persistent radiating pain from the lower back to the right calf, attributed to an L4-L5 disc protrusion. This condition is characterized by nerve root compression, leading to pain and functional limitations. The patient's activity level is moderate, and there are no chronic diseases reported, which is favorable for rehabilitation. The pain exacerbates with prolonged sitting and forward bending, indicating the need for careful management of these activities during the rehabilitation process.

Immediate Physiotherapy Recommendations:

A tailored physiotherapy program focusing on core stabilization, flexibility, and strengthening exercises for the lumbar and pelvic regions is recommended. Modalities such as heat therapy and electrical stimulation may be beneficial for pain management. Gradual progression of exercises should be emphasized to avoid exacerbation of symptoms.

Precautions:

The patient should avoid prolonged sitting and forward bending activities that may aggravate pain. Caution should be exercised during exercises that involve heavy lifting or twisting motions. Regular monitoring of symptoms during rehabilitation is essential.

Estimated Recovery Timeline:

Given the nature of the condition and the patient's response to therapy, a recovery timeline of approximately 8 to 12 weeks is anticipated, contingent upon adherence to the rehabilitation program and symptom management.

Risk Assessment:

Potential risks include worsening of symptoms, development of chronic pain, and possible neurological deficits if the disc protrusion continues to exert pressure on the nerve roots. Close monitoring is necessary to mitigate these risks.

Red Flags:

Emergency warning signs include sudden worsening of pain, loss of bowel or bladder control, significant weakness in the lower extremities, or any signs of cauda equina syndrome.

Immediate medical attention is required if these symptoms occur.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:
Normal knee flexion range for adults is typically 135 degrees.

Clinical Interpretation:
The recorded measurements indicate a progressive improvement in right knee flexion, moving from 95 degrees to 115 degrees over the assessment period. This suggests an increase in range of motion, which is a positive sign in the context of rehabilitation.

Progress Evaluation:
The trend shows a consistent improvement in knee flexion, indicating effective rehabilitation strategies and potential recovery of function in the lower extremity.

Physiotherapy Focus:
Continue to emphasize knee flexion exercises, along with stretching and strengthening of the surrounding musculature to support further improvement.


HIGHEST
115.0

LOWEST
95.0

AVERAGE
106.0

OVERALL PROGRESS

Increased by 21.1% ↑ Since first test (2026-04-14)



Date	Result
2026-04-14	95
2026-04-18	100
2026-04-22	110
2026-04-26	110
2026-04-28	115

Important Notice

This report is AI-assisted and intended for informational support only. It is not a medical diagnosis and does not replace professional clinical judgment. Please consult your doctor or therapist for any medical decisions.