



Physiotherapy & Rehabilitation Platform

Medical Report: Chronic Lumbar Disc Herniation with Sciatica

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #22

Clinical Details

Doctor: Dr. Alarkany

Date: 2026-04-28

Description: Patient presents with persistent radiating pain from the lower back down to the right calf. Diagnosed with L4-L5 disc protrusion. Pain is aggravated by prolonged sitting and forward bending.

Physical Measurements

Height: 178 cm

Weight: 85 kg

Blood Group: A+

Health History

Smoker: No

Chronic Diseases: No

Activity Level: Moderate

Smart Clinical Analysis

Clinical Explanation:

The patient presents with persistent radiating pain from the lower back to the right calf, attributed to an L4-L5 disc protrusion. This condition is characterized by nerve root compression, leading to pain exacerbated by prolonged sitting and forward bending. The patient's moderate activity level and absence of chronic diseases suggest a potential for effective rehabilitation. The focus will be on pain management, improving functional mobility, and restoring strength and flexibility in the lumbar region and lower extremities.

Immediate Physiotherapy Recommendations:

A tailored physiotherapy program should include:

- Core stabilization exercises to support the lumbar spine.
- Stretching exercises for the hamstrings and hip flexors to alleviate tension.
- Gradual introduction of low-impact aerobic activities to enhance overall fitness.
- Education on proper body mechanics and posture during daily activities.

Precautions:

- Avoid activities that involve heavy lifting or twisting motions.
- Limit prolonged sitting and forward bending to prevent exacerbation of symptoms.
- Monitor for any signs of increased pain or neurological deficits.

Estimated Recovery Timeline:

Given the nature of the condition and the patient's progress, a recovery timeline of 6 to 12 weeks is anticipated, contingent upon adherence to the rehabilitation program and response to treatment.

Risk Assessment:

Potential risks include worsening of symptoms, development of chronic pain, or progression of neurological deficits if the condition is not managed appropriately. Close monitoring of symptoms is essential.

Red Flags:

- Sudden onset of severe pain.
- Loss of bowel or bladder control.
- Progressive weakness in the lower extremities.
- Numbness or tingling in the genital or perianal region.

Test Group: Range of Motion (ROM) Assessment

Test: Knee Flexion (Right)

Clinical AI Analysis

Normal Reference:

Normal knee flexion range is typically between 135 to 150 degrees for adults.

Clinical Interpretation:

The patient's knee flexion measurements show a progressive increase from 95 degrees to 115 degrees over the assessment period, indicating improvement in range of motion and functional mobility.

Progress Evaluation:

The trend demonstrates a positive response to rehabilitation efforts, suggesting effective management of the underlying condition and improved joint function.

Physiotherapy Focus:

Continue to emphasize knee flexion exercises, incorporating both active and passive range of motion activities to further enhance flexibility and strength in the lower extremities.

HIGHEST

115.0

LOWEST

95.0


AVERAGE

106.0

OVERALL PROGRESS

Increased by 21.1% ↑

Since first test (2026-04-14)



Date	Result
2026-04-14	95
2026-04-18	100
2026-04-22	110
2026-04-26	110
2026-04-28	115