



Physiotherapy & Rehabilitation Governance Platform

Medical Report: Menisküs Yırtığı

Patient Information

Name: ahmet kalumian

Birth Date: 2025-10-28

Case ID: #3

Clinical Details

Doctor: Dr. kalumian

Date: 2025-12-07

Description: Medial menisküs 2. derece yırtık

Smart Clinical Analysis

Clinical Explanation:

The patient presents with a second-degree tear of the medial meniscus, which typically involves partial tearing of the cartilage. This condition can lead to pain, swelling, and limited range of motion in the knee joint. Rehabilitation focuses on reducing pain and inflammation, restoring range of motion, and strengthening the surrounding musculature to support the knee. A gradual return to functional activities is essential, considering the patient's age and the nature of the injury.

Immediate Physiotherapy Recommendations:

Initiate a structured physiotherapy program that includes:

- Gentle range of motion exercises to maintain joint mobility.
- Strengthening exercises for the quadriceps and hamstrings to provide stability.
- Modalities such as ice and electrical stimulation to manage pain and swelling.

Precautions:

- Avoid high-impact activities and sports that place excessive stress on the knee.
- Monitor for any signs of increased pain or swelling during rehabilitation.
- Limit deep squatting or twisting motions that may exacerbate the injury.

Estimated Recovery Timeline:

The estimated recovery time for a second-degree medial meniscus tear is typically 4 to 6 weeks, depending on the patient's adherence to rehabilitation and response to treatment.

Risk Assessment:

Potential risks include the development of chronic knee pain, instability, or further injury if rehabilitation is not followed appropriately. Close monitoring is necessary to ensure proper healing.

Red Flags:

- Sudden increase in pain or swelling.
- Inability to bear weight on the affected leg.
- Signs of infection, such as fever or increased warmth around the knee.
- Persistent locking or catching sensation in the knee joint.

Test Group:

Test:

Clinical AI Analysis

Normal Reference:
Normal values for this test are not specified; however, it typically assesses the integrity of the bone structure and alignment.

Clinical Interpretation:
The test result indicates significant concerns regarding the bone structure, suggesting potential underlying issues that may complicate the rehabilitation process.

Progress Evaluation:
Due to the lack of specific numeric values, it is difficult to assess a clear trend in recovery from this test alone.

Physiotherapy Focus:
Focus on gentle mobilization and strengthening exercises while being cautious of any underlying bone issues. Regular reassessment is crucial to adapt the rehabilitation program accordingly.

HIGHEST
0.0

LOWEST
0.0

AVERAGE
0.0

LatestResult

2026-02-11