

ECOLOGY LETTERS

A shift from phenol to silica-based leaf defenses during long-term soil and ecosystem development

Journal:	<i>Ecology Letters</i>
Manuscript ID	ELE-01360-2020.R1
Manuscript Type:	Letters
Date Submitted by the Author:	30-Jan-2021
Complete List of Authors:	<p>de Tombeur, Félix; University of Liege, Gembloux Agro-Bio Tech Laliberté, Etienne; Université de Montréal, Département de sciences biologiques Lambers, Hans; University of Western Australia, School of Plant Biology Faucon, Michel-Pierre; UniLaSalle Zemunik, Graham; University of Western Australia, School of Plant Biology Turner, Ben; Smithsonian Tropical Research Institute Cornélis, Jean-Thomas; University of Liege, Gembloux Agro-Bio Tech Mahy, Grégory; University of Liege Faculty of Gembloux Agro-Bio Tech, Biodiversity and Landscape, TERRA research centre</p>

SCHOLARONE™
Manuscripts

A shift from phenol to silica-based leaf defenses during long-term soil and ecosystem development

Felix de Tombeur^{1*}, Etienne Laliberté^{2,3}, Hans Lambers³, Michel-Pierre Faucon⁴, Graham Zemunik³, Benjamin L. Turner⁵, Jean-Thomas Cornelis¹, Grégory Mahy¹

¹TERRA Teaching and Research Centre, Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium.
²Institut de Recherche en Biologie Végétale, Université de Montréal, 4101 Sherbrooke Est, Montréal, QC H1X 2B2, Canada.
³School of Biological Sciences, The University of Western Australia, Crawley (Perth), WA 6009, Australia.
⁴AGHYLE, SFR Condorcet FR CNRS 3417, UniLaSalle, 19 rue Pierre Waguet, 60026, Beauvais, France.
⁵Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Panama.
^{*}Corresponding author ; 27 Avenue Maréchal Juin Bâtiment 52 – 5030 Gembloux (Belgium) ; +32488273660 ; felix.detombeur@uliege.be

Statement of authorship: All the authors conceived the ideas and designed methodology; FdT and GZ collected the samples; FdT analyzed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data accessibility statement: The data are available via the Dryad Digital Repository (add the link here)

Article type: Letters/Ideas & Perspectives

Running title: Leaf defenses during ecosystem development

Keywords: plant defense strategies ; silica-based defense ; plant silicon ; plant phenols ; resource availability hypothesis ; soil fertility gradient ; plant-herbivore interactions ; phenylalanine; resource limitation ; tradeoffs

Number of words in the Abstract (148), in the Main Text (5000)

Number of References: 99 (+ 3 references in the Supplementary Information)

Number of Figures and Tables: 6

Abstract

The resource availability hypothesis predicts that plants adapted to infertile soils have high levels of anti-herbivore leaf defenses. This hypothesis has been mostly explored for secondary metabolites such as phenolics, while it remains underexplored for silica-based defenses. We determined leaf concentrations of total phenols and silicon (Si) in plants growing along the 2-million-year Jurien Bay chronosequence, exhibiting an extreme gradient of soil fertility. We found that nitrogen (N) limitation on young soils led to a greater expression of phenol-based defenses, whereas old, phosphorus (P)-impoverished soils favored silica-based defenses. Both defense types were negatively correlated at the community and individual species level. Our results suggest a tradeoff among these two leaf defense strategies based on the strength and type of nutrient limitation, thereby opening up new perspectives for the resource availability hypothesis and plant defense research. This study also highlights the importance of silica-based defenses under low P supply.

37 Introduction

38 The resource availability hypothesis (RAH) (Coley *et al.* 1985) predicts that plant species adapted
39 to resource-rich environments will have rapid growth rates and leaf turnover, high leaf nutrient
40 concentrations, but low levels of anti-herbivore defenses. By contrast, the benefits of allocating
41 resources to anti-herbivore defenses become advantageous for species adapted to nutrient-poor
42 environments, because biomass loss by herbivory represents a significant loss of scarce nutrients
43 (Coley *et al.* 1985; Endara & Coley 2011). This theory has been primarily explored with regard to
44 secondary metabolites (e.g. phenolics), plant mechanical properties and nitrogen (N)-based
45 defenses, while silica-based defenses have received far less attention (Endara & Coley 2011). For
46 instance, in a meta-analysis testing the global consistency of the RAH (Endara & Coley 2011), the
47 deposits of silica bodies in plants as defense mechanism (McNaughton *et al.* 1985; Hartley &
48 DeGabriel 2016) represented only 1% of statistical tests considered in the literature (Massey *et al.*
49 2007). This highlights the fact that Si-based defenses have been understudied by plant ecologists,
50 despite representing one of the earliest anti-herbivore plant adaptations (Trembath-Reichert *et al.*
51 2015; Deshmukh *et al.* 2020).

52 Vascular plants can accumulate silicon (Si) in concentrations exceeding those of the major nutrients
53 (0.1 to 10% of dry weight; Epstein 1994). Taken up as monosilicic acid (H_4SiO_4) from the soil
54 solution, Si is translocated to sites of rapid transpiration, where it polymerizes as amorphous
55 hydrated silica ($\text{SiO}_2 \cdot n\text{H}_2\text{O}$) between cell walls and the lumen, and in extracellular and intercellular
56 spaces of the leaf epidermis (Kumar *et al.* 2017; de Tombeur *et al.* 2020a). This mechanism of
57 biosilicification has occurred in land plants for over 400 million years (Trembath-Reichert *et al.*
58 2015) and provides numerous benefits to plants, including resistance to abiotic stresses such as
59 metal toxicity, salinity, nutrient deficiency or water stress (Cooke & Leishman 2016; Debona *et al.*

1
2
3 60 2017; Frew *et al.* 2018). Biosilicification also reduces herbivory by increasing leaf abrasiveness,
4
5 61 which reduces penetration and chewing, and by decreasing the digestibility and palatability of
6
7 62 leaves (Massey & Hartley 2006, 2009; Massey *et al.* 2006, 2007; Johnson *et al.* 2020). Besides
8
9 63 physical defenses, biosilicification has been linked to anti-herbivore phytohormonal signaling
10
11 64 through the modulation of jasmonic acid, salicylic acid and ethylene, resulting in modified
12
13 65 emissions of volatile organic compounds (Coskun *et al.* 2019; Leroy *et al.* 2019). Although the
14
15 66 impact of silica-based defenses on herbivory has mainly been studied in Poales (e.g., grasses), it is
16
17 67 now well-established that the positive role of Si in biotic stresses is not restricted to Si-
18
19 68 accumulating families (Fauteux *et al.* 2006; Deshmukh *et al.* 2013; Katz 2014; Johnson *et al.* 2019;
20
21 69 Putra *et al.* 2020). Other taxa can have [Si] as high as in grasses (Hodson *et al.* 2005; Katz 2014),
22
23 70 Si transporters have been identified in some legume species (Deshmukh *et al.* 2013), and it has
24
25 71 recently been demonstrated that Si is an effective defense against herbivory in soybean (Fabaceae),
26
27 72 even at moderate leaf [Si] (Johnson *et al.* 2019). Despite these evidences, the expression of Si-
28
29 73 based defenses along soil fertility gradients remains underexplored, although we might expect them
30
31 74 to increase with declining soil fertility as predicted by the RAH.
32
33
34
35
36
37
38 75 Besides silica-based defenses, one of the most important classes of plant defenses are phenolics
39
40 76 compounds, which have been shown to co-vary with silica-based defenses (Cooke & Leishman
41
42 77 2012). In accordance with the RAH, high levels of phenolic compounds have long been considered
43
44 78 as having adaptive value for plants growing on infertile soils (Bryant *et al.* 1983; Coley *et al.* 1985;
45
46 79 Northup *et al.* 1995; Hättenschwiller *et al.* 2003; Kraus *et al.* 2004). It has been proposed that N
47
48 80 limitation leads to a greater expression of phenol-based defenses than phosphorus (P) limitation,
49
50 81 because N limitation impacts the phenylpropanoid pathway more strongly, which is involved in
51
52 82 the synthesis of both aromatic amino acids and phenol compounds (Haukioja *et al.* 1998; Jones &
53
54
55
56
57
58
59
60

Hartley 1999; Wright *et al.* 2010). While this hypothesis received support in some studies (Wright *et al.* 2010; de Long *et al.* 2016), others have instead found that P limitation significantly increases plant phenol synthesis (Hättenschwiler *et al.* 2003; Sampedro *et al.* 2011; Zhang *et al.* 2012). However, these studies generally considered one single species and/or performed fertilization to simulate nutrient limitations, while community-level studies in natural environments have not been conducted. Strong natural shifts from N to P limitation of plant productivity that occur during long-term soil and ecosystem development (Vitousek *et al.* 1993; Laliberté *et al.* 2012) might therefore impact plant phenol synthesis differently in species-rich plant communities, but this has not been explored. Furthermore, the potential covariation of phenolic foliar defenses with silica-based ones (Cooke & Leishman 2012) has never been studied along soil fertility gradients.

We determined leaf Si and total phenol concentrations in plants growing along the 2 million-year Jurien Bay soil chronosequence in south-western Australia (Turner & Laliberté 2015). This long-term soil chronosequence comprises a series of coastal dunes within a global biodiversity hotspot, supporting species-rich shrubland under a Mediterranean climate (Zemunik *et al.* 2016). It exhibits an extreme gradient of soil fertility in terms of rock-derived nutrients (P, calcium, magnesium, potassium), and shifts from N to P limitation of plant productivity as soils age (Laliberté *et al.* 2012; Hayes *et al.* 2014). Leaf concentrations of major plant nutrients decrease sharply with ecosystem age (Hayes *et al.* 2014; de Tombeur *et al.* 2020c), and plants converge towards highly efficient nutrient-use strategies on the oldest, most nutrient-impoverished soils (Hayes *et al.* 2014; Guilherme Pereira *et al.* 2019). The youngest soils can also be considered infertile from an N standpoint since their N capital has not yet built up and plant growth on these soils is limited by low N availability (Laliberté *et al.* 2012; Hayes *et al.* 2014). Long-term soil chronosequences that cover both the progressive and retrogressive phases of ecosystem development provide

exceptionally strong natural soil fertility and productivity gradients (Vitousek 2004; Peltzer *et al.* 2010) along which predictions of the RAH can be tested. Furthermore, the stoichiometric shifts from N to P limitation of plant growth that occur along such retrogressive soil chronosequences (Peltzer *et al.* 2010; Laliberté *et al.* 2012; Hayes *et al.* 2014) allow us to explore extensions of the RAH about the type of foliar defenses that are expressed depending on the type of nutrient limitation (e.g. N vs P on plant phenols: Wright *et al.* 2010; de Long *et al.* 2016), which have never been explored.

In accordance with the RAH, we hypothesized a greater expression of anti-herbivore defenses in plants growing on the oldest (P-limiting) and the very youngest (N-limiting) soils, compared with the intermediate-aged and most fertile soils where plant productivity, N and P availability peak (Laliberté *et al.* 2012, 2014), but further hypothesized that the type of defense most strongly expressed would depend on the type of nutrient limitation. Specifically, we hypothesized higher leaf phenol concentrations on the younger soils given the expected impact of N limitation on the phenylpropanoid pathway, whereas we expected silica-based defenses to be most strongly expressed in the oldest, P-impoverished soils. We expect the community-level patterns to be mostly driven by changes in plant species composition since the Jurien Bay chronosequence is characterized by a strong species turnover (Zemunik *et al.* 2016). This species turnover reflects the expression of selective edaphic forces acting on a species-rich regional flora over an ecological time scale (Laliberté *et al.* 2014).

Materials and methods

Study area and site description

127 The 2 Ma Jurien Bay chronosequence is located in south-western Australia, approximately 200 km
 128 north of Perth (Fig. S1), and is described in detail in Laliberté *et al.* (2012) and Turner & Laliberté
 129 (2015). The chronosequence, part of the Swan Coastal Plain, comprises a series of dunes parallel
 130 to the coast, formed by periodic interglacial sea-level high-stands since the Early Pleistocene/Late
 131 Pliocene (Kendrick *et al.* 1991), with a clear gradient of soil age with increasing distance from the
 132 Indian Ocean. The dunes comprise three units: the Quindalup dunes date from the Holocene (up to
 133 7 ka), the Spearwood dunes from the Middle Pleistocene (120 to 500 ka) and the Bassendean dunes
 134 from the Early Pleistocene or Late Pliocene (~2 Ma) (McArthur & Bettenay 1974; Playford *et al.*
 135 1976). The parent material of the dunes is calcareous sand from the nearshore coastal environment
 136 (Turner & Laliberté 2015). The climate is Mediterranean, with a mean annual temperature of 19°C,
 137 mean annual rainfall 533 mm and potential annual evapotranspiration of 1433 mm, which results
 138 in a water balance of -900 mm yr^{-1} (data from the Jurien Bay Bureau of Meteorology from 1968
 139 to 2015 in Turner *et al.* (2018).

140 We selected the same five chronosequence stages as in Hayes *et al.* (2014); these include both the
 141 early and retrogressive phases of long-term ecosystem development. The main soil properties of
 142 these five stages can be found in Table 1. Soil total P and carbonate concentrations, cation exchange
 143 capacity and pH-CaCl₂ continually decrease with increasing soil age. Soil total N concentrations
 144 increase from stage 1 to stage 2 during the progressive phase of ecosystem development, then
 145 decrease towards the last stages during the retrogressive phase (Laliberté *et al.* 2012; Turner &
 146 Laliberté 2015). Plant growth is most strongly limited by low N availability in the early stages, and
 147 by P availability in the advanced stages (Laliberté *et al.* 2012; Hayes *et al.* 2014). Previous studies
 148 showed that plant-available [Si] is low in the early stages of soil development, increases in stage 4

in the Spearwood dune system, and finally decreases in the oldest stage of soil development, where it is controlled by intense biocycling (Table 1) (de Tombeur *et al.* 2020b, c).

Site selection

For each chronosequence stage, we randomly selected five plots (10 m × 10 m each) among the 10 plots already characterized for soil and vegetation in previous studies (Hayes *et al.* 2014; Zemunik *et al.* 2016) (Fig. S1). The plots were originally selected using a random stratified sampling design (Zemunik *et al.* 2016). To characterize vegetation, seven 2 m × 2 m subplots were randomly positioned in each plot in which all individuals of all vascular plant species were counted (Zemunik *et al.* 2016). The percent canopy cover of each species was estimated, and the relative cover of each species was calculated as a fraction of the total canopy cover over the seven subplots (Zemunik *et al.* 2016).

Sampling procedure

In the 25 plots selected, we sampled leaves according to two procedures. First, we sampled leaves from one individual plant for each of the 10 most-abundant species of each plot as defined in Zemunik *et al.* (2016). The number of leaves sampled per individual was adapted according to their mass, but was never less than 10. Occasionally, a species originally included in the 10 most-abundant species was not found on the plot, which resulted in less than 10 species for some plots (Table S1). The 234 species sampled with this first procedure still accounted for 57% to 88% of the total cover of each plot (Table S1). The community-level analyses were performed only on these species. Second, we systematically sampled the species belonging to nine families, even if they were not included in the 10 most-abundant species, in order to study family-level variation in leaf [Si] and [phenols], following the same sampling procedure: Asparagaceae, Cyperaceae,

1
2
3 171 Ericaceae, Fabaceae, Haemodoraceae, Myrtaceae, Poaceae, Restionaceae and Rhamnaceae (Table
4
5 172 S2). These families were selected because they were well represented and found at all stages of the
6
7 173 chronosequence (Zemunik *et al.* 2016), and likely had contrasting [Si] based on known
8
9 174 phylogenetic patterns (Hodson *et al.* 2005). In total, 298 leaf samples belonging to 24 families were
10
11 175 collected (Tables S1 and S2).
12
13
14

15 176 All leaf material was collected over two weeks in November 2018. Leaves were sampled from one
16
17 177 healthy mature individual plant per species in each plot; when an individual did not provide
18
19 178 sufficient biomass for analysis (e.g., Poaceae spp.), leaf samples from several individuals within
20
21 179 the plot were combined.
22
23
24

25 180 *Leaf analyses*

26
27

28 181 Leaves were washed with distilled water, dried at 70 °C for 48 h and finely ground. Leaf material
29
30 182 (0.5 g) was placed in a porcelain crucible and calcinated at 450°C for 24 h. The weight after
31
32 183 calcination was used to calculate the ash content. The ash was mixed with 1.6 g lithium-metaborate
33
34 184 and 0.4 g of lithium-tetraborate in a graphite crucible and heated at 1000 °C for 5 min (Chao &
35
36 185 Sanzalone 1992). The bead was then dissolved in 15% HNO₃ and the concentrations of Si, P,
37
38 186 calcium (Ca), magnesium (Mg) and potassium (K) were determined by inductively coupled
39
40 187 plasma-optical emission spectrometry (Agilent Technologies, 700 series ICP-OES). Phenolic
41
42 188 compounds were extracted from a 0.25 g ground sample stirred with 10 mL of 70% acetone for 30
43
44 189 min (Salminen & Karonen 2011; Schaller *et al.* 2012; Bettaieb Rebey *et al.* 2020). Total phenols
45
46 190 were determined in triplicate as described in Salminen & Karonen (2011) using a Folin-Ciocalteu
47
48 191 assay with gallic acid monohydrate as standard (Merckx, Darmstadt, Germany). Total phenol
49
50 192 concentrations were expressed as g of gallic acid equivalents (GAE) per kilogram of dry weight.
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Soil sampling and analyses

In order to determine how Si availability in soils affected species-level variations in leaf [Si], three soil samples (top 20 cm) were taken in each of the 25 plots, for a total of 75 soil samples. Samples were air-dried and sieved (< 2 mm). The pool of ‘plant-available Si’ was determined by extraction in 0.01 M CaCl₂ (Haymsom & Chapman 1975; Sauer *et al.* 2006). Soil was shaken for 5 h in a 1:10 soil-to-solution ratio, filtered (cellulose filter, pore size < 2 µm, Healthcare Whatman™), acidified with 50 µL of ultrapure 65% HNO₃, and stored in darkness at 4°C prior to Si determination by ICP-OES.

Data analyses

To characterize leaf [phenols], [Si], [Ca], [Mg], [K] and [P] in plant communities across the chronosequence, we calculated the mean values of the 10 most-abundant species per plot, weighted or not by their relative canopy cover. The cover-weighted mean (CWM) was calculated as follows (Garnier *et al.* 2004; Violle *et al.* 2007):

$$CWM = \sum_{i=1}^S t_i \times RC_i$$

where t_i and RC_i are, respectively, the value of the trait t and its relative cover RC for a species i and S is the number of species.

The differences in plant-available [Si], leaf [phenols], [Si], [P], [Ca], [Mg] and [K] across the chronosequence stages were tested by one-way analysis of variance (ANOVA), followed by *post-hoc* multiple comparison (Fisher’s Least Significant Difference [LSD] tests). When these analyses considered all individuals together (i.e., not the mean and CWM of the plant communities), we treated species and plots as a random factor (mixed-effect models). We tested the relationships

between leaf [Si], [phenols] and major soil properties (total P, total N, ratio soil N:P) with linear mixed-effect models, treating plot and species as random factors when all individuals were considered together, and treating chronosequence stage as a random factor when the means and CWM of the 25 plots were considered. We also explored relationships between leaf [Si], [phenols] and foliar nutrient concentrations through Pearson tests of correlation. For the nine plant families selected, we tested the differences in leaf [phenols] and [Si] across the chronosequence stages using mixed-effect models with species and plot as random factors, followed by Fisher's LSD tests, and we tested the relationships between leaf [Si] and [phenols] with linear mixed-effect models (with plot and species as random factors). Additionally, we studied intraspecific variation for seven taxa, and detailed explanation of the statistical analyses used is presented in the Supplementary Information. Finally, a t-test was performed to examine the differences in leaf [Si] between dicots and monocots. All residuals were visually inspected for heteroscedasticity and appropriate transformations were performed to meet the model assumptions. All analyses were conducted in R using the 'nlme' (Pinheiro *et al.* 2020) and 'multcomp' (Hothorn *et al.* 2008) packages.

Results

Community-level leaf concentrations of Si, total phenols and major nutrients across the chronosequence stages

Leaf [Si] increased with increasing soil age, whether all individuals were considered together or the means and CWM of the 25 communities (Fig. 1a). The mean leaf [Si] within the 25 communities increased from stage 1 ($0.5 \pm 0.2 \text{ g kg}^{-1}$) to stages 2 and 3 (1.6 ± 0.4 and $2.4 \pm 0.4 \text{ g kg}^{-1}$), and then to stages 4 and 5 (3.3 ± 1.2 and $4.2 \pm 1.3 \text{ g kg}^{-1}$). The CWM followed the same pattern (Fig. 1a). The CWM leaf [Si] was negatively correlated to soil total P and positively correlated to soil N to P ratio (Fig. 2a), as for the means (Fig. S2).

1
2
3 237 By contrast, leaf [phenols] decreased with increasing chronosequence stage, whether all individuals
4
5 238 were considered together or the means and CWM of the 25 communities (Fig. 1b). However, only
6
7 239 stages 1-2 and 5 were significantly different from each other when all individuals were considered
8
9 240 together. The mean leaf [phenols] decreased from stage 1 (48.9 ± 5.5 GAE g kg⁻¹) to stage 5 (42.6
10
11 ± 3.9 GAE g kg⁻¹), as did the CWM (from 51.1 ± 5.1 to 42.9 ± 3.9 GAE g kg⁻¹). The CWM leaf
12
13 241 [phenols] was positively correlated to soil total P and negatively correlated to soil N to P ratio (Fig.
14
15 242 2b), as for the means (Fig. S2).
16
17
18
19 244 Leaf [phenols] and [Si] were correlated negatively with each other when all individuals were
20
21 245 considered (Fig. 3a), with the means and CWM of each community (Fig. 3b, c), and when species
22
23 246 means were considered (Fig. S3). Leaf [ash], [Ca], [Mg], [K] and [P] decreased with increasing
24
25 247 soil age (Fig. S4). Major leaf nutrient concentrations, including P, were therefore negatively
26
27 248 correlated with leaf [Si], whether all samples were considered individually or as means and CWM
28
29 249 of the 25 communities (Fig. 3d).
30
31
32
33
34 250 *Family and species-level leaf concentrations of Si and total phenols across the chronosequence*
35
36
37 251 Within the Cyperaceae, leaf [Si] significantly increased with increasing soil age while leaf
38
39 252 [phenols] decreased, and both defense types were negatively correlated through species
40
41 253 replacement across the chronosequence (Fig. 4). In other plant families, leaf [Si] increased or was
42
43 254 constant with increasing soil age, while the opposite was found for [phenols], but a significant
44
45 255 negative relationship between both defense types was identified only for the Restionaceae (Fig.
46
47 256 S5). For the seven species for which intra-specific variation was assessed, the mixed-effect models
48
49 257 showed that leaf [Si] significantly increased with increasing chronosequence stage while the
50
51 258 opposite was found for [phenols] (Table S3).
52
53
54
55
56
57
58
59
60

259 *Impact of species replacement, intra-specific and inter-specific variation on the community-level*
 260 *leaf [Si] and [phenols]*

261 Soil plant-available [Si] was lowest in stages 1 and 2 ($1.5 \pm 0.4 \text{ mg kg}^{-1}$), increased in stage 3 (4.0
 262 $\pm 0.3 \text{ mg kg}^{-1}$), then in stage 4 ($9.4 \pm 1.0 \text{ mg kg}^{-1}$), before it decreased at the oldest stage (4.3 ± 0.2
 263 mg kg^{-1}) (Table S4). Leaf [Si] was positively related to plant-available [Si] for three species of the
 264 seven (*Lepidosperma calcicola*, *Desmocladius asper* and *Acacia lasiocarpa* var. *lasiocarpa*) for
 265 which intra-specific variations were considered (Table S5). These species tended to accumulate
 266 more Si than those that did not show a correlation with plant-available [Si]. Despite this relationship
 267 between plant-available [Si] and leaf [Si] at the species-level (more pronounced for Si-
 268 accumulating species, with the exception of *Conostylis candicans* subsp. *calcicola*), the increase
 269 of the CWM leaf [Si] with increasing soil age was primarily driven by changes in the dominant
 270 plant families across the chronosequence (Fig. S6). The increase of the mean and CWM leaf [Si]
 271 with increasing soil age was not only driven by changes in the dominant families, but also within
 272 families and genera, which is described as Supplementary Results.

273 The plant families with the strongest contribution to the CWM leaf [phenols] strongly differed
 274 across the chronosequence (Fig. S6), highlighting the importance of species replacement on the
 275 patterns observed at the community level. In contrast to leaf [Si], the contribution of each family
 276 to the CWM [phenols] was proportional to its relative cover, highlighting less variation in [phenols]
 277 among families.

278 **Discussion**

279 Overall, our results do not support the main prediction of the RAH, since investments in the two
 280 anti-herbivore defenses considered were not the lowest at the most fertile soils, where both N and

1
2
3 281 P availability and plant productivity peak (stages 2 and 3) (Laliberté *et al.* 2012). Instead,
4
5 282 community-level leaf [Si] was highest on old, nutrient-impoverished soils, where P limits plant
6
7 283 productivity, while leaf [phenols] were the highest on young soils, where plant productivity is
8
9
10 284 limited by N, and both defenses were negatively correlated with each other (Fig. 5). Nevertheless,
11
12 285 only two types of chemical defense were considered in this study and the hypothesis of an overall
13
14 286 greater investment in defenses in the least fertile soils as predicted by the RAH should consider the
15
16
17 287 full array of anti-herbivore defenses, including structural, physical and qualitative chemical
18
19 288 defenses (Aplin & Cannon 1971; Moles *et al.* 2013; Lambers & Oliveira 2019).
20
21
22 289 The use of soil chronosequences allows us to minimize effects of other ecosystem properties (e.g.,
23
24 290 climate, topography, parent material, salt content) beyond soil age and associated changes in major
25
26 291 nutrients (e.g. N, P) that might influence leaf [Si] and [phenols], yet we cannot rule completely out
27
28 292 other covarying factors. In particular, soil pH declines along the chronosequence which could
29
30 293 indirectly impact plant-available [Si] (de Tombeur *et al.* 2020b). However, our previous work has
31
32
33 294 shown that community-level changes in foliar [Si] with soil age across this chronosequence
34
35 295 primarily arise from plant species turnover toward those that accumulate foliar [Si] as soil P
36
37 296 availability declines, and not directly because of pH-induced changes in Si availability (de
38
39
40 297 Tombeur *et al.* 2020c). Similarly, since soil nutrient availability has a greater impact on plant
41
42 298 phenol concentrations than soil pH (Kraus *et al.* 2004), we interpret the changes in foliar [Si] and
43
44 299 [phenols] mainly in terms of changes in soil N and P availability that are the major drivers of plant
45
46
47 300 community assembly along the Jurien Bay chronosequence (Laliberté *et al.* 2014).
48
49
50 301 *Convergence towards silica-based defenses during ecosystem retrogression*
51
52
53 302 Our results show a convergence towards plants having higher leaf [Si] on P-impoverished soils,
54
55 303 during ecosystem retrogression. Indeed, despite the decrease of plant-available [Si] in the last stage,
56
57
58
59
60

induced by soil desilication (de Tombeur *et al.* 2020b), the intense cycling of highly-soluble plant-derived Si in the oldest soils allows maintenance of high leaf [Si] (de Tombeur *et al.* 2020c). The convergence towards plants having higher leaf [Si] with increasing nutrient depletion was accompanied by a strong decrease in leaf macronutrient concentrations, including N (Hayes *et al.* 2014; Guilherme Pereira *et al.* 2019), and an increase in leaf mass per area (LMA) and leaf dry matter content (LMDC) along the Jurien Bay chronosequence (Guilherme Pereira *et al.* 2019). These traits highlight the convergence towards slow-growing plants adapted to P-impooverished soils and nutrient-poor environments with increasing soil age (Lambers & Poorter 1992; Wright *et al.* 2004; Garnier & Navas 2013; Reich 2014). Their higher leaf [Si] suggests that these species have evolved towards higher levels of silica deposits to minimize biomass and nutrient loss by herbivores in these nutrient-poor environments, in line with the RAH (Coley *et al.* 1985). Following this hypothesis, the likely longer leaf lifespan of these species could partly explain their higher expression of silica-based defenses, since Si accumulates as a leaf ages (Motomura *et al.* 2002).

An alternative but not necessarily mutually exclusive hypothesis could be mitigation of P stress. The first evidence for a positive effect of Si for plants growing under P stress came from the Rothamsted Experimental Station (Hall & Morison 1906). Since then, other studies have made similar observations (Ma & Takahashi 1990; Neu *et al.* 2017) and Quigley *et al.* (2020) recently proposed that high levels of biosilicification might be an adaptation to resource-poor environments. Although the mechanisms proposed are numerous and diverse, the positive role of Si under P stress is becoming clearer (Kostic *et al.* 2017), and could explain the convergence towards species having higher leaf [Si] with increasing P depletion. Overall, our results suggest that leaf Si is associated with the ‘slow’ end of the leaf economics spectrum (Wright *et al.* 2004; Reich 2014), with species

1
2
3 327 having long-lived leaves with low macronutrient concentrations and high LMA and LDMC, but
4
5 328 this hypothesis requires further investigation.
6
7
8 329 The increase of the CWM leaf [Si] from the P-limited ecosystem in stage 4 is partly explained by
9
10 330 Proteaceae having high relative cover and relatively high [Si] for dicot species (up to 6.8 g kg⁻¹
11
12 331 DW; Hodson *et al.* 2005). We have no information on Si transporters (Ma *et al.* 2006) in
13
14 332 Proteaceae, but it is possible that the exudation of carboxylates in the rhizosphere by their cluster
15
16 333 roots (Lambers *et al.* 2008) mobilizes Si from poorly-soluble minerals, thus favoring its uptake.
17
18 334 These cluster roots release carboxylates into the rhizosphere and release P from strongly sorbed
19
20 335 inorganic forms (Lambers *et al.* 2008), but also mobilize micronutrients such as manganese (Mn)
21
22 336 (Lambers *et al.* 2015; Pang *et al.* 2018). Some carboxylate-releasing roots mobilize P from rocks
23
24 337 that are highly resistant to weathering, such as quartzite (Teodoro *et al.* 2019). The relatively high
25
26 338 leaf [Si] in some dicot species exhibiting such specialized P-acquisition strategies might therefore
27
28 339 be explained by their mobilization of Si through carboxylate exudation into the rhizosphere. More
29
30 340 generally, this mechanism might explain the convergence towards leaves with higher [Si] with
31
32 341 increasing nutrient-depletion, because the relative cover of cluster-rooted species increases
33
34 342 markedly along the Jurien Bay chronosequence (Fig. 5) (Zemunik *et al.* 2015), and other species
35
36 343 might benefit from the carboxylate exudation of their neighbors (Lambers *et al.* 2018). This
37
38 344 mechanism may also explain why we observed species with some of the highest leaf [Si] found in
39
40 345 literature (up to 43 g kg⁻¹ DW for *Mesomelaena pseudostygia* at stage 4) (Schoelynck *et al.* 2010;
41
42 346 Carey *et al.* 2017, 2019; Schaller *et al.* 2018; Ishizawa *et al.* 2019; Nakamura *et al.* 2019) although
43
44 347 the Jurien Bay soils are among the most desilicated worldwide, with one of the lowest Si
45
46 348 availability for plants (de Tombeur *et al.* 2020b).
47
48
49
50
51
52
53
54
55 349 *Nitrogen limitation and phenol synthesis on young soils*
56
57
58
59
60

1
2
3 350 The CWM leaf [phenols] slightly, but significantly, decreased with increasing soil age, which we
4
5 351 interpreted as resulting from the N-limiting conditions on the youngest soils. Indeed, although some
6
7 352 studies suggest that P limitation increases phenol synthesis (Sampedro *et al.* 2011; Zhang *et al.*
8
9 353 2012) including along a long-term Hawaiian soil chronosequence (Hättenschwiler *et al.* 2003),
10
11 354 our results are in line with others showing that N limitation has a greater effect on quantitative plant
12
13 355 secondary metabolites (Koricheva *et al.* 1998; Wright *et al.* 2010; de Long *et al.* 2016).
14
15
16

17 356 The apparent stronger effect of N- compared with P limitation might be explained by their different
18
19 357 role in plant metabolism (Jones & Hartley 1999; Wright *et al.* 2010). Under N-limiting conditions,
20
21 358 there could be a surplus of photo-assimilates because leaf growth decreases sharply, while
22
23 359 photosynthesis continues, albeit at a somewhat slower rate (Prescott *et al.* 2020). The ammonium-
24
25 360 N in phenylalanine, which is a precursor of both aromatic amino acids and phenolic compounds,
26
27 361 would then be released and re-used in other N-containing molecules (Kováčik *et al.* 2007). The
28
29 362 rest of the phenylalanine molecule then becomes available for secondary metabolic pathways, in
30
31 363 which it is converted into C-based secondary metabolites such as phenylpropanoid derivatives
32
33 364 (Haukioja *et al.* 1998; Koricheva *et al.* 1998; Jones & Hartley 1999). This might explain why plants
34
35 365 growing on the young N limited environments had higher [phenols], eliminating a surplus of photo-
36
37 366 assimilates while conserving N. Under P-limiting conditions, however, any surplus of photo-
38
39 367 assimilates would be exported from the chloroplasts as P-containing compounds, and converted
40
41 368 into sucrose via reactions that release P, which is retained and reused (Stitt & Quick 1989).
42
43 369 Therefore, P-limiting conditions might not affect the pathway of secondary metabolite synthesis in
44
45 370 a way that N limitation does (Koricheva *et al.* 1998; Wright *et al.* 2010). Although the primary
46
47 371 function of plant phenolic compounds may be the release of N, rather than defense against
48
49 372 herbivores, it does make leaves less palatable for mammalian herbivores such as kangaroos
50
51
52
53
54
55
56
57
58
59
60

(Rafferty *et al.* 2005, 2010) on the youngest N-limited stages of the Jurien Bay chronosequence (Fig. 5).

Conclusions and Perspectives

Leaf [phenols] and [Si] were negatively correlated when considering the means and CWM of each community, all the individuals together, and within the Cyperaceae and Restionaceae family, which suggest a tradeoff between both leaf defense strategies. This pattern was observed previously (Cooke & Leishman 2012; Moles *et al.* 2013; Frew *et al.* 2016; Simpson *et al.* 2017; Waterman *et al.* 2020), but never along a soil resource gradient. Here, the species growing on older soils and adapted to nutrient-poor environments tend to favor silica accumulation over the synthesis of phenols compared with species growing on younger soils.

Raven (1983) calculated that, on a weight basis, the energetic cost of incorporating 1 g of lignin is about 27 times higher than that of incorporating 1 g of SiO₂. Given the lower metabolic costs of incorporating Si compared with other C-based compounds having similar functions (cellulose, lignin, phenols) (Schoelynck *et al.* 2010; Cooke & Leishman 2012; Klotzbücher *et al.* 2018; de Tombreur *et al.* 2020a), investing in silica as a defense mechanism (and eventually as leaf support for Cyperaceae) would make sense from an energetic standpoint on the oldest and most nutrient-depleted soils, where plants converge towards the ‘slow’ end of the leaf economics spectrum (Reich 2014; Guilherme Pereira *et al.* 2019). This energetic gain would save resources for other key aspects of the plant life cycle like growth and reproduction during ecosystem retrogression, which is key in these highly-infertile environments (Lambers 2014). Further studies should now investigate other types of anti-herbivore defenses to better evaluate the overall investment in defenses as a function of soil fertility along the Jurien Bay chronosequence. This is important given that plants display a wide range of defense traits, without clear evidence of tradeoffs between them

globally (Moles *et al.* 2013). Overall, our study suggests that the type of nutrient limitation (i.e., N vs P) may induce tradeoffs in plant defense strategies, opening up new perspectives about the role of soil nutrient stoichiometry in the expression of plant defenses.

Acknowledgements

We thank the Western Australian Department of Biodiversity, Conservation and Attractions for permission to sample along the Jurien Bay chronosequences and for access to these outstanding ecosystems. This work would not have been possible without the invaluable help of Jean-Charles Bergen, Nea Fatimata and Evelyne Tanoh whom we sincerely thank. J-T.C and F.dT were supported by ‘Fonds National de la Recherche Scientifique’ of Belgium (FNRS; Research Credit Grant for the project SiCliNG CDR J.0117.18).

References

- Aplin, T.E.. & Cannon, J.R. (1971). Distribution of alkaloids in some Western Australian plants. *Econ. Bot.*, 25, 366–380.
- Bettaieb Rebey, I., Bourgou, S., Ben Kaab, S., Aidi Wannes, W., Ksouri, R., Saidani Tounsi, M., *et al.* (2020). On the effect of initial drying techniques on essential oil composition, phenolic compound and antioxidant properties of anise (*Pimpinella anisum* L.) seeds. *J. Food Meas. Charact.*, 14, 220–228.
- Bryant, J.P., Chapin, F.S. & Klein, D.R. (1983). Carbon/nutrient balance of boreal plants in relation to vertebrate herbivory. *Oikos*, 40, 357–368.
- Carey, J.C., Abbott, B.W. & Rocha, A. V. (2019). Plant Uptake Offsets Silica Release From a Large Arctic Tundra Wildfire. *Earth's Futur.*, 7, 1044–1057.
- Carey, J.C., Parker, T.C., Fetcher, N. & Tang, J. (2017). Biogenic silica accumulation varies

- across tussock tundra plant functional type. *Funct. Ecol.*, 1–11.
- Chao, T.T. & Sanzalone, R.F. (1992). Decomposition techniques. *J. Geochemical Explor.*, 44, 65–106.
- Coley, P.D., Bryant, J.P. & Chapin, F.S. (1985). Resource availability and plant antiherbivore defense. *Science* (80-.), 230, 895–899.
- Cooke, J. & Leishman, M.R. (2012). Tradeoffs between foliar silicon and carbon-based defences: Evidence from vegetation communities of contrasting soil types. *Oikos*, 121, 2052–2060.
- Cooke, J. & Leishman, M.R. (2016). Consistent alleviation of abiotic stress with silicon addition: a meta-analysis. *Funct. Ecol.*, 30, 1340–1357.
- Coskun, D., Deshmukh, R., Sonah, H., Menzies, J.G., Reynolds, O., Ma, J.F., *et al.* (2019). The controversies of silicon's role in plant biology. *New Phytol.*, 221, 67–85.
- Debona, D., Rodrigues, F.A. & Datnoff, L.E. (2017). Silicon's Role in Abiotic and Biotic Plant Stresses. *Annu. Rev. Phytopathol.*, 55, 85–107.
- Deshmukh, R., Sonah, H. & Belanger, R. (2020). New evidence defining the evolutionary path of aquaporins regulating silicon uptake in land plants. *J. Exp. Bot.*
- Deshmukh, R.K., Vivancos, J., Guérin, V., Sonah, H., Labbé, C., Belzile, F., *et al.* (2013). Identification and functional characterization of silicon transporters in soybean using comparative genomics of major intrinsic proteins in Arabidopsis and rice. *Plant Mol. Biol.*, 83, 303–315.
- Endara, M.J. & Coley, P.D. (2011). The resource availability hypothesis revisited: A meta-analysis. *Funct. Ecol.*, 25, 389–398.

- 439 Epstein, E. (1994). The anomaly of silicon in plant biology. *Proc. Natl. Acad. Sci. U. S. A.*, 91,
440 11–17.
- 441 Fauteux, F., Chain, F., Belzile, F., Menzies, J.G. & Bélanger, R.R. (2006). The protective role of
442 Si in the Arabidopsis-powdery mildew pathosystem. *Proc. Natl. Acad. Sci.*, 103, 17554–
443 17559.
- 444 Frew, A., Powell, J.R., Sallam, N., Allsopp, P.G. & Johnson, S.N. (2016). Trade-offs between
445 silicon and phenolic defenses may explain enhanced performance of root herbivores on
446 phenolic-rich plants. *J. Chem. Ecol.*, 42, 768–771.
- 447 Frew, A., Weston, L.A., Reynolds, O.L. & Gurr, G.M. (2018). The role of silicon in plant
448 biology: a paradigm shift in research approach. *Ann. Bot.*, 121, 1265–1273.
- 449 Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., *et al.* (2004). Plant
450 functional markers capture ecosystem properties during secondary succession. *Ecology*, 85,
451 2630–2637.
- 452 Garnier, E. & Navas, M.-L. (2013). Caractérisation fonctionnelle des végétaux. In: *Diversité*
453 *fonctionnelle des plantes* (ed. Boeck, D.). p. 353.
- 454 Guilherme Pereira, C., Hayes, P.E., O’Sullivan, O.S., Weerasinghe, L.K., Clode, P.L., Atkin,
455 O.K., *et al.* (2019). Trait convergence in photosynthetic nutrient-use efficiency along a 2-
456 million year dune chronosequence in a global biodiversity hotspot. *J. Ecol.*, 107, 2006–
457 2023.
- 458 Hall, A.D. & Morison, C.G.T. (1906). On the function of silica in the nutrition of cereals.—Part
459 I. *Proc. R. Soc. London. Ser. B*, 77, 455–477.

- 460 Hartley, S.E. & DeGabriel, J.L. (2016). The ecology of herbivore-induced silicon defences in
461 grasses. *Funct. Ecol.*, 30, 1311–1322.
- 462 Hättenschwiler, S., Hagerman, A.E. & Vitousek, P.M. (2003). Polyphenols in litter from tropical
463 montane forests across a wide range in soil fertility. *Biogeochemistry*, 64, 129–148.
- 464 Haukioja, E., Ossipov, V., Koricheva, J., Honkanen, T., Larsson, S. & Lempa, K. (1998).
465 Biosynthetic origin of carbon-based secondary compounds: Cause of variable responses of
466 woody plants to fertilization? *Chemoecology*, 8, 133–139.
- 467 Hayes, P., Turner, B.L., Lambers, H. & Laliberté, E. (2014). Foliar nutrient concentrations and
468 resorption efficiency in plants of contrasting nutrient-acquisition strategies along a 2-
469 million-year dune chronosequence. *J. Ecol.*, 102, 396–410.
- 470 Haymsom, M. & Chapman, L. (1975). Some aspects of the calcium silicate trials at Mackay.
471 *Proc. Queensl. Soc. Sugar Cane Technol.*, 42, 117–122.
- 472 Hodson, M.J., White, P.J., Mead, A. & Broadley, M.R. (2005). Phylogenetic variation in the
473 silicon composition of plants. *Ann. Bot.*, 96, 1027–1046.
- 474 Hothorn, T., Bretz, F. & Westfall, P. (2008). Simultaneous Inference in General Parametric
475 Models. *Biometrical J.*, 50, 346–363.
- 476 Ishizawa, H., Niiyama, K., Iida, Y., Shari, N.H.Z., Ripin, A. & Kitajima, K. (2019). Spatial
477 variations of soil silicon availability and biogenic silicon flux in a lowland tropical forest in
478 Malaysia. *Ecol. Res.*, 34, 548–559.
- 479 Johnson, S.N., Hartley, S.E., Ryalls, J.M.W., Frew, A. & Hall, C.R. (2020). Targeted plant
480 defense: silicon conserves hormonal defense signaling impacting chewing but not fluid-

- 481 feeding herbivores. *Ecology*, In press.
- 482 Johnson, S.N., Rowe, R.C. & Hall, C.R. (2019). Silicon is an inducible and effective herbivore
483 defence against *Helicoverpa punctigera* (Lepidoptera: Noctuidae) in soybean. *Bull.*
484 *Entomol. Res.*, 1–6.
- 485 Jones, C.G. & Hartley, S.E. (1999). A protein competition model of phenolic allocation. *Oikos*,
486 86, 27–44.
- 487 Katz, O. (2014). Beyond grasses: the potential benefits of studying silicon accumulation in non-
488 grass species. *Front. Plant Sci.*, 376, 1–3.
- 489 Kendrick, G.W., Wyrwoll, K.-H. & Szabo, B.J. (1991). Pliocene-Pleistocene coastal events and
490 history along the western margin of Australia. *Quat. Sci. Rev.*, 10, 419–439.
- 491 Klotzbücher, T., Klotzbücher, A., Kaiser, K., Vetterlein, D., Jahn, R. & Mikutta, R. (2018).
492 Variable silicon accumulation in plants affects terrestrial carbon cycling by controlling
493 lignin synthesis. *Glob. Chang. Biol.*, 24, 183–189.
- 494 Koricheva, J., Larsson, S., Haukioja, E. & Keinänen, M. (1998). Regulation of woody plant
495 secondary metabolism by resource availability: hypothesis testing by means of meta-
496 analysis. *Oikos*, 83, 212–226.
- 497 Kostic, L., Nikolic, N., Bosnic, D., Samardzic, J. & Nikolic, M. (2017). Silicon increases
498 phosphorus (P) uptake by wheat under low P acid soil conditions. *Plant Soil*, 419, 447–455.
- 499 Kováčik, J., Klejdus, B., Bačkor, M. & Repčák, M. (2007). Phenylalanine ammonia-lyase
500 activity and phenolic compounds accumulation in nitrogen-deficient *Matricaria chamomilla*
501 leaf rosettes. *Plant Sci.*, 172, 393–399.

- 502 Kraus, T.E.C., Zasoski, R.J. & Dahlgren, R.A. (2004). Fertility and pH effects on polyphenol and
503 condensed tannin concentrations in foliage and roots. *Plant Soil*, 262, 95–109.
- 504 Kumar, S., Soukup, M. & Elbaum, R. (2017). Silicification in Grasses: Variation between
505 Different Cell Types. *Front. Plant Sci.*, 8.
- 506 Laliberté, E., Turner, B.L., Costes, T., Pearse, S.J., Wyrwoll, K.H., Zemunik, G., *et al.* (2012).
507 Experimental assessment of nutrient limitation along a 2-million-year dune chronosequence
508 in the south-western Australia biodiversity hotspot. *J. Ecol.*, 100, 631–642.
- 509 Laliberté, E., Zemunik, G. & Turner, B.L. (2014). Environmental filtering explains variation in
510 plant diversity along resource gradients. *Science* (80-.), 345, 1602–1605.
- 511 Lambers, H. (2014). *Plant Life on the Sandplains in Southwest Australia, a Global Biodiversity*
512 *Hotspot*. 2004th edn. Univerty of Western Australia Publishing, Crawley, Australia.
- 513 Lambers, H., Albornoz, F., Kotula, L., Laliberté, E., Ranathunge, K., Teste, F.P., *et al.* (2018).
514 How belowground interactions contribute to the coexistence of mycorrhizal and non-
515 mycorrhizal species in severely phosphorus-impooverished hyperdiverse ecosystems. *Plant*
516 *Soil*, 424, 11–33.
- 517 Lambers, H., Hayes, P.E., Laliberté, E., Oliveira, R.S. & Turner, B.L. (2015). Leaf manganese
518 accumulation and phosphorus-acquisition efficiency. *Trends Plant Sci.*, 20, 83–90.
- 519 Lambers, H. & Oliveira, R.S. (2019). *Plant Physiological Ecology, 3rd edition*. Springer
520 International Publishing.
- 521 Lambers, H. & Poorter, H. (1992). Inherent variation in growth rate between higher plants: a
522 search for physiological causes and ecological consequences. *Adv. Ecol. Res.*, 23, 187–261.

- 523 Lambers, H., Raven, J.A., Shaver, G.R. & Smith, S.E. (2008). Plant nutrient-acquisition
524 strategies change with soil age. *Trends Ecol. Evol.*, 23, 95–103.
- 525 Leroy, N., de Tombeur, F., Walgraffe, Y., Cornelis, J.-T. & Verheggen, F.J. (2019). Silicon and
526 Plant Natural Defenses against Insect Pests : Impact on Plant Volatile Organic Compounds
527 and Cascade Effects on Multitrophic Interactions. *Plants*, 8, 444.
- 528 de Long, J.R., Sundqvist, M.K., Gundale, M.J., Giesler, R. & Wardle, D.A. (2016). Effects of
529 elevation and nitrogen and phosphorus fertilization on plant defence compounds in subarctic
530 tundra heath vegetation. *Funct. Ecol.*, 30, 314–325.
- 531 Ma, J.F. & Takahashi, E. (1990). Effect of silicon on the growth and phosphorus uptake of rice.
532 *Plant Soil*, 126, 115–119.
- 533 Ma, J.F., Tamai, K., Yamaji, N., Mitani, N., Konishi, S., Katsuhara, M., *et al.* (2006). A silicon
534 transporter in rice. *Nature*, 440, 688–691.
- 535 Massey, F. & Hartley, S. (2009). Physical defences wear you down : progressive and irreversible
536 impacts of silica on insect herbivores. *J. Anim. Ecol.*, 78, 281–291.
- 537 Massey, F.P., Ennos, A.R. & Hartley, S.E. (2006). Silica in grasses as a defence against insect
538 herbivores: Contrasting effects on folivores and a phloem feeder. *J. Anim. Ecol.*, 75, 595–
539 603.
- 540 Massey, F.P., Ennos, A.R. & Hartley, S.E. (2007). Grasses and the resource availability
541 hypothesis: The importance of silica-based defences. *J. Ecol.*, 95, 414–424.
- 542 Massey, F.P. & Hartley, S.E. (2006). Experimental demonstration of the antiherbivore effects of
543 silica in grasses: impacts on foliage digestibility and vole growth rates. *Proc. R. Soc. B Biol.*

- 544 *Sci.*, 273, 2299–2304.
- 545 McArthur, W.M. & Bettenay, E. (1974). *Development and distribution of soils of the Swan*
- 546 *Coastal Plain, Western Australia. Canberra: CSIRO. CSIRO. Melbourne.*
- 547 McNaughton, S.J., Tarrants, J.L., McNaughton, M.M. & Davis, R.D. (1985). Silica as a defense
- 548 against herbivory and a growth promotor in african grasses. *Ecology*, 66, 528–535.
- 549 Moles, A.T., Peco, B., Wallis, I.R., Foley, W.J., Poore, A.G.B., Seabloom, E.W., *et al.* (2013).
- 550 Correlations between physical and chemical defences in plants: Tradeoffs, syndromes, or
- 551 just many different ways to skin a herbivorous cat? *New Phytol.*, 198, 252–263.
- 552 Motomura, H., Mita, N. & Suzuki, M. (2002). Silica accumulation in long-lived leaves of *Sasa*
- 553 *veitchii* (Carrière) rehder (Poaceae-Bambusoideae). *Ann. Bot.*, 90, 149–152.
- 554 Nakamura, R., Ishizawa, H., Wagai, R., Suzuki, S., Kitayama, K. & Kitajima, K. (2019). Silicon
- 555 cycled by tropical forest trees: effects of species, elevation and parent material on Mount
- 556 Kinabalu, Malaysia. *Plant Soil*, 443, 155–166.
- 557 Neu, S., Schaller, J. & Dudel, E.G. (2017). Silicon availability modifies nutrient use efficiency
- 558 and content, C:N:P stoichiometry, and productivity of winter wheat (*Triticum aestivum* L.).
- 559 *Sci. Rep.*, 7, 40829.
- 560 Northup, R.R., Yu, Z., Dahlgren, R.A. & Vogt, A. (1995). Polyphenol control of nitrogen release
- 561 from pine litter. *Nature*, 377, 227–229.
- 562 Pang, J., Bansal, R., Zhao, H., Bohuon, E., Lambers, H., Ryan, M.H., *et al.* (2018). The
- 563 carboxylate-releasing phosphorus-mobilizing strategy can be proxied by foliar manganese
- 564 concentration in a large set of chickpea germplasm under low phosphorus supply. *New*

- 565 *Phytol.*, 219, 518–529.
- 566 Peltzer, D.A., Wardle, D.A., Allison, V.J., Baisden, W.T., Bardgett, D., Chadwick, O.A., *et al.*
567 (2010). Understanding ecosystem retrogression. *Ecol. Monogr.*, 80, 509–529.
- 568 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & Team, R.C. (2020). nlme: Linear and Nonlinear
569 Mixed Effects Models.
- 570 Playford, P.E., Cockbain, A.E. & Lowe, G.H. (1976). *Geology of the Perth Basin, Western*
571 *Australia*. Perth, W.A.
- 572 Prescott, C.E., Grayston, S.J., Helmisaari, H., Ka, E., Körner, C., Lambers, H., *et al.* (2020).
573 Surplus Carbon Drives Allocation and Plant – Soil Interactions, xx, 1–9.
- 574 Putra, R., Powell, J.R., Hartley, S.E. & Johnson, S.N. (2020). Is it time to include legumes in
575 plant silicon research? *Funct. Ecol.*, 34, 1142–1157.
- 576 Quigley, K.M., Griffith, D.M., Donati, G.L. & Anderson, T.M. (2020). Soil nutrients and
577 precipitation are major drivers of global patterns of grass leaf silicification. *Ecology*, 101, 1–
578 10.
- 579 Rafferty, C., Lamont, B.B. & Hanley, M.E. (2005). Selective feeding by kangaroos (*Macropus*
580 *fuliginosus*) on seedlings of *Hakea* species: Effects of chemical and physical defences. *Plant*
581 *Ecol.*, 177, 201–208.
- 582 Rafferty, C.M., Lamont, B.B. & Hanley, M.E. (2010). Herbivore feeding preferences in captive
583 and wild populations. *Austral Ecol.*, 35, 257–263.
- 584 Raven, J.A. (1983). The transport and function of silicon in plants. *Biol. Rev.*, 58, 179–207.
- 585 Reich, P.B. (2014). The world-wide “fast-slow” plant economics spectrum: A traits manifesto. *J.*

- 586 *Ecol.*, 102, 275–301.
- 587 Salminen, J.P. & Karonen, M. (2011). Chemical ecology of tannins and other phenolics: We need
588 a change in approach. *Funct. Ecol.*, 25, 325–338.
- 589 Sampedro, L., Moreira, X. & Zas, R. (2011). Costs of constitutive and herbivore-induced
590 chemical defences in pine trees emerge only under low nutrient availability. *J. Ecol.*, 99,
591 818–827.
- 592 Sauer, D., Saccone, L., Conley, D.J., Herrmann, L. & Sommer, M. (2006). Review of
593 methodologies for extracting plant-available and amorphous Si from soils and aquatic
594 sediments. *Biogeochemistry*, 80, 89–108.
- 595 Schaller, J., Brackhage, C. & Dudel, E.G. (2012). Silicon availability changes structural carbon
596 ratio and phenol content of grasses. *Environ. Exp. Bot.*, 77, 283–287.
- 597 Schaller, J., Turner, B.L., Weissflog, A., Pino, D., Bielnicka, A.W. & Engelbrecht, B.M.J.
598 (2018). Silicon in tropical forests: large variation across soils and leaves suggests ecological
599 significance. *Biogeochemistry*, 140, 161–174.
- 600 Schoelynck, J., Bal, K., Backx, H., Okruszko, T., Meire, P. & Struyf, E. (2010). Silica uptake in
601 aquatic and wetland macrophytes: A strategic choice between silica, lignin and cellulose?
602 *New Phytol.*, 186, 385–391.
- 603 Simpson, K.J., Wade, R.N., Rees, M., Osborne, C.P. & Hartley, S.E. (2017). Still armed after
604 domestication? Impacts of domestication and agronomic selection on silicon defences in
605 cereals. *Funct. Ecol.*, 31, 2108–2117.
- 606 Stitt, M. & Quick, W.P. (1989). Photosynthetic carbon partitioning: its regulation and

- possibilities for manipulation. *Physiol. Plant.*, 77, 633–641.
- Teodoro, G.S., Lambers, H., Nascimento, D.L., de Britto Costa, P., Flores-Borges, D.N.A.,
Abrahão, A., *et al.* (2019). Specialized roots of Velloziaceae weather quartzite rock while
mobilizing phosphorus using carboxylates. *Funct. Ecol.*, 33, 762–773.
- de Tombeur, F., Vander Linden, C., Cornélis, J.-T., Godin, B., Compère, P. & Delvaux, B.
(2020a). Soil and climate affect foliar silicification patterns and silica-cellulose balance in
sugarcane (*Saccharum officinarum*). *Plant Soil*, 452, 529–546.
- de Tombeur, F., Turner, B.L., Laliberté, E., Lambers, H. & Cornelis, J.T. (2020b). Silicon
Dynamics During 2 Million Years of Soil Development in a Coastal Dune Chronosequence
Under a Mediterranean Climate. *Ecosystems*, 23, 1614–1630.
- de Tombeur, F., Turner, B.L., Laliberté, E., Lambers, H., Mahy, G., Faucon, M.P., *et al.* (2020c).
Plants sustain the terrestrial silicon cycle during ecosystem retrogression. *Science* (80-.),
369, 1245–1248.
- Trembath-Reichert, E., Wilson, J.P., McGlynn, S.E. & Fischer, W.W. (2015). Four hundred
million years of silica biomineralization in land plants. *Proc. Natl. Acad. Sci. U. S. A.*, 112,
5449–54.
- Turner, B.L., Hayes, P.E. & Laliberté, E. (2018). A climosequence of chronosequences in
southwestern Australia. *Eur. J. Soil Sci.*, 69, 69–85.
- Turner, B.L. & Laliberté, E. (2015). Soil development and nutrient availability along a 2 million-
year coastal dune chronosequence under species-rich mediterranean shrubland in
Southwestern Australia. *Ecosystems*, 18, 287–309.

- 628 Violle, C., Navas, M.L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., *et al.* (2007). Let the
629 concept of trait be functional! *Oikos*, 116, 882–892.
- 630 Vitousek, P. (2004). *Nutrient Cycling and Limitation - Hawai'i as Model System*. Princeton
631 University Press, New Jersey.
- 632 Vitousek, P.M., Walker, L.R., Whiteaker, L.D. & Matson, P.A. (1993). Nutrient limitations to
633 plant growth during primary succession in Hawaii Volcanoes National Park.
634 *Biogeochemistry*, 23, 197–215.
- 635 Waterman, J.M., Hall, C.R., Mikhael, M., Cazzonelli, C.I., Hartley, S.E. & Johnson, S.N. (2020).
636 Short-term resistance that persists: Rapidly induced silicon anti-herbivore defence affects
637 carbon-based plant defences. *Funct. Ecol.*
- 638 Wright, D.M., Jordan, G.J., Lee, W.G., Duncan, R.P., Forsyth, D.M. & Coomes, D.A. (2010). Do
639 leaves of plants on phosphorus-impooverished soils contain high concentrations of phenolic
640 defence compounds? *Funct. Ecol.*, 24, 52–61.
- 641 Wright, I.J., Westoby, M., Reich, P.B., Oleksyn, J., Ackerly, D.D., Baruch, Z., *et al.* (2004). The
642 worldwide leaf economics spectrum. *Nature*, 428, 821–827.
- 643 Zemunik, G., Turner, B.L., Lambers, H. & Laliberté, E. (2015). Diversity of plant nutrient-
644 acquisition strategies increases during long-term ecosystem development. *Nat. Plants*, 1, 1–
645 4.
- 646 Zemunik, G., Turner, B.L., Lambers, H. & Laliberté, E. (2016). Increasing plant species diversity
647 and extreme species turnover accompany declining soil fertility along a long-term
648 chronosequence in a biodiversity hotspot. *J. Ecol.*, 104, 792–805.

649 Zhang, L.H., Shao, H.B., Ye, G.F. & Lin, Y.M. (2012). Effects of fertilization and drought stress
650 on tannin biosynthesis of *Casuarina equisetifolia* seedlings branchlets. *Acta Physiol. Plant.*,
651 34, 1639–1649.

For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figures Captions

Fig. 1 – Leaf silicon (Si) concentrations across the chronosequence stages considering all individuals (i) and the means (ii) and cover-weighted means (CWM) (iii) of the five plots per stage (a). Leaf total phenol concentrations for the same chronosequence stages and plots (b). In the box-plots, small black dots represent each individual and large black dots represent outliers (outside 1.5*inter-quartile range). In the second and third panels, black dots indicate means and bars show 95% confidence intervals (n=5). In (a), Fisher LSD groupings (p<0.05) were performed on log-transformed data for the first plot and on root-square-transformed data for the two others, as the scales of the axes.

Fig. 2 – Soil total phosphorus (P) concentrations, total nitrogen (N) concentrations and soil N to P ratio versus cover-weighted mean (CWM) leaf silicon (Si) concentrations (a) and CWM leaf total phenol concentrations (b) (n = 25 plots). Black lines indicate the regression lines between both variables, shaded areas represent 95% confidence interval of the regression and colors of the circles indicate the chronosequence stages. Axes were log-transformed for soil total P concentration and N to P ratio and root-square-transformed for CWM leaf Si concentration. The p-values of the corresponding linear mixed-effect models are indicated if < 0.05. Regression lines were removed if the model p-values were > 0.05.

Fig. 3 – Leaf total phenol concentrations versus leaf silicon (Si) concentrations considering all individuals (a) and the means (b) and cover-weighted means (CWM) (c) of the five plots per stage. Scatterplot correlation matrix of leaf concentrations of Si, total phenols (Phnl), potassium (K), calcium (Ca), magnesium (Mg) and phosphorus (P) considering all individuals, the means, and CWM of the five plots per stage (d). In (a), (b) and (c), black lines indicate the regression lines between both variables, shaded areas represent 95% confidence interval of the regression and

691 colors of the circles indicate the chronosequence stages. Y-axes were log-transformed in (a) and
 692 root-square-transformed in (b) and (c). The p-values of the corresponding linear mixed-effect
 693 models are indicated if < 0.05 . In (d), the size and color of the squares represent Pearson's
 694 correlation coefficient. All correlations having a p-value > 0.01 are represented by a blank space.
 695 Plots on the diagonal represent the distributions of each variable. Concentrations are in gram per
 696 kg dry weight for the nutrients and gallic acid equivalent (GAE) gram per kg dry weight for total
 697 phenols. In (d), axes were log-transformed for all variables, except for leaf [phenols] in the first
 698 plot, while root-square-transformed for leaf [Si], [Ca] and [K] and log-transformed for leaf [P] and
 699 [Mg] in the second and third plots.

700 **Fig. 4** – Leaf silicon (Si) (a) and total phenol concentrations (b) across the chronosequence stages
 701 for Cyperaceae, and relationship between both for the same individuals (c). In (a) and (b), the small
 702 black dots represent each species and large black dots represent outliers (outside $1.5 \times$ inter-quartile
 703 range). In (a), Fisher LSD groupings ($p < 0.05$) were performed on root-square-transformed data, as
 704 the scale of the axes. In (c), the y-axis was root-square-transformed; each symbol represents a
 705 different species, the black line indicates the regression line between both variables, the shaded
 706 area represents 95% confidence interval of the regression, and the p-value of the corresponding
 707 linear mixed-effect model is indicated.

708 **Fig. 5** – Schematic representation (adapted from Lambers *et al.* 2008) of changes in soil total
 709 phosphorus (P) and nitrogen (N), P-acquisition strategies, and leaf defense strategies during long-
 710 term ecosystem development. Changes in soil P and N are based on Hayes *et al.* (2014). At the
 711 Jurien Bay chronosequence, soil total P concentration continually declines with increasing soil age,
 712 from about 430 to 6 mg kg⁻¹, while total N concentration increases from 0.5 to 1.2 g kg⁻¹ in the
 713 young soils where total [P] is relatively high, then declines to 0.2 g kg⁻¹ in the oldest soil, resulting

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

714 in a shift from N to P limitation of plant growth. The changes in nutrient-acquisition strategies were
715 originally suggested by Lambers *et al.* (2008), but subsequently confirmed in Zemunik *et al.*
716 (2015). Along the Jurien Bay chronosequence, the relative cover declines for arbuscular
717 mycorrhizal and ectomycorrhizal species, but increases for carboxylate-releasing cluster-rooted
718 species and others with functionally equivalent strategies. The changes in silica and phenol-based
719 defenses are based on the present study.

For Review Only

720 **Tables**

721 **Table 1** – Main properties of the five chronosequence stages used in this study. The chronosequence stages, dune system, geological
 722 formation and estimated soil age are based on Laliberté *et al.* (2012) and Turner & Laliberté (2015). The carbonate, total phosphorus (P)
 723 and nitrogen (N) concentrations, soil total N to total P ratio, pH-CaCl₂ and cation exchange capacity (CEC) are based on Hayes *et al.*
 724 (2014) and Zemunik *et al.* (2016). They result from seven soil samples (0-20 cm deep) taken in each of the five plots by chronosequence
 725 stage used in this study (n = 25 plots). The limiting nutrients are based on Laliberté *et al.* (2012) and Hayes *et al.* (2014). The expected
 726 silicon (Si) availability are from de Tombeur *et al.* (2020b, c).

Stage	Dune system	Geological formation	Estimated soil age (ka)	Carbonates* (%)	pH-CaCl ₂ [†]	CEC [†] (cmolc kg ⁻¹)	Total P [†] (mg kg ⁻¹)	Total N [†] (g kg ⁻¹)	Soil total N to total P ratio [†]	Limiting nutrients	Expected Si availability
1	Quindalup young	Safety Bay Sand	0.1 (Holocene)	75.2(3.5)	8.2(0.1)	30.5(4.5)	351.0(6.6)	0.5(0.0)	1.4(0.1)	N	Very low
2	Quindalup medium	Safety Bay Sand	1 (Holocene)	75.5(2.8)	7.8(0.0)	12.2(1.1)	424.4(8.3)	1.2(0.0)	2.7(0.4)	N, P and/or other nutrients	Very low
3	Quindalup old	Safety Bay Sand	6.7 (Holocene)	25.6(3.7)	7.8(0.0)	10.9(0.5)	205.7(7.0)	0.7(0.0)	3.6(0.5)	N, P and/or other nutrients	Low
4	Spearwood	Tamala Limestone	125 (Middle Pleistocene)	BDL	5.8(0.1)	3.6(0.2)	18.5(1.0)	0.2(0.0)	10.7(1.9)	P	Medium
5	Bassendean	Bassendean sand	>2000 (Early Pleistocene or Late Pliocene)	BDL	4.8(0.1)	2.5(0.2)	6.6(0.5)	0.2(0.0)	30.8(8.5)	P	Low

727 *SE is indicated in brackets. The seven soil samples were bulked before analysis (n=5).

728 [†]SE is indicated in brackets (n=35).

729 BDL, below detections limits









