Kepler's Third Law

r _{Earth}

The key to using the third law is to ensure that you use two orbiting bodies that share the same focus mass.

ratio of distances:

A.U. = \underline{r}_{object}




	body radius	orbital period	mean orbital radius	
	m		km	
Earth	6.3713 x 10 ⁶	l y	1.4957 x 10 ⁸	
Moon	1.737 x 10 ⁶	\mathbb{N}	3,85 x 10 ⁵	
ISS	$\supset <$	92.75 min		

conversion factors:


1 y = 365.2422 d 1 d = 23 h 56 min 4 s

(based on 400 year average) (solar day based on orbit of Sun)

- The biggest separation distance between Comet 67P and the Sun is 5.683 AU. Express this measurement in kilometres.
- By comparison, the biggest separation distance between Halley's Comet and the Sun is 5.25 x 10 9 km. Express this measurement in astronomical units.
- Halley's Comet has an orbital period of 75.3 y around the Sun. Use Kepler's third law to predict the mean orbital radius for Halley's orbit in kilometres.
- Mercury has an orbital period of 87.77 d. Use Kerpler's third law to predict the mean orbital radius of Mercury's orbit using astronomical units.
- A satellite placed in orbit around the Earth has a mean orbital radius equal to one third of the radius of the Moon's orbit around Earth. Use Kepler's third law to derive an expression for the period of the satellite based on the Moon's period for orbiting the Earth. (Not looking for a numeric answer. Looking for
- Suppose the ISS orbits at an altitude of 435 km above the Earth. Calculate the period of the Moon, in solar Earth days, orbiting around the Earth. (Numeric answer,)



https://en.wikipedia.org/ wiki/Halley% 27s Comet



https://en.wikipedia.org/wiki/ 67P/Churyumov%E2%80% 93Gerasimenko

2. 2.57 × 10 ° km 3. 2.57 × 10 ° km 4. 0.3865 AU 4. 5. 19 T_{moon} 5. 0.19 T_{moon} 6. 25.7 d

answers: 1, 8,500 x 10 * km 2, 35,1 AU

e distribute que qua estra a en como e



e de la companya de	en e	ara di	and the same of the same
			er er er i state i sta

Section of the section of			and the second of the second
the second second	:		and the second
	Carlos de para en la carlo	والأنابي والماء	and the second second
1 45-	1200		
errage contraction	and Art March	and the same	and the second of the
			The same of the
	Participation of the con-	Sec. 13.	de en likelinger
and from the second			* *

The control of the co


A second of the property of the control of the property of the control of the contr

which the standard contribution of the contribution of the γ - γ - γ


If the property of the property of the control of the

A supplied particular and a second of the control of

The second section of the second section of the second section of the second section of the second section of the second section of the second section of the second section of the second section of the second section section



4.4



Keplers 3rd Can Ws

$$\frac{\Gamma_{H}^{3}}{\Gamma_{H}^{2}} = \frac{\Gamma_{E}^{3}}{\Gamma_{E}^{2}} \rightarrow \Gamma_{H}^{3} = \sqrt{\frac{\Gamma_{E}^{3} \Gamma_{H}^{3}}{\Gamma_{E}^{2}}}$$

$$= \sqrt[3]{(1.4957 \times 10^{8} \text{Km})^{3} (75.34)^{2}}$$

$$(1.0004)^{2}$$