# The Constants of Nature

$$E_h \vdots \mathrm{K} = ж_1{}^4\left(\frac{T_p m_e}{m_p}\right)\left(1 - 14\, C_U{}^{-4}\, ж_\theta \boxtimes\right)$$
$$\mathrm{K} \vdots E_h = \frac{\text{kelvin}}{ж_1{}^4}\left(\frac{m_p}{T_p m_e}\right)\left(1 + 14\, C_U{}^{-4}\, ж_\theta \boxtimes\right)$$
$$\Delta_{\nu_{Cs}} = \frac{E_1}{2\pi}\left(\frac{t_p}{l_p{}^2 m_p}\right)\left(1 - \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\mathrm{eV} \vdots \mathrm{Hz} = \frac{\mathrm{eV}}{2\pi}\left(\frac{t_p}{l_p{}^2 m_p}\right)\left(1 - \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\mathrm{J} \vdots \mathrm{Hz} = \frac{\text{joule}}{2\pi}\left(\frac{t_p}{l_p{}^2 m_p}\right)\left(1 - \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\mathrm{Hz} \vdots \mathrm{J} = \frac{2\pi}{\text{second}}\left(\frac{l_p{}^2 m_p}{t_p}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\mathrm{Hz} \vdots \mathrm{eV} = \frac{2\pi\ \text{joule}}{\text{second eV}}\left(\frac{l_p{}^2 m_p}{t_p}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\hbar = \frac{\text{joule}}{\mathrm{eV}}\left(\frac{l_p{}^2 m_p}{t_p}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\hbar = \left(\frac{l_p{}^2 m_p}{t_p}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$h = 2\pi\left(\frac{l_p{}^2 m_p}{t_p}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$q_c = \pi\left(\frac{l_p{}^2 m_p}{t_p m_e}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$2q_c = 2\pi\left(\frac{l_p{}^2 m_p}{t_p m_e}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_B}{e} = \frac{\text{coulomb}}{2}\left(\frac{l_p{}^2 m_p}{t_p m_e}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_N}{e} = \frac{\text{coulomb}}{2}\left(\frac{l_p{}^2 m_p}{t_p m_+}\right)\left(1 + \sqrt{\zeta(2)}\, ж_\theta{}^2 \boxtimes\right)$$
$$G_0{}^{-1} = \frac{\pi}{ж_1{}^2}\left(\frac{l_p{}^2 m_p}{t_p q_p{}^2}\right)\left(1 + \sqrt{\zeta(3)}\, ж_\theta{}^2 \boxtimes\right)$$
$$G_0 = \frac{ж_1{}^2}{\pi}\left(\frac{t_p q_p{}^2}{l_p{}^2 m_p}\right)\left(1 - \sqrt{\zeta(3)}\, ж_\theta{}^2 \boxtimes\right)$$

$$\mathrm{eV} = \frac{\mathrm{eV}}{\text{coulomb}}\, ж_1\left(q_p\right)\left(1 + \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$
$$\mathrm{J} \vdots \mathrm{eV} = \frac{\text{joule coulomb}}{ж_1}\left(\frac{1}{q_p}\right)\left(1 - \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$
$$-\frac{e}{m_e} = -ж_1\left(\frac{q_p}{m_e}\right)\left(1 + \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$
$$\frac{e}{m_+} = ж_1\left(\frac{q_p}{m_+}\right)\left(1 + \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_B}{h} = \text{tesla}^2\,\frac{ж_1}{4\pi}\left(\frac{q_p}{m_e}\right)\left(1 + \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_N}{h} = \frac{1}{\text{second MHz}}\,\frac{ж_1}{4\pi}\left(\frac{q_p}{m_+}\right)\left(1 + \frac{e^{-\gamma}}{6}\, ж_\theta{}^2 \boxtimes\right)$$

$$g_e = -\frac{4\, ж_1 ж_2{}^2}{P_{up}}\left(1 + 4\left(4\, C_{CF} + 6\,(2\pi)\right) ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_B} = -\frac{2\, ж_1 ж_2{}^2}{P_{up}}\left(1 + 4\left(4\, C_{CF} + 6\,(2\pi)\right) ж_\theta{}^2 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_N} = -\frac{2\, ж_1 ж_2{}^2}{P_{up}}\left(\frac{m_+}{m_e}\right)\left(1 + 4\left(4\, C_{CF} + 6\,(2\pi)\right) ж_\theta{}^2 \boxtimes\right)$$

$$\varepsilon_0 = \frac{1}{4\pi}\left(\frac{t_p{}^2 q_p{}^2}{l_p{}^3 m_p}\right)\left(1 - \left(\frac{C_d}{6}\right) ж_\theta{}^4 \boxtimes\right)$$
$$\mathrm{A}_{\text{perm}} = \left(\frac{t_p{}^2 q_p{}^2}{l_p{}^3 m_p}\right)\left(1 - \left(\frac{C_d}{6}\right) ж_\theta{}^4 \boxtimes\right)$$
$$\kappa = \left(\frac{l_p{}^3 m_p}{t_p{}^2 q_p{}^2}\right)\left(1 + \left(\frac{C_d}{6}\right) ж_\theta{}^4 \boxtimes\right)$$
$$\mu_0 = 4\pi\left(\frac{l_p m_p}{q_p{}^2}\right)\left(1 + \sqrt{7}\left(\frac{2K}{6}\right) ж_\theta{}^4 \boxtimes\right)$$
$$R_\infty = \frac{ж_1{}^4}{4\pi}\left(\frac{m_e}{l_p m_p}\right)\left(1 - \frac{7}{5}\left(\frac{M}{18}\right) ж_\theta{}^4 \boxtimes\right)$$
$$E_h \vdots \frac{1}{\mathrm{m}} = \frac{ж_1{}^4}{2\pi}\left(\frac{m_e}{l_p m_p}\right)\left(1 - \frac{7}{5}\left(\frac{M}{18}\right) ж_\theta{}^4 \boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots E_h = \frac{1}{\text{meter}}\,\frac{2\pi}{ж_1{}^4}\left(\frac{l_p m_p}{m_e}\right)\left(1 + \frac{7}{5}\left(\frac{M}{18}\right) ж_\theta{}^4 \boxtimes\right)$$

$$\mathrm{A}_{\text{time}} = \frac{1}{ж_1{}^4}\left(\frac{t_p m_p}{m_e}\right)\left(1 + \frac{\sqrt{P}}{6}\, ж_\theta{}^4 \boxtimes\right)$$
$$\mathrm{Hz} \vdots E_h = \text{watt}\,\frac{2\pi}{ж_1{}^4}\left(\frac{t_p m_p}{m_e}\right)\left(1 + \frac{\sqrt{P}}{6}\, ж_\theta{}^4 \boxtimes\right)$$
$$E_h \vdots \mathrm{Hz} = \frac{ж_1{}^4}{2\pi}\left(\frac{m_e}{t_p m_p}\right)\left(1 - \frac{\sqrt{P}}{6}\, ж_\theta{}^4 \boxtimes\right)$$
$$R_\infty c = \frac{ж_1{}^4}{4\pi}\left(\frac{m_e}{t_p m_p}\right)\left(1 - \frac{\sqrt{P}}{6}\, ж_\theta{}^4 \boxtimes\right)$$

$$m_{\text{de}} = \frac{3s}{ж_3 ж_4}\left(\frac{16}{4\pi}\right)\left(m_n + m_+\right)\left(1 + \frac{Im(\omega_1)}{ж_1 ж_2}\left(\frac{L_2}{14}\right) ж_r{}^3 \boxtimes\right)$$
$$m_\alpha = \frac{1}{ж_3 ж_4}\sqrt{\frac{16}{4\pi}}\left(m_n + m_+\right)\left(1 + \left(\frac{16}{2\pi}\right) ж_r{}^3 \boxtimes\right)$$
$$m_+ = \frac{2\pi}{\sqrt{ж_3 ж_4}}\left(\frac{\mathrm{GeV}}{\sqrt{5}}\right)\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 + \frac{1}{6^{1/3}}\left(\frac{2\pi}{2s}\right) ж_r{}^3 \boxtimes\right)$$
$$m_\alpha = \frac{16}{18}\left(\frac{7\,\mathrm{GeV}}{2 G_{Gi}}\right)\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 - s\left(\frac{s}{2\pi}\right)^{1/3} ж_r{}^3 \boxtimes\right)$$
$$m_{\text{he}} = \frac{\sqrt{3}}{ж_1{}^2 ж_2{}^2}\left(\frac{\mathrm{GeV}}{2\pi}\right)\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 + \frac{3^{-1}}{ж_1 ж_2}\left(\frac{s}{18}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{G}{\hbar c} = \mathrm{GeV}^2\left(\frac{t_p{}^2}{l_p{}^2 m_p}\right)^2\left(1 - \left(\frac{4\pi}{18}\right) ж_r{}^3 \boxtimes\right)$$
$$G = \left(\frac{l_p{}^3}{t_p{}^2 m_p}\right)\left(1 - \left(\frac{4\pi}{18}\right) ж_r{}^3 \boxtimes\right)$$
$$m_e = 2V_{fe}\left(\frac{m_p{}^4}{\text{kilogram}^3}\right)\left(1 + \sqrt{\frac{18}{2V_{fe}}}\, ж_r{}^3 \boxtimes\right)$$
$$m_{\text{tri}} = \frac{ж_1{}^{-5}}{18}\sqrt{\frac{4s}{2\pi}}\left(\frac{m_p{}^4}{\text{kilogram}^3}\right)\left(1 + 3\,\omega_1{}^3\, ж_r{}^3 \boxtimes\right)$$
$$K_{cd} = \frac{ж_1{}^{-7}}{f_1}\sqrt{\frac{4s}{2\pi}}\left(\frac{t_p{}^2}{l_p{}^2 \mathrm{A}_{\text{mass}}}\right)\left(1 + 3\,\omega_1{}^3\, ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{de}}}{\mu_n} = -C_{CFP}\sqrt{\frac{2s}{6\pi}}\left(\frac{m_n}{m_{\text{de}}}\right)\left(1 - ж_1 ж_2\left(\frac{35}{L_1}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_{\text{de}}} = -\frac{1}{P_{up}}\sqrt{\frac{6\pi}{2s}}\left(\frac{m_{\text{de}}}{m_e}\right)\left(1 + \frac{5}{ж_1 ж_2}\left(\frac{L_1}{G_{Gi}}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{de}}}{\mu_e} = -P_{up}\sqrt{\frac{2s}{6\pi}}\left(\frac{m_e}{m_{\text{de}}}\right)\left(1 - \frac{5}{ж_1 ж_2}\left(\frac{L_1}{G_{Gi}}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{de}}}{\mu_B} = S\sqrt{\frac{2s}{6\pi}}\left(\frac{m_+}{m_{\text{de}}}\right)\left(1 - \frac{1}{ж_1 ж_2}\left(\frac{7s}{12}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_{\text{de}} = ж_1{}^2 ж_2{}^2\sqrt{\frac{2s}{6\pi}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_{\text{de}}}\right)\left(1 - \frac{5}{ж_1 ж_2}\left(\frac{7s}{!5}\right) ж_r{}^3 \boxtimes\right)$$
$$\mathrm{A}_{\text{mdm}} = ж_1\left(\frac{l_p{}^2 q_p m_p}{t_p m_e}\right)\left(1 + ж_1 ж_2\left(\frac{2\pi}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_B = \frac{ж_1}{2}\left(\frac{l_p{}^2 q_p m_p}{t_p m_e}\right)\left(1 + ж_1 ж_2\left(\frac{2\pi}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_N = \frac{ж_1}{2}\left(\frac{l_p{}^2 q_p m_p}{t_p m_+}\right)\left(1 + ж_1 ж_2\left(\frac{2\pi}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_+ = \frac{ж_1{}^2 ж_2{}^2}{S}\left(\frac{l_p{}^2 q_p m_p}{t_p m_+}\right)\left(1 - \frac{S\, P_{up}}{ж_1 ж_2}\left(\frac{6\pi}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_+{}' = \frac{ж_1{}^2 ж_2{}^2}{S}\left(\frac{l_p{}^2 q_p m_p}{t_p m_+}\right)\left(1 - \frac{1}{ж_1 ж_2}\left(\frac{C_{QA}}{G_{Gi}}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_e = -\frac{ж_1{}^2 ж_2{}^2}{P_{up}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_e}\right)\left(1 + \frac{14\, F_{FF}}{ж_1 ж_2}\left(\frac{2\pi}{9s}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_\mu = -\frac{ж_1{}^2 ж_2{}^2}{P_{up}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_\mu}\right)\left(1 + \frac{1}{ж_1 ж_2}\left(\frac{3s}{2\pi}\right) ж_r{}^3 \boxtimes\right)$$
$$\mu_{\text{tri}} = ж_1{}^2 ж_2{}^2\left(D_{Do} + \pi\right)\left(\frac{l_p{}^2 q_p m_p}{t_p m_{\text{tri}}}\right)\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{4\pi}{s}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{tri}}}{\mu_+} = S\left(D_{Do} + \pi\right)\left(\frac{m_+}{m_{\text{tri}}}\right)\left(1 + ж_3 ж_4\left(\frac{2\pi}{7s}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{tri}}}{\mu_N} = ж_1 ж_2{}^2\left(2D_{Do} + 2\pi\right)\left(\frac{m_+}{m_{\text{tri}}}\right)\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{2\pi}{\sqrt{3}}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{tri}}}{\mu_B} = ж_1 ж_2{}^2\left(2D_{Do} + 2\pi\right)\left(\frac{m_e}{m_{\text{tri}}}\right)\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{2\pi}{\sqrt{3}}\right) ж_r{}^3 \boxtimes\right)$$
$$g_{\text{tri}} = ж_1 ж_2{}^2\left(4D_{Do} + 4\pi\right)\left(\frac{m_+}{m_{\text{tri}}}\right)\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{2\pi}{\sqrt{3}}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{he}}{}'}{\mu_B} = -\frac{1}{2}\sqrt{\frac{1}{4\pi}\frac{1}{4s}}\left(\frac{m_e}{m_{\text{he}}}\right)\left(1 + ж_3 ж_4\, s\left(\frac{L_{LL}}{18}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{he}}{}'}{\mu_N} = -\frac{1}{2}\sqrt{\frac{1}{4\pi}\frac{1}{4s}}\left(\frac{m_+ m_e}{m_e m_{\text{he}}}\right)\left(1 + ж_3 ж_4\, s\left(\frac{L_{LL}}{18}\right) ж_r{}^3 \boxtimes\right)$$
$$g_{\text{he}} = -\sqrt{\frac{1}{4\pi}\frac{1}{4s}}\left(\frac{m_+ m_e}{m_e m_{\text{he}}}\right)\left(1 + ж_3 ж_4\, 7\left(\frac{j_{0,1}}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{he}}}{\mu_N} = -\frac{1}{2}\sqrt{\frac{1}{4\pi}\frac{1}{4s}}\left(\frac{m_+ m_e}{m_e m_{\text{he}}}\right)\left(1 + ж_3 ж_4\, 7\left(\frac{j_{0,1}}{32}\right) ж_r{}^3 \boxtimes\right)$$
$$\frac{\mu_{\text{he}}}{\mu_B} = -\frac{1}{2}\sqrt{\frac{1}{4\pi}\frac{1}{4s}}\left(\frac{m_e}{m_{\text{he}}}\right)\left(1 + ж_3 ж_4\, 7\left(\frac{j_{0,1}}{32}\right) ж_r{}^3 \boxtimes\right)$$

$$\frac{\mu_+}{\mu_n} = -\frac{C_{CFP}}{S}\left(\frac{m_n}{m_+}\right)\left(1 - \frac{6}{ж_3 ж_4}\left(2\pi\right) ж_r{}^4 \boxtimes\right)$$
$$\frac{\mu_n}{\mu_+} = -\frac{S}{C_{CFP}}\left(\frac{m_+}{m_n}\right)\left(1 + \frac{6}{ж_3 ж_4}\left(2\pi\right) ж_r{}^4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_n} = \frac{C_{CFP}}{P_{up}}\left(\frac{m_n}{m_e}\right)\left(1 + \frac{ж_3 ж_4}{6}\left(\frac{1}{2\pi}\right) ж_r{}^4 \boxtimes\right)$$
$$\frac{\mu_n}{\mu_e} = \frac{P_{up}}{C_{CFP}}\left(\frac{m_e}{m_n}\right)\left(1 - \frac{ж_3 ж_4}{6}\left(\frac{1}{2\pi}\right) ж_r{}^4 \boxtimes\right)$$

$$E_p{}' = \frac{\text{joule}}{\mathrm{GeV}}\left(\frac{l_p{}^2 m_p}{t_p{}^2}\right)\left(1 - \frac{1}{2}Re\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$R_K = \frac{2\pi}{ж_1{}^2}\left(\frac{l_p{}^2 m_p}{t_p q_p{}^2}\right)\left(1 + \frac{1}{2}Re\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$Z_0 = 4\pi\left(\frac{l_p{}^2 m_p}{t_p q_p{}^2}\right)\left(1 + \frac{1}{2}Re\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$Z_p = \left(\frac{l_p{}^2 m_p}{t_p q_p{}^2}\right)\left(1 + \frac{1}{2}Re\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{Hz} \vdots \frac{1}{\mathrm{m}} = \frac{1}{\text{second}}\left(\frac{t_p}{l_p}\right)\left(1 + \frac{1}{2}Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{A}_{\text{vel}} = ж_1{}^2\left(\frac{l_p}{t_p}\right)\left(1 + \frac{1}{2}Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{A}_{\text{mom}} = ж_1{}^2\left(\frac{l_p m_e}{t_p}\right)\left(1 - \frac{1}{2}Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathcal{N}_{\text{mom}} = \left(\frac{l_p m_e}{t_p}\right)\left(1 - \frac{1}{2}Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathcal{N}_{\text{mom}}{}' = \frac{\text{joule second}}{\text{MeV meter}}\left(\frac{l_p{}^2 m}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\mathrm{A}_{\text{mass}}} = \frac{\mathrm{A}_{\text{mass}}}{\mathrm{MeV}}\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{kg} \vdots \mathrm{J} = \text{kilogram}\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{J} \vdots \mathrm{kg} = \text{joule}\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 + Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} \vdots \mathrm{J} = \mathrm{A}_{\text{mass}}\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{J} \vdots \mathrm{A}_{\text{mass}} = \frac{\text{joule}}{\mathrm{A}_{\text{mass}}}\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 + Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{kg} \vdots \mathrm{eV} = \frac{\text{joule kilogram}}{\mathrm{eV}}\left(\frac{t_p{}^2}{l_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{eV} \vdots \mathrm{kg} = \mathrm{eV}\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 + Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\Delta = \left(m_n - m_+\right)\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\Delta{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(m_n - m_+\right)\left(\frac{l_p{}^2}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_e{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_e}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\mu{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_\mu}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_+{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_+}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_n{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_n}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\tau{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_\tau}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{de}}{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_{\text{de}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{he}}{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_{\text{he}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{tri}}{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_{\text{tri}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\alpha{}' = \frac{\text{joule}}{\mathrm{MeV}}\left(\frac{l_p{}^2 m_\alpha}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_e = \left(\frac{l_p{}^2 m_e}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\mu = \left(\frac{l_p{}^2 m_\mu}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_+ = \left(\frac{l_p{}^2 m_+}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_n = \left(\frac{l_p{}^2 m_n}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\tau = \left(\frac{l_p{}^2 m_\tau}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{de}} = \left(\frac{l_p{}^2 m_{\text{de}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{he}} = \left(\frac{l_p{}^2 m_{\text{he}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_{\text{tri}} = \left(\frac{l_p{}^2 m_{\text{tri}}}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{E}_\alpha = \left(\frac{l_p{}^2 m_\alpha}{t_p{}^2}\right)\left(1 - Im\left(i^{i^{\cdot^{\cdot}}}\right) ж_3 ж_4 \boxtimes\right)$$

$$\frac{\mu_B}{\hbar c} = \frac{ж_1}{4\pi}\left(\frac{t_p q_p}{l_p m_e}\right)\left(1 + \sqrt{\frac{4}{D_{Do}}}\, ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_N}{\hbar c} = \frac{ж_1}{4\pi}\left(\frac{t_p q_p}{l_p m_+}\right)\left(1 + \sqrt{\frac{4}{D_{Do}}}\, ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{A}_{90} = \text{ampere}\left(1 + \sqrt{\frac{9}{j_{0,1}}}\, ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\mathrm{C}_{90} = \text{coulomb}\left(1 + \sqrt{\frac{9}{j_{0,1}}}\, ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_+{}'}{\mu_n} = \frac{2\, ж_1 ж_2{}^2}{S}\left(1 - 18\left(6\pi\right) ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_+{}'}{\mu_B} = \frac{2\, ж_1 ж_2{}^2}{S}\left(\frac{m_e}{m_+}\right)\left(1 - 18\left(6\pi\right) ж_1 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_\mu} = \left(\frac{m_\mu}{m_e}\right)\left(1 - 18\left(V_{fe}\right) ж_1 ж_3 ж_4 \boxtimes\right)$$
$$R = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{\text{coulomb}}{\text{meter}}\right)\left(\frac{l_p{}^2}{t_p{}^2 q_p T_p}\right)\left(1 - \frac{8}{C_d}\, ж_1{}^4 ж_3{}^4 ж_4{}^4 \boxtimes\right)$$
$$V_{m_0} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{\text{coulomb}\ T_0}{\text{meter}\ p_0}\right)\left(\frac{l_p{}^2}{t_p{}^2 q_p T_p}\right)\left(1 - \frac{8}{C_d}\, ж_1{}^4 ж_3{}^4 ж_4{}^4 \boxtimes\right)$$
$$V_{m_1} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{\text{coulomb}\ T_0}{\text{meter}\ p_1}\right)\left(\frac{l_p{}^2}{t_p{}^2 q_p T_p}\right)\left(1 - \frac{8}{C_d}\, ж_1{}^4 ж_3{}^4 ж_4{}^4 \boxtimes\right)$$
$$\frac{\mu_n}{\mu_+{}'} = -\frac{S}{C_{CFP}}\left(\frac{m_+}{m_n}\right)\left(1 + 35\left(\frac{2\pi}{16}\right) ж_1{}^8 ж_3{}^8 ж_4{}^8 \boxtimes\right)$$
$$\frac{\mu_+}{\mu_e} = -\frac{S}{P_{up}}\left(\frac{m_e}{m_+}\right)\left(1 + 35\left(\frac{2\pi}{3s}\right) ж_1{}^8 ж_3{}^8 ж_4{}^8 \boxtimes\right)$$
$$\gamma_e = \frac{2\, ж_1{}^2 ж_2{}^2}{P_{up}}\left(\frac{q_p}{m_e}\right)\left(1 + 7\left(\frac{3s}{4\pi}\right) ж_1{}^8 ж_3{}^8 ж_4{}^8 \boxtimes\right)$$
$$\dot{\gamma}_e = \frac{ж_1{}^2 ж_2{}^2}{\pi P_{up}}\,\frac{1}{\text{MHz second}}\left(\frac{q_p}{m_e}\right)\left(1 + 7\left(\frac{3s}{4\pi}\right) ж_1{}^8 ж_3{}^8 ж_4{}^8 \boxtimes\right)$$
$$\dot{\gamma}_+{}' = \frac{ж_1{}^2 ж_2{}^2}{\pi S}\,\frac{1}{\text{MHz second}}\left(\frac{q_p}{m_+}\right)\left(1 - \frac{\sqrt{6}}{C_d}\left(\frac{6\pi}{ж_3 ж_4}\right) ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\gamma_+{}' = \frac{2\, ж_1{}^2 ж_2{}^2}{S}\left(\frac{q_p}{m_+}\right)\left(1 - \frac{\sqrt{6}}{C_d}\left(\frac{6\pi}{ж_3 ж_4}\right) ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_+} = -\frac{S}{P_{up}}\left(\frac{m_+}{m_e}\right)\left(1 + \sqrt{12}\left(\frac{ж_3 ж_4}{2\pi}\right) ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_+{}'} = -\frac{S}{P_{up}}\left(\frac{m_+}{m_e}\right)\left(1 + \left(\frac{s}{ж_3 ж_4}\right) ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_+}{\mu_B} = \frac{2\, ж_1 ж_2{}^2}{S}\left(\frac{m_e}{m_+}\right)\left(1 - \frac{1}{18}\left(\frac{2\pi}{4K}\right)^2 ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_+}{\mu_N} = \frac{2\, ж_1 ж_2{}^2}{S}\left(1 - \frac{1}{18}\left(\frac{2\pi}{4K}\right)^2 ж_2 ж_3 ж_4 \boxtimes\right)$$
$$g_+ = \frac{4\, ж_1 ж_2{}^2}{S}\left(1 - \frac{1}{12}\left(\frac{2\pi}{4K}\right)^2 ж_2 ж_3 ж_4 \boxtimes\right)$$
$$d_{220} = \frac{1}{\sqrt{8}}\,\frac{4\pi}{32\,s}\,\frac{1}{ж_1{}^4}\left(\frac{l_p m_p}{m_e}\right)\left(1 + 18\, G_{Gi}\, ж_2 ж_3 ж_4 \boxtimes\right)$$
$$a = \frac{4\pi}{32\,s}\,\frac{1}{ж_1{}^4}\left(\frac{l_p m_p}{m_e}\right)\left(1 + 18\, G_{Gi}\, ж_2 ж_3 ж_4 \boxtimes\right)$$
$$g_p = -\frac{4\, ж_1 ж_2{}^2}{P_{up}}\left(1 + 16\, C_Q\, ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_B} = -\frac{2\, ж_1 ж_2{}^2}{P_{up}}\left(\frac{m_e}{m_e}\right)\left(1 + 16\, C_Q\, ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\frac{\mu_e}{\mu_N} = -\frac{2\, ж_1 ж_2{}^2}{P_{up}}\left(\frac{m_+}{m_e}\right)\left(1 + 16\, C_Q\, ж_2 ж_3 ж_4 \boxtimes\right)$$
$$\gamma_+ = \frac{2\, ж_1{}^2 ж_2{}^2}{S}\left(\frac{q_p}{m_+}\right)\left(1 - \frac{1}{\sqrt{8}}\left(\frac{C_U}{5}\right) ж_2{}^2 ж_3{}^2 ж_4{}^2 \boxtimes\right)$$
$$\dot{\gamma}_+ = \frac{ж_1{}^2 ж_2{}^2}{\pi S}\,\frac{\text{second}}{\text{MHz}}\left(\frac{q_p}{m_+}\right)\left(1 - \frac{1}{\sqrt{8}}\left(\frac{C_U}{5}\right) ж_2{}^2 ж_3{}^2 ж_4{}^2 \boxtimes\right)$$
$$m_n = \sqrt{G_{Gi}}\left(\mathrm{A}_{\text{mass}}\right)\left(1 + \sqrt{!5 \cdot 32}\left(\frac{ж_3 ж_4}{s}\right) ж_2{}^2 ж_3{}^2 ж_4{}^2 \boxtimes\right)$$
$$m_\tau = \frac{\Gamma(5)}{4\pi}\left(\mathrm{A}_{\text{mass}}\right)\left(1 - \sqrt{!5 \cdot 32}\left(\frac{ж_3 ж_4}{s}\right) ж_2{}^2 ж_3{}^2 ж_4{}^2 \boxtimes\right)$$
$$a_\mu = \frac{2\, ж_1 ж_2{}^2}{P_{up}} - 1\left(1 - \frac{18}{35}\sqrt{\frac{L_1}{!5}}\left(\frac{ж_1 ж_2}{2s}\right) ж_2{}^3 ж_3{}^3 ж_4{}^3 \boxtimes\right)$$
$$a_e = \frac{2\, ж_1 ж_2{}^2}{P_{up}} - 1\left(1 - \frac{36}{!5}\sqrt{\frac{\sqrt{8}}{4\pi}}\left(\frac{ж_1 ж_2}{2\pi}\right)^2 ж_2{}^3 ж_3{}^3 ж_4{}^3 \boxtimes\right)$$

$$c = \left(\frac{l_p}{t_p}\right)\left(1 + \sqrt{G_{Gi}}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{Hz} = \frac{1}{\text{meter}}\left(\frac{l_p}{t_p}\right)\left(1 + \sqrt{G_{Gi}}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$r_e = ж_1{}^2\left(\frac{l_p m_p}{m_e}\right)\left(1 - \sqrt{6}\, C_{R1}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$a_0 = \frac{1}{ж_1{}^2}\left(\frac{l_p m_p}{m_e}\right)\left(1 - \sqrt{6}\, C_{R1}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\sigma_e = \sqrt{\frac{32}{18}}\, 2\pi\, ж_1{}^4\left(\frac{l_p m_p}{m_e}\right)^2\left(1 - 2\sqrt{6}\, C_{R1}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{G_F}{(\hbar c)^3} = \sqrt{\frac{32}{18}}\,\frac{(2\pi)^2}{\text{kilogram}}\left(\frac{\text{joule}}{\mathrm{GeV}}\right)^2\left(\frac{t_p{}^4}{l_p{}^4 m_p}\right)\left(1 + 28\sqrt{6}\, C_{R1}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{eV} \vdots \mathrm{A}_{\text{mass}} = \frac{\text{joule}^2}{\text{coulomb}}\sqrt{\frac{32\, ж_1{}^8}{18\, P_{up}}}\left(\frac{t_p{}^2 q_p}{l_p{}^2 m_e}\right)\left(1 - \alpha_F\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} \vdots \mathrm{eV} = \text{coulomb}\sqrt{\frac{18\, P_{up}}{32\, ж_1{}^8}}\left(\frac{l_p{}^2 m_e}{t_p{}^2 q_p}\right)\left(1 + \alpha_F\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{J} \vdots \frac{1}{\mathrm{m}} = \frac{\text{joule}}{2\pi}\left(\frac{t_p{}^2}{l_p{}^3 m_p}\right)\left(1 + \frac{2}{\delta_F}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{eV} \vdots \frac{1}{\mathrm{m}} = \frac{\mathrm{eV}}{2\pi}\left(\frac{t_p{}^2}{l_p{}^3 m_p}\right)\left(1 + \frac{2}{\delta_F}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{eV} = \frac{2\pi}{\text{meter eV}}\left(\frac{l_p{}^3 m_p}{t_p{}^2}\right)\left(1 - \frac{2}{\delta_F}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{J} = \frac{2\pi}{\text{meter}}\left(\frac{l_p{}^3 m_p}{t_p{}^2}\right)\left(1 - \frac{2}{\delta_F}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\hbar c = \frac{\text{joule meter}}{\text{MeV fm}}\left(\frac{l_p{}^3 m_p}{t_p{}^2}\right)\left(1 - \frac{2}{\delta_F}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} = \sqrt{\frac{18\, P_{up}}{32\, ж_1{}^6}}\left(m_e\right)\left(1 + \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{kg} \vdots \mathrm{A}_{\text{mass}} = \text{kilogram}\sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{1}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(e) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(+) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_+}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(n) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_n}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(\text{de}) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{de}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(\text{he}) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{he}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(\text{tri}) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{tri}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_r(\alpha) = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_\alpha}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_e}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_\Delta}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_n - m_+}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_\mu}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_\mu}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_+}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_+}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_n}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_n}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_\tau}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_\tau}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_{\text{de}}}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{de}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_{\text{he}}}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{he}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_{\text{tri}}}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_{\text{tri}}}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\frac{m_\alpha}{\mathrm{A}_{\text{mass}}} = \sqrt{\frac{32\, ж_1{}^6}{18\, P_{up}}}\left(\frac{m_\alpha}{m_e}\right)\left(1 - \frac{1}{L}\left(ж_1 - ж_2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{e pot}} = \sqrt{\frac{32\, ж_1{}^{12}}{18\, P_{up}}}\left(\frac{l_p{}^2 \mathrm{A}_{\text{mass}}}{t_p{}^2 q_p}\right)\left(1 + \left(2\pi - 6\, Im(\omega_1)\right)\left(ж_1 + ж_2\right)\boxtimes\right)$$
$$R_\infty hc = \frac{\text{coulomb}}{2}\sqrt{\frac{32\, ж_1{}^{12}}{18\, P_{up}}}\left(\frac{l_p{}^2 \mathrm{A}_{\text{mass}}}{t_p{}^2 q_p}\right)\left(1 + \left(2\pi - 6\, Im(\omega_1)\right)\left(ж_1 + ж_2\right)\boxtimes\right)$$
$$E_h \vdots \mathrm{eV} = \text{coulomb}\sqrt{\frac{32\, ж_1{}^{12}}{18\, P_{up}}}\left(\frac{l_p{}^2 \mathrm{A}_{\text{mass}}}{t_p{}^2 q_p}\right)\left(1 + \left(2\pi - 6\, Im(\omega_1)\right)\left(ж_1 + ж_2\right)\boxtimes\right)$$
$$\mathrm{eV} \vdots E_h = \frac{1}{\text{coulomb}}\sqrt{\frac{18\, P_{up}}{32\, ж_1{}^{12}}}\left(\frac{t_p{}^2 q_p}{l_p{}^2 \mathrm{A}_{\text{mass}}}\right)\left(1 - \left(2\pi - 6\, Im(\omega_1)\right)\left(ж_1 + ж_2\right)\boxtimes\right)$$

$$\mathrm{F}_{90} = \text{farad}\left(1 + \frac{C_d}{6}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\mathrm{H}_{90} = \text{henry}\left(1 - \frac{C_d}{6}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\Omega_{90} = \text{ohm}\left(1 - \frac{C_d}{6}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$ST_0 = \frac{5}{2} + \log\left(\left(\frac{1}{2\pi}\,\frac{\mathrm{K}\,\mathrm{A}_{\text{mass}}}{l_p{}^2 T_p m_p}\right)^{3/2}\frac{\mathrm{K}\, E_p}{T_p\, p_0}\right)\left(1 - \frac{1}{12}\left(\frac{s^2}{ж_1 ж_2}\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$ST_1 = \frac{5}{2} + \log\left(\left(\frac{1}{2\pi}\,\frac{\mathrm{K}\,\mathrm{A}_{\text{mass}}}{l_p{}^2 T_p m_p}\right)^{3/2}\frac{\mathrm{K}\, E_p}{T_p\, p_1}\right)\left(1 - \frac{1}{12}\left(\frac{s^2}{ж_1 ж_2}\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\sigma = \frac{3}{5}\left(\frac{4\pi}{\Gamma(5)}\right)^2\left(\frac{m_p}{t_p{}^3 T_p{}^4}\right)\left(1 + \sqrt{\frac{!5}{8}}\left(\frac{2\pi}{ж_1 ж_2}\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\sigma_+{}' = 1 - \left(1 - 35\sqrt{\frac{s^8}{4^2\,(2\pi)^5}}\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\sigma_{\text{he}} = 1 - \left(1 - 2\cdot 35\sqrt{7\, G_{Ga}{}^{-7}}\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{ef}} = ж_1{}^5\left(\frac{l_p m_e{}^2}{t_p{}^2 q_p m_p}\right)\left(1 + 18\sqrt{\frac{s^8}{24\,(2\pi)^7}}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{current}} = ж_1{}^5\left(\frac{q_p m_e}{t_p m_p}\right)\left(1 + 18\sqrt{\frac{s^8}{4^{\sqrt{2}}\,(2\pi)^8}}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{force}} = ж_1{}^6\left(\frac{l_p m_e{}^2}{t_p{}^2 m_p}\right)\left(1 + 18\sqrt{\frac{s^8}{4\,(2\pi)^8}}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{cd}} = ж_1{}^7\left(\frac{q_p m_e{}^3}{l_p{}^3 m_p{}^3}\right)\left(1 + 18\sqrt{\frac{s^9}{8\,(2\pi)^8}}\left(ж_1 + ж_3 + ж_4\right)\boxtimes\right)$$
$$\mu_n = -\frac{ж_1{}^2 ж_2{}^2}{C_{CFP}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\dot{\gamma}_n = \frac{ж_1{}^2 ж_2{}^2}{\pi\, C_{CFP}}\,\frac{1}{\text{MHz second}}\left(\frac{q_p}{m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\gamma_n = \frac{2\, ж_1{}^2 ж_2{}^2}{C_{CFP}}\left(\frac{q_p}{m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\frac{\mu_n}{\mu_B} = -\frac{2\, ж_1 ж_2{}^2}{C_{CFP}}\left(\frac{m_e}{m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\frac{\mu_n}{\mu_N} = -\frac{2\, ж_1 ж_2{}^2}{C_{CFP}}\left(\frac{m_+}{m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$g_n = -\frac{4\, ж_1 ж_2{}^2}{C_{CFP}}\left(\frac{m_+}{m_n}\right)\left(1 + 18\left(2\pi\right)\left(ж_1 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{Hz} \vdots \mathrm{kg} = \frac{2\pi}{\text{second}}\left(t_p m_p\right)\left(1 + \frac{2}{C_{CFP}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{Hz} \vdots \mathrm{A}_{\text{mass}} = \frac{2\pi}{\text{second}}\left(\frac{t_p m_p}{\mathrm{A}_{\text{mass}}}\right)\left(1 + \frac{2}{C_{CFP}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathcal{N}_{\text{time}} = \left(\frac{t_p m_p}{m_e}\right)\left(1 + \frac{2}{C_{CFP}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} \vdots \mathrm{Hz} = \frac{1}{2\pi}\left(\frac{\mathrm{A}_{\text{mass}}}{t_p m_p}\right)\left(1 - \frac{2}{C_{CFP}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{kg} \vdots \mathrm{Hz} = \frac{\text{kilogram}}{2\pi}\left(\frac{1}{t_p m_p}\right)\left(1 - \frac{2}{C_{CFP}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$R_\infty hc = \frac{ж_1{}^4}{2}\left(\frac{l_p{}^2 m_e}{t_p{}^2}\right)\left(1 + \left(s - 2\pi\right)\frac{6\pi}{ж_3 ж_4}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$E_h = ж_1{}^4\left(\frac{l_p{}^2 m_e}{t_p{}^2}\right)\left(1 + \left(s - 2\pi\right)\frac{6\pi}{ж_3 ж_4}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{J} \vdots E_h = \frac{\text{joule}^2}{ж_1{}^4}\left(\frac{t_p{}^2}{l_p{}^2 m_e}\right)\left(1 - \left(s - 2\pi\right)\frac{6\pi}{ж_3 ж_4}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$k_B = \left(\frac{l_p{}^2 m_p}{t_p{}^2 T_p}\right)\left(1 + 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\dddot{k}_B = \frac{\text{joule}}{\mathrm{eV}}\left(\frac{l_p{}^2 m_p}{t_p{}^2 T_p}\right)\left(1 + 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{K} \vdots \mathrm{J} = \text{kelvin}\left(\frac{l_p{}^2 m_p}{t_p{}^2 T_p}\right)\left(1 + 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{K} \vdots \mathrm{eV} = \frac{\text{kelvin joule}}{\mathrm{eV}}\left(\frac{l_p{}^2 m_p}{t_p{}^2 T_p}\right)\left(1 + 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{eV} \vdots \mathrm{K} = \mathrm{eV}\left(\frac{t_p{}^2 T_p}{l_p{}^2 m_p}\right)\left(1 - 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\mathrm{J} \vdots \mathrm{K} = \text{joule}\left(\frac{t_p{}^2 T_p}{l_p{}^2 m_p}\right)\left(1 - 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$n_0 = \frac{p_0}{T_0}\left(\frac{t_p{}^2 T_p}{l_p{}^2 m_p}\right)\left(1 - 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$n_1 = \frac{p_1}{T_0}\left(\frac{t_p{}^2 T_p}{l_p{}^2 m_p}\right)\left(1 - 18\, C_{HSM}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\frac{\mu_B}{k_B} = \frac{ж_1}{2}\left(\frac{t_p q_p T_p}{m_e}\right)\left(1 - \sqrt{\frac{32}{G_{Gi}}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$
$$\frac{\mu_N}{k_B} = \frac{ж_1}{2}\left(\frac{t_p q_p T_p}{m_+}\right)\left(1 - \sqrt{\frac{32}{G_{Gi}}}\left(ж_2 - ж_3 - ж_4\right)\boxtimes\right)$$

$$K_J = \frac{ж_1}{\pi}\left(\frac{t_p q_p}{l_p{}^2 m_p}\right)\left(1 - \frac{7}{8}\left(\frac{8}{4\pi}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\frac{e}{\hbar} = ж_1\left(\frac{t_p q_p}{l_p{}^2 m_p}\right)\left(1 - \frac{7}{8}\left(\frac{8}{4\pi}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\Phi_0 = \frac{\pi}{ж_1}\left(\frac{l_p{}^2 m_p}{t_p q_p}\right)\left(1 + \frac{7}{8}\left(\frac{8}{4\pi}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\frac{\hbar}{e} = \frac{\text{coulomb}}{ж_1}\left(\frac{l_p{}^2 m_p}{t_p q_p}\right)\left(1 + \frac{7}{8}\left(\frac{8}{4\pi}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\frac{h}{e} = 2\pi\,\frac{\text{coulomb}}{ж_1}\left(\frac{l_p{}^2 m_p}{t_p q_p}\right)\left(1 + \frac{7}{8}\left(\frac{8}{4\pi}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$c_1 = (2\pi)^2\left(\frac{l_p{}^4 m_p}{t_p{}^3}\right)\left(1 - \left(\frac{4\pi}{32}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$c_{1L} = 4\pi\left(\frac{l_p{}^4 m_p}{t_p{}^3}\right)\left(1 - \left(\frac{4\pi}{32}\right)^2\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\gamma_{\text{he}}{}' = \frac{C_{CFP}{}^2}{\sqrt{7}}\left(\frac{q_p}{m_{\text{he}}}\right)\left(1 - C_d\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\dot{\gamma}_{\text{he}}{}' = \frac{C_{CFP}{}^2}{2\pi\sqrt{7}\ \text{MHz second}}\left(\frac{q_p}{m_{\text{he}}}\right)\left(1 - C_d\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\mu_{\text{he}} = -\frac{C_{CFP}{}^2}{2\sqrt{7}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_{\text{he}}}\right)\left(1 + 5\left(\frac{16}{D_d}\right)\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\mu_{\text{he}}{}' = -\frac{C_{CFP}{}^2}{2\sqrt{7}}\left(\frac{l_p{}^2 q_p m_p}{t_p m_{\text{he}}}\right)\left(1 + \frac{1}{7}\left(\frac{8}{4\pi}\right)\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\frac{\mu_{\text{he}}{}'}{\mu_+} = -\frac{C_{CFP}{}^2}{2\sqrt{7}}\,\frac{S}{ж_1{}^2 ж_2{}^2}\left(\frac{m_+}{m_{\text{he}}}\right)\left(1 - \sqrt{12}\, L^2\left(ж_1{}^2 - ж_2{}^2\right)\boxtimes\right)$$
$$\frac{\mu_{\text{he}}{}'}{\mu_+{}'} = -\frac{C_{CFP}{}^2}{2\sqrt{7}}\,\frac{S}{ж_1{}^2 ж_2{}^2}\left(\frac{m_+}{m_{\text{he}}}\right)\left(1 + \frac{\pi}{\sqrt{12}}\, L^4\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\frac{\mu_e}{\mu_{\text{he}}{}'} = \frac{2\sqrt{7}}{C_{CFP}{}^2}\,\frac{ж_1{}^2 ж_2{}^2}{P_{up}}\left(\frac{m_{\text{he}}}{m_e}\right)\left(1 + \frac{2e^\gamma}{\sqrt{12}}\, L^4\left(ж_1{}^2 + ж_2{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} \vdots E_h = \sqrt{\frac{18}{32}\,\frac{P_{up}}{ж_1{}^{14}}}\left(1 + \frac{1}{ж_3 ж_4}\left(\frac{2}{C_{R1}}\right)\left(ж_1{}^2 - ж_2{}^2\right)\boxtimes\right)$$
$$E_h \vdots \mathrm{A}_{\text{mass}} = \sqrt{\frac{32}{18}\,\frac{ж_1{}^{14}}{P_{up}}}\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{2}{C_{R1}}\right)\left(ж_1{}^2 - ж_2{}^2\right)\boxtimes\right)$$
$$\mathrm{kg} \vdots E_h = \sqrt{\frac{18}{32}\,\frac{P_{up}}{ж_1{}^{14}}}\left(\frac{\text{kilogram}}{\mathrm{A}_{\text{mass}}}\right)\left(1 + \frac{1}{ж_3 ж_4}\left(\frac{2}{C_{R1}}\right)\left(ж_1{}^2 - ж_2{}^2\right)\boxtimes\right)$$
$$E_h \vdots \mathrm{kg} = \sqrt{\frac{32}{18}\,\frac{ж_1{}^{14}}{P_{up}}}\left(\mathrm{A}_{\text{mass}}\right)\left(1 - \frac{1}{ж_3 ж_4}\left(\frac{2}{C_{R1}}\right)\left(ж_1{}^2 - ж_2{}^2\right)\boxtimes\right)$$
$$V_m(Si) = \frac{6}{e^\gamma}\left(\frac{2\pi}{32\,s}\right)^3\frac{1}{ж_1{}^{10}}\left(\frac{l_p{}^3 m_p{}^2}{q_p m_e{}^3}\right)\left(1 - \frac{32\pi}{ж_1 ж_2}\left(\frac{2\pi}{14}\right)\left(ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$F = \frac{6}{e^\gamma}\, ж_1{}^3\left(\frac{1}{m_p}\right)\left(1 + \frac{\omega_2}{ж_1 ж_2}\left(\frac{14}{2\pi}\right)\left(ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_p = \frac{6}{e^\gamma}\,\frac{2\pi\, ж_1{}^2}{\text{second}^2}\left(\frac{l_p{}^2}{t_p q_p}\right)\left(1 - \frac{ж_3 ж_4}{\sqrt{-6}}\left(\frac{14}{4\pi}\right)\left(ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$m_\Delta = \left(m_n - m_+\right)\left(1 - \frac{\sqrt{-1}}{ж_3 ж_4}\left(\frac{4\pi}{14}\right)\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{kg} \vdots \frac{1}{\mathrm{m}} = \frac{\text{kilogram}}{2\pi}\left(\frac{1}{l_p m_p}\right)\left(1 - \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mass}} \vdots \frac{1}{\mathrm{m}} = \frac{1}{2\pi}\left(\frac{\mathrm{A}_{\text{mass}}}{l_p m_p}\right)\left(1 - \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{kg} = \frac{2\pi}{\text{meter}}\left(l_p m_p\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{A}_{\text{mass}} = \frac{2\pi}{\text{meter}}\left(\frac{l_p m_p}{\mathrm{A}_{\text{mass}}}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_C = 2\pi\left(\frac{l_p m_p}{m_e}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_\mu = 2\pi\left(\frac{l_p m_p}{m_\mu}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_+ = 2\pi\left(\frac{l_p m_p}{m_+}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_n = 2\pi\left(\frac{l_p m_p}{m_n}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_\tau = 2\pi\left(\frac{l_p m_p}{m_\tau}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_{\bar{}} = \left(\frac{l_p m_p}{m_e}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_{\mu_-} = \left(\frac{l_p m_p}{m_\mu}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_{+_-} = \left(\frac{l_p m_p}{m_+}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_{n_-} = \left(\frac{l_p m_p}{m_n}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\lambda_{\tau_-} = \left(\frac{l_p m_p}{m_\tau}\right)\left(1 + \frac{P_{up}\, i}{8}\left(ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$

$$g_{\text{de}} = \frac{2\pi}{8K}\left(1 + \frac{16}{P_{up}}\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\frac{\mu_{\text{de}}}{\mu_N} = \frac{2\pi}{8K}\left(1 + \frac{16}{P_{up}}\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\frac{\mu_{\text{de}}}{\mu_B} = \frac{2\pi}{8K}\left(\frac{m_e}{m_+}\right)\left(1 + \frac{16}{P_{up}}\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$b' = \frac{\nu_{\text{peak}}}{2\pi}\left(\frac{1}{t_p T_p}\right)\left(1 + \frac{18}{16}\left(\frac{2\pi}{s}\right)^2\left(ж_1{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\ddot{k}_B = \frac{1}{2\pi}\left(\frac{1}{t_p T_p}\right)\left(1 + \frac{18}{16}\left(\frac{2\pi}{s}\right)^2\left(ж_1{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{K} \vdots \mathrm{Hz} = \frac{\text{kelvin}}{2\pi}\left(\frac{1}{t_p T_p}\right)\left(1 + \frac{18}{16}\left(\frac{2\pi}{s}\right)^2\left(ж_1{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{Hz} \vdots \mathrm{K} = \frac{2\pi}{\text{second}}\left(t_p T_p\right)\left(1 - \frac{18}{16}\left(\frac{2\pi}{s}\right)^2\left(ж_1{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$b = \frac{2\pi}{\lambda_{\text{peak}}}\left(l_p T_p\right)\left(1 + P^2\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$c_2 = 2\pi\left(l_p T_p\right)\left(1 + P^2\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\frac{1}{\mathrm{m}} \vdots \mathrm{K} = \frac{2\pi}{\text{meter}}\left(l_p T_p\right)\left(1 + P^2\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{K} \vdots \frac{1}{\mathrm{m}} = \frac{\text{kelvin}}{2\pi}\left(\frac{1}{l_p T_p}\right)\left(1 - P^2\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\dot{k}_B = \frac{1}{2\pi}\left(\frac{1}{l_p T_p}\right)\left(1 - P^2\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_e = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_e}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_\mu = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_\mu}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_{\mathrm{A}_{\text{mass}}} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{\mathrm{A}_{\text{mass}}}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_+ = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_+}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_n = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_n}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_\tau = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_\tau}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_{\text{de}} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_{\text{de}}}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_{\text{he}} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_{\text{he}}}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_{\text{tri}} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_{\text{tri}}}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_\alpha = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{m_\alpha}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{M}_{^{12}\mathrm{C}} = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{12\,\mathrm{A}_{\text{mass}}}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$N_A = \frac{6}{e^\gamma}\, ж_1{}^2\left(\frac{1}{q_p m_p}\right)\left(1 + 18\, C_Q\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{V}_{90} = \text{volt}\left(1 + \frac{L_1}{32}\left(ж_1{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{W}_{90} = \text{watt}\left(1 - \frac{j_{0,1}}{32}\left(ж_1{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\sigma_{\text{dp}} = 1 - \left(1 + \frac{3}{18}\left(\frac{1}{S^2}\right)\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\sigma_{\text{tp}} = 1 - \left(1 - \frac{18}{35}\left(\frac{L_1}{S^4}\right)\left(ж_2{}^2 - ж_3{}^2 - ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{edm}} = \frac{1}{ж_1}\left(\frac{l_p q_p m_p}{m_e}\right)\left(1 - \frac{2}{7}\left(\frac{16}{2\pi}\right)\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mag}} = \frac{1}{ж_1{}^2}\left(\frac{l_p{}^2 q_p{}^2 m_p{}^2}{m_e{}^3}\right)\left(1 - \frac{4}{7}\left(\frac{16}{2\pi}\right)\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{eqm}} = \frac{1}{ж_1{}^3}\left(\frac{l_p{}^2 q_p m_p{}^2}{m_e{}^2}\right)\left(1 - \frac{4}{\sqrt{3}}\left(\frac{2\pi}{8}\right)^2\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{mfd}} = ж_1{}^3\left(\frac{m_e{}^2}{t_p q_p m_p}\right)\left(1 + \frac{8}{3}\left(\frac{2\pi}{8}\right)^4\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{ep}} = \frac{1}{ж_1{}^6}\left(\frac{t_p{}^2 q_p{}^2 m_p{}^2}{m_e{}^3}\right)\left(1 + \frac{8}{6}\left(\frac{s}{2\pi}\right)^4\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{\text{ef}\,\nabla} = ж_1{}^7\left(\frac{m_e{}^3}{t_p{}^2 q_p m_p{}^2}\right)\left(1 + \frac{s^2}{35}\left(\frac{16}{2\pi}\right)\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{1^{\text{st}}\,\text{hp}} = \frac{1}{ж_1{}^{11}}\left(\frac{t_p{}^4 q_p{}^3 m_p{}^3}{l_p m_e{}^5}\right)\left(1 - \frac{18}{35}\left(\frac{16}{2\pi}\right)^2\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$
$$\mathrm{A}_{2^{\text{nd}}\,\text{hp}} = \frac{1}{ж_1{}^{16}}\left(\frac{t_p{}^6 q_p{}^4 m_p{}^4}{l_p{}^2 m_e{}^7}\right)\left(1 - \frac{18}{8}\left(\frac{2\pi}{s}\right)^4\left(ж_2{}^2 + ж_3{}^2 + ж_4{}^2\right)\boxtimes\right)$$

## legend

$t_p = 5.39125836832313 \ldots \times 10^{-44}$ s — Planck time
$l_p = 1.61625918175645 \ldots \times 10^{-35}$ m — Planck length
$q_p = 1.87554596713962 \ldots \times 10^{-18}$ C — Planck charge
$T_p = 1.41678698590795 \ldots \times 10^{32}$ K — Planck temperature
$m_p = 2.17642683817579 \ldots \times 10^{-8}$ kg — Planck mass

$\boxtimes = 0.999999199973626 \ldots 10^{-7}$ — hyperbolic inversion boundary
**5** — number of dimensions
**7** — break in scale symmetry

$ж_1 = 0.0854245431533304 \ldots$ — 1st hyperbolic partition constant
$ж_2 = 3.66756753485501 \ldots$ — 2nd hyperbolic partition constant
$ж_3 = -1.87649603900417 \ldots + 4.06615262615972 \ldots i$ — 3rd hpc
$ж_4 = -1.87649603900417 \ldots - 4.06615262615972 \ldots i$ — 4th hpc
$ж_r = 4.47826244916751 \ldots$ — hyperbolic radius constant
$ж_\theta = 2.00316562310924 \ldots$ — hyperbolic radian constant

$G_{Gi} = 1.01494160640965 \ldots$ — Gieseking's constant
$V_{fe} = 2.02988321281930 \ldots$ — figure eight knot hyperbolic volume
$K = 0.9159655941772190 \ldots$ — Catalan's constant
$C_d = 0.291560904030818 \ldots$ — domino tiling constant
$D_d = 1.79162281206959 \ldots$ — dimer constant
$\omega_1 = 0.764977018528596 \ldots + 1.32497062714087 \ldots i$ — omega_1 constant
$\omega_2 = 1.529954037057192 \ldots$ — omega_2 constant

$i^{i^{\cdot^{\cdot}}} = 0.438282936727032 \ldots + 0.360592471871385 \ldots i$ — power tower of $i$
$i = (-1)^{1/2}$ — imaginary unit
$\varphi_i = (-1)^{1/3}$ — imaginary golden ratio
$G_g = 1.15872847301812 \ldots$ — tether length for grazing half the unit circle
$\pi = 3.14159265358979 \ldots$ — Archimedes' constant
$P_{up} = 2.29558714939263 \ldots$ — universal parabolic constant
$s = 5.24411510858423 \ldots$ — arc length of the unit lemniscate
$L = 2.622057554292119 \ldots$ — lemniscate constant
$L_1 = 1.31102877714605 \ldots$ — 1st lemniscate constant
$L_2 = 0.599070117367796 \ldots$ — 2nd lemniscate constant
$C_U = 0.847213084793979 \ldots$ — ubiquitous constant
$G_{Ga} = 0.834626841674073 \ldots$ — Gauss's constant

$C_{CF} = 0.697774657964819 \ldots$ — continued fraction constant
$C_{R1} = 0.998136044598509 \ldots$ — Ramanujan's 1st continued fraction constant
$F_{FF} = 1.22674201072035 \ldots$ — Fibonacci factorial constant
$M = -1.74756459463318 \ldots$ — Madelung constant

$e = 2.71828182845904 \ldots$ — Euler's number
$\gamma = 0.577215664901532 \ldots$ — Euler-Mascheroni constant
$S = 0.822825249678847 \ldots$ — Sierpiński's constant
$j_{0,1} = 2.404825557695772 \ldots$ — 1st root of the Bessel function
$C_{CFP} = 1.19967864025773 \ldots$ — Real fixed point of the hyperbolic cotangent
$D_{Do} = 0.739085133215160 \ldots$ — Dottie number
$L_{LL} = 0.662743419349181 \ldots$ — Laplace limit

$\alpha_F = 2.50290787509589 \ldots$ — alpha Feigenbaum constant
$\delta_F = 4.66920160910299 \ldots$ — delta Feigenbaum constant
$\mathcal{S} = 3.24697960371714 \ldots$ — silver constant
$P = 1.32471795724474 \ldots$ — plastic constant
$C_Q = 0.605443657196732 \ldots$ — QRS constant
$C_{QA} = 2.03816937970215 \ldots$ — associated QRS constant
$C_{HSM} = 0.353236371854995 \ldots$ — Hafner-Sarnak-McCurley constant

$T_0 = 2.73150000000000 \ldots \times 10^2$ kelvin — freezing point of water
$p_0 = 1.00000000000000 \ldots \times 10^5$ pascal — 1 bar
$p_1 = 1.01325003754773 \ldots \times 10^5$ pascal — standard atmospheric pressure
$E_1 = 6.09110229711386 \ldots \times 10^{-24}$ joules — energy associated with $\Delta_{\nu_{Cs}}$
$f_1 = 5.40000000000000 \ldots \times 10^{14}$ Hz — frequency of max light conversion
$\lambda_{\text{peak}} = 4.96511423174427 \ldots$ — peak $\lambda$ of spectral radiance/unit wavelength
$\nu_{\text{peak}} = 2.82143937212207 \ldots$ — peak emission of spectral flux/unit frequency

$\log(x)$ = natural logarithm
$\Gamma(x)$ = gamma function
$\zeta(x)$ = Riemann zeta function
$!n$ = derangement function

lm = lumen = cd · sr
N = newton = m kg/s²
$\mathcal{N}$ = noether = m kg/s
Ω = ohm = m² kg/s C²
Pa = pascal = kg/s² m
S = siemens = s C²/m² kg
T = tesla = kg/s C
V = volt = m² kg/s² C
W = watt = m²kg/s³
Wb = weber = m² kg/s C

**units**

s = second
m = meter
C = coulomb
K = kelvin
kg = kilogram

**composite units**

A = ampere = C/s
F = farad = s² C²/m² kg
H = henry = m² kg/C²
Hz = hertz = 1/s
J = joule = m² kg/s²

**derived units**

MHz = megahertz = $1.00000000000000 \ldots \times 10^6$ 1/s
fm = femtometer = $1.00000000000000 \ldots \times 10^{-15}$ m
eV = electron-volt = $1.00000000000000 \ldots$ eV
MeV = megaelectron-volt = $1.00000000000000 \ldots \times 10^6$ eV
GeV = gigaelectron-volt = $1.00000000000000 \ldots \times 10^9$ eV

$$\frac{r_n}{r_e} = \left(\frac{4\pi}{\Gamma(5)}\right)^2 \qquad \frac{r_n}{r_+} = K \qquad \frac{r_n}{r_\alpha} = \sqrt{\frac{C_U}{4}} \qquad \frac{r_e}{r_{\text{de}}} = \sqrt{\frac{2\pi}{\log(36)}} \qquad E_p = \frac{l_p{}^2 m_p}{t_p{}^2}$$

**external transform space**

$$\mathrm{V}_{24} = \left(\frac{4\pi}{5!}\right)\left(\frac{4\pi}{!5}\right)\left(\frac{4\pi}{35}\right)^1\left(\frac{4\pi}{18}\right)^2\left(\frac{4\pi}{32}\right)^3\left(\frac{4\pi}{8}\right)^4$$

**figure eight knot hyperbolic volume**

$$\mathrm{V}_{fe} = i\left[Li_2\left(\frac{1}{\varphi_i}\right) - Li_2\left(\varphi_i\right)\right]$$

**binomial constructor**

$$\mathrm{A}_{\text{ext}}\,\mathrm{B}_{\text{ext}}\left(1 + \mathrm{A}_{\text{int}}\boxtimes\right) \qquad \boxtimes = \frac{l_p m_p}{q_p{}^2} \times \frac{\text{coulomb}^2}{\text{meter kilogram}}$$

**hyperbolic partition equation**

$$\frac{1}{x} + x + \frac{x^3}{2\pi} = \left(i^i\right)^{-\frac{4\pi}{8}} - \frac{m_p}{\text{kilogram}}$$
$$ж_1ж_2ж_3ж_4 = 2\pi$$
$$ж_1 + ж_2 + ж_3 + ж_4 = 0$$
$$ж_1{}^2 + ж_2{}^2 + ж_3{}^2 + ж_4{}^2 = -4\pi$$

$\alpha = ж_1{}^2$
$e = ж_1 q_p$


Physics Monastery
Thad Roberts
Thad Roberts
December-8-2025

with:
Angela Arvizu
Vanessa Moon
Matthew Fox
David Heggli
Eschelon Azha
Avi Rubin
Wayne Eskridge
Dan Whitaker
Kevin Sirios
Mike Ritter
Jerry Gardner
Shauna Montgomery


WolframAlpha
NIST/CODATA