

Global Emissions Timeseries

This manuscript ([permalink](#)) was automatically generated from [openclimatedata/global-emissions@e1c2171](#) on June 1, 2018.

Authors

- Robert Gieseke

 [XXXX-XXXX-XXXX-XXXX](#) ·  [rgieseke](#) ·  [openclimatedata](#)

Potsdam Institute for Climate Impact Research

Unchecked, unreviewed, work-in-progress

Abstract

Introduction

Easily updateable emissions input data for simple climate models like MAGICC6[1], Hector [2], or FAIR[3].

Global-Only.

Inputs

CO₂

Fossil Fuel Industrial

The main contributor to global emissions are fossil fuel and industrial emissions which are available from multiple sources. They are usually based on energy statistics and cement production data. The longest time series from 1750 to 2017 (as a projection for the last year) is available in the Global Carbon Budget 2017 [4], mainly based on data from CDIAC [5] and data submitted to the UNFCCC. PRIMAP-hist [6] in its version 1.2 [7] has data from 1850 to 2015 and is based on the same and other sources as the Global Carbon Budget, but does not include bunker emissions (international aviation and shipping). The EDGAR v4.3.2_FT2016 dataset [8] (in review) covers the years 1970 to 2016.

Figure 1 shows the sources together from 1990.




Figure 1: Global Fossil Fuel and Industrial CO₂ Emissions from various datasets ([High Resolution](#))

Land-Use

Methane (CH₄)

PRIMAP-hist from 1850 to 2015 and RCP data [\[9\]](#) until 2005.




Figure 2: Global Methane (CH₄) Emissions ([High Resolution](#))

N₂O

PRIMAP-hist from 1850 to 2015 and RCP data [9] until 2005.




Figure 3: Global N₂O Emissions ([High Resolution](#))

SO_x

CO

CEDS [[10](#)]

Figure 4: Global CO Emissions (High Resolution)

Figure 4: Global CO Emissions ([High Resolution](#))

NMVOC

CEDS [[10](#)]




Figure 5: Global NMVOC Emissions ([High Resolution](#))

NO_x

CEDS [[10](#)]




Figure 6: Global NOx Emissions ([High Resolution](#))

BC

CEDS [[10](#)]




Figure 7: Global BC Emissions ([High Resolution](#))

OC

CEDS [[10](#)]




Figure 8: Global OC Emissions ([High Resolution](#))

NH₃

Halocarbons

- CF₄
- C₂F₆
- C₆F₁₄
- HFC23
- HFC32
- HFC43-10
- HFC125
- HFC134a
- HFC143a
- HFC227ea
- HFC245fa
- SF₆

References

1. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration

M. Meinshausen, S. C. B. Raper, T. M. L. Wigley

Atmospheric Chemistry and Physics (2011-02-16) <https://doi.org/10.5194/acp-11-1417-2011>

2. A simple object-oriented and open-source model for scientific and policy analyses of the global climate system – Hector v1.0

C. A. Hartin, P. Patel, A. Schwarber, R. P. Link, B. P. Bond-Lamberty

Geoscientific Model Development (2015-04-01) <https://doi.org/10.5194/gmd-8-939-2015>

3. FAIR v1.1: A simple emissions-based impulse response and carbon cycle model

Christopher J. Smith, Piers M. Forster, Myles Allen, Nicholas Leach, Richard J. Millar, Giovanni A. Passerello, Leighton A. Regayre

Geoscientific Model Development Discussions (2017-12-07) <https://doi.org/10.5194/gmd-2017-266>

4. Global Carbon Budget 2017

Corinne Le Quéré, Robbie M. Andrew, Pierre Friedlingstein, Stephen Sitch, Julia Pongratz, Andrew C. Manning, Jan Ivar Korsbakken, Glen P. Peters, Josep G. Canadell, Robert B. Jackson, ... Dan Zhu

Earth System Science Data (2018-03-12) <https://doi.org/10.5194/essd-10-405-2018>

5. Global, Regional, and National Fossil-Fuel CO₂ Emissions (1751 - 2014) (V. 2017)

T. Boden, R. Andres, G. Marland

Carbon Dioxide Information Analysis Center (CDIAC), Oak Ridge National Laboratory (ORNL), Oak Ridge, TN (United States) (2017) https://doi.org/10.3334/cdiac/00001_v2017

6. The PRIMAP-hist national historical emissions time series

Johannes Gütschow, M. Louise Jeffery, Robert Gieseke, Ronja Gebel, David Stevens, Mario Krapp, Marcia Rocha

Earth System Science Data (2016-11-09) <https://doi.org/10.5194/essd-8-571-2016>

7. The PRIMAP-hist national historical emissions time series (1850-2015)

Johannes Gütschow, Louise Jeffery, Robert Gieseke, Ronja Gebel

GFZ Data Services (2018) <https://doi.org/10.5880/pik.2018.003>

8. EDGAR v4.3.2 Global Atlas of the three major Greenhouse Gas Emissions for the period 1970–2012

Greet Janssens-Maenhout, Monica Crippa, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, Frank Dentener, Peter Bergamaschi, Valerio Pagliari, Jos G. J. Olivier, Jeroen A. H. W. Peters, ...

A. M. Roxana Petrescu

Earth System Science Data Discussions (2017-08-28) <https://doi.org/10.5194/essd-2017-79>

9. The RCP greenhouse gas concentrations and their extensions from 1765 to 2300

Malte Meinshausen, S. J. Smith, K. Calvin, J. S. Daniel, M. L. T. Kainuma, J-F. Lamarque, K. Matsumoto, S. A. Montzka, S. C. B. Raper, K. Riahi, ... D.P. P. van Vuuren

Climatic Change (2011-08-09) <https://doi.org/10.1007/s10584-011-0156-z>

10. Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS)

Rachel M. Hoesly, Steven J. Smith, Leyang Feng, Zbigniew Klimont, Greet Janssens-Maenhout, Tyler Pitkanen, Jonathan J. Seibert, Linh Vu, Robert J. Andres, Ryan M. Bolt, ... Qiang Zhang

Geoscientific Model Development (2018-01-29) <https://doi.org/10.5194/gmd-11-369-2018>