

Supporting Information for “Changes in damaging hail in major Australian cities with global warming”

Timothy H. Raupach^{1,2,3}, Joanna Aldridge^{4,5}

¹UNSW Institute for Climate Risk and Response, UNSW Sydney, New South Wales, Australia

²UNSW Climate Change Research Centre, UNSW Sydney, New South Wales, Australia

³ARC Centre of Excellence for Climate Extremes, Sydney, New South Wales, Australia

⁴School of Geosciences, University of Sydney, Sydney, New South Wales, Australia

⁵QBE Australia, Sydney, New South Wales, Australia

Contents of this file

1. Figures S1 to S8.
2. Tables S1 to S2.

Corresponding author: T. H. Raupach, UNSW Sydney Climate Change Research Centre, Mathews Building Level 4, UNSW Sydney, New South Wales 2052, Australia (timothy.h.raupach@gmail.com)

September 16, 2024, 3:52pm

References

- Hong, S.-Y., Noh, Y., & Dudhia, J. (2006). A new vertical diffusion package with an explicit treatment of entrainment processes. *Mon Weather Rev*, *134*(9), 2318 - 2341. doi: 10.1175/MWR3199.1
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., & Collins, W. D. (2008). Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models. *J Geophys Res-Atmos*, *113*(D13). doi: 10.1029/2008JD009944
- Jiménez, P. A., Dudhia, J., González-Rouco, J. F., Navarro, J., Montávez, J. P., & García-Bustamante, E. (2012). A revised scheme for the WRF surface layer formulation. *Mon Weather Rev*, *140*(3), 898 - 918. doi: 10.1175/MWR-D-11-00056.1
- Milbrandt, J. A., Morrison, H., II, D. T. D., & Pukert, M. (2021). A triple-moment representation of ice in the predicted particle properties (P3) microphysics scheme. *J Atmos Sci*, *78*(2), 439 - 458. doi: 10.1175/JAS-D-20-0084.1
- Niu, G.-Y., Yang, Z.-L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., ... Xia, Y. (2011). The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements. *J Geophys Res-Atmos*, *116*(D12). doi: 10.1029/2010JD015139
- Zhang, C., & Wang, Y. (2017). Projected future changes of tropical cyclone activity over the Western North and South Pacific in a 20-km-mesh regional climate model. *J Climate*, *30*(15), 5923 - 5941. doi: 10.1175/JCLI-D-16-0597.1




Figure S1. Approximate extents of the model domains on a map of Australia. The coarse-resolution domains are in yellow, medium-resolution domains in dark green, and fine-resolution domains in blue. The solid and dotted lines group the two sets of nested domains that were calculated together. Approximate city locations (with city extents not shown) are marked with red points for Perth (a), Kalgoorlie (b), Adelaide (c), Melbourne (d), Canberra (e), Sydney (f), and Brisbane (g).




Figure S2. Maximum hail size diameters for the Adelaide, Brisbane, Kalgoorlie, Melbourne, Perth, and Sydney/Canberra domains under the historical and future scenarios, with and without ocean areas removed.




Figure S3. Maximum 10 m wind speed at hail times in historical and future climates, for Adelaide (a, b), Brisbane (c, d), Kalgoorlie (e, f), Melbourne (g, h), Perth (i, j) and Sydney/Canberra (k, l) domains.




Figure S4. Time series of daily maximum hail sizes by domain and epoch. Gaps exist because only the convective season was simulated each year.




Figure S5. As for Figure S4, but for daily maximum 10 m wind collocated with hail.




Figure S6. Quantile-quantile plots for GEV models fitted to daily maximum hail sizes, per domain and epoch.




Figure S7. Quantile-quantile plots for GEV models fitted to daily maximum 10 m wind collocated with hail.




Figure S8. Distributions of p values from KS tests, comparing empirical values GEV models, and historical GEVs to future GEVs, per variable and domain. For each test, the KS test was applied 100 times with 1000 random values drawn from the relevant GEV distribution(s) each time, to obtain a distribution of p values. Bars show medians, box hinges show the interquartile ranges (IQRs), whiskers show the largest (smallest) values no more than $1.5 \times \text{IQR}$ from the upper (lower) hinge, and points show outlier points beyond the whisker ranges. The red horizontal line shows $p = 0.05$; when p values are below this line the null hypothesis that the two samples come from the same distribution can be rejected.

Table S1. Parameterization schemes used in the WRF simulations.

Microphysics	P3-3moment (Milbrandt et al., 2021)
Cumulus (medium and coarse nests only)	New Tiedtke (Zhang & Wang, 2017)
Long wave and shortwave radiation	RRTMG (Iacono et al., 2008)
Planetary boundary layer	YSU (Hong et al., 2006)
Surface layer	Revised MM5 (Jiménez et al., 2012)
Land surface	Noah-MP (Niu et al., 2011)

Table S2. Probability of exceeding hail sizes of 20, 50, or 100 mm, and of coincident wind exceeding 80 and 100 km h⁻¹, per domain and epoch.

Domain	Epoch	Hail probability [%]			Wind probability [%]	
		20 mm	50 mm	100 mm	80 km h ⁻¹	100 km h ⁻¹
Adelaide	Historical	49.99	10.49	3.45	1.70	0.00
	Future	48.70	9.23	2.62	5.41	1.00
Brisbane	Historical	68.34	12.65	2.48	1.64	0.02
	Future	72.17	13.12	2.30	2.06	0.01
Kalgoorlie	Historical	43.85	6.22	1.51	9.03	0.00
	Future	46.79	8.89	2.57	13.02	0.23
Melbourne	Historical	45.82	4.29	0.65	1.26	0.05
	Future	60.50	12.77	3.66	0.59	0.00
Perth	Historical	45.57	6.02	1.11	1.84	0.00
	Future	49.06	6.98	1.35	0.89	0.00
Sydney/Canberra	Historical	49.26	6.01	1.04	0.57	0.01
	Future	61.83	9.93	1.78	0.24	0.00