Supplemental Material for:

Evaluation of predicted Medfly (*Ceratitis capitata*) quarantine length in the United States utilizing degree-day and agent-based models

Travis C. Collier 1,2 and Nicholas C. Manoukis 1,3

¹Daniel K. Inouye US Pacific Basin Agricultural Research Center (PBARC), United States Department of Agriculture, Agricultural Research Service, Hilo, Hawaii, 96720, USA

Contents

1	"Supernorm" figures	2
	1.1 Daily normal of hourly temperatures	2
	1.2 Degree day based PQL supernorm	3
	1.3 MED-FOES PQL based supernorm	4
2	Difference in PQL at nearby sites (LA basin example)	5
3	Historical quarantines	ť

²corresponding author; email: Travis.Collier@ARS.USDA.gov

³email: Nicholas.Manoukis@ARS.USDA.gov

"Supernorm" figures

Daily normal of hourly temperatures




Figure 1. Hourly temperature data aggregated by day of year.

Degree day based PQL supernorm




Figure 2. Single sine degree day based predicted quarantine lengths aggregated by day of year.

MED-FOES PQL based supernorm




Figure 3. MED-FOES based predicted quarantine lengths aggregated by day of year.

Difference in PQL at nearby sites (LA basin example)




Figure 4. The difference in PQL values for the same start date (including year) between sites in the LA basin. For each date over the range of available data (1973 through 2015 for LAX-BUR and RIV-BUR, 1950 through 2015 for LAX-RIV), PQL values for each site are computed and the differences taken. The resulting differences are then aggregated by day-of-year. Lines show the median while the shaded region is the 25% to 75% quantile range for each day-of-year aggregation of differences.

Historical quarantines

Table 1. Historical quarantines in California.