

Evaluation of predicted Medfly (*Ceratitis capitata*) quarantine length in the United States utilizing degree-day and agent-based models

Travis C. Collier^{1,2} and Nicholas C. Manoukis^{1,3}

¹Daniel K. Inouye US Pacific Basin Agricultural Research Center (PBARC), United States Department of Agriculture, Agricultural Research Service, Hilo, Hawaii, 96720, USA

²corresponding author; email: Travis.Collier@ARS.USDA.gov

³email: Nicholas.Manoukis@ARS.USDA.gov

Abstract Invasions by pest insects pose a significant threat to agriculture worldwide. In the case of *Ceratitis capitata* incursions on the US mainland, where it is not officially established, repeated detections are followed by quarantines to eliminate the invading population. However, it is notoriously difficult to set quarantine duration because the non-detection of the pest does not necessarily mean the pest is extirpated. Most programs determine quarantine lengths by calculating the amount of time required for 3 generations to elapse under a thermal unit accumulation development model ("degree day"). A newer approach is to use an Agent-Based Simulation (ABS) to explicitly simulate population demographics and extirpation. Here, the predicted quarantine length for 11 sites in the continental United States were evaluated. Results indicate that strong seasonality in quarantine length, with longer predictions in the second half of the year compared with the first; this pattern was more extreme in the degree day predictions compared with the ABS. Geographically, quarantine lengths increased with latitude, though again this feature was less pronounced under the ABS. Variation in quarantine lengths for particular times and places was significantly higher for degree day versus ABS, and generally peaked in the middle of the year for the former and second half of the year for the latter. Analysis of 34 *C. capitata* quarantines from 1975 to 2017 in California shows that all but two were declared in the second half of the year, when degree day quarantine lengths are longest and have the highest uncertainty. For this set of hypothetical quarantines, the ABS produced significantly shorter quarantines than degree day calculations. Overall, ABS quarantine lengths were more consistent than degree day predictions, avoided unrealistically long values seen in the northernmost site studied (San Francisco), and realistically modeled the effects of rare events such as cold snaps.

Keywords

biosecurity, Mediterranean fruit fly, eradication, invasive pest, agriculture

Introduction

Invasions by insects, pathogens and pests are increasingly a defining challenge of the 21st century, facilitated by global connectivity, climatic shifts, and other factors [1, 2], with a particularly severe impact on agriculture[3]. Invasions by insects that do not become established have a lower public profile than those that are “successful” from the point of view of the insect. However, when the invading species is of environmental, human health, or economic concern there is a greater chance that cases of invasion followed by extirpation would be detected and studied [4]. Eradication of such insects can be desirable and feasible [5] depending on several factors.

One factor is if the new environment is only marginally or seasonally suitable to the invading insect, facilitating its eradication. Another is when the high cost of allowing establishment leads to extensive efforts for eradication. The invasion of the malaria mosquito species *Anopheles gambiae* into Northeastern Brazil in the 1930's [6] is one example of an invasive insect that was successfully eradicated primarily due to the second of these reasons [7, 8]. In the case of *An. gambiae* there have been no reports of reinvasion, but there are examples of insects that recurrently invade areas outside their native range and are recurrently extirpated within relatively few generations. The Gypsy moth *Lymantria dispar* in Canada [9] is one such species. Arguably, another example is the screw-worm *Cochlyomyia hominivorax* along the current northernmost edge of its range in Panama[10] and more recently in Florida[11].

One of the most important instances of repeated invasion and extirpation by an economically important insect pest is that of the Mediterranean fruit fly *Ceratitidis capitata* (Wiedemann), aka. “Medfly”, in California. The last four decades have seen a repeated pattern of invasion, detection, and response interspersed by periods of no detections [12, 13]. While it has been suggested that this pattern is the result of cryptic establishment of Medfly in California[14], others have argued that Medfly in California is an example of a “metainvasion”, consisting of multiple sequential or overlapping introductions [15]. Still other researchers have maintained that Medfly is repeatedly eradicated there [16], or that there are different situations in different regions of the state [17, 18]. Medfly is occasionally found in other parts of the mainland US such as Florida[19], and in other countries or areas that are considered free of the pest including Australia, Mexico and Chile[20].

The response plan to Medfly in California, and the other “free” regions mentioned above, is extensive and costly, including a quarantine when detections exceed an established threshold [21]. A practical and important problem is how long to maintain the countermeasures and quarantine after flies are no longer detected. Predicting the likely duration of required quarantines would help with management decision-making and planning, including potential cost savings by having sufficient but not excessive resources available.

Currently most programs determine quarantine lengths

by calculating the amount of time required for a given number of generations (usually three) to elapse under a thermal unit accumulation (“degree day”) physiological development model. Degree day based quarantine lengths have been codified in some legal regulations, including United States Federal (@TCC make ref:Title 7 Subtitle B Chapter III Part 301.32-10 Treatments), California[22], and Florida[?]. However, the procedure prescribed only defines when the end of a quarantine period has been reached after the fact. For planning and resource allocation, policy makers and managers typically attempt to predict the quarantine lengths by using normal temperatures for forward projection [?]. Although it frequently works fairly well, this approach is mathematically flawed and also provides no indication as to the variance or uncertainty of those predictions.

Recently, another approach to determining effective quarantine durations against Medfly via Agent-Based Simulations (ABS)[23] was introduced. The MED-FOES system simulates a population of individual Medflies under inundative sterile insect technique (SIT) and other control, explicitly modeling extirpation as opposed to the degree day approach which only determines the time for a specific number of generations to elapse to estimate quarantine duration. MED-FOES also allows for the sampling of parameter space (temperature dependent mortality for each stage, fecundity, etc.) producing a distribution of possible outcomes.

In this paper, predicted quarantine length (PQL for short) for 11 sites in the continental United States were analyzed (figure 1 and table 1) based on both the standard thermal accumulation degree day method[24] as well as the MED-FOES ABS[25]. Seasonal variation dominates quarantine duration, so we aggregated the PQL values for each day of the year (Jan. 1, Jan. 2, ect.) across a large number of years (65 for most locations) to produce normals. This approach enables comparison of the standard degree day method to the ABS, but more importantly provides insight into seasonal and spatial variations, prediction uncertainties, and model reliability.



Figure 1. Location of sites analyzed.

Methods

Sites and Temperature Data

Hourly air temperature data for 11 sites was downloaded from NOAA's Integrated Surface Database (ISD) dataset[26, 27]. The airport sites shown in figure 1 were chosen for their biological relevance and availability of high quality hourly data over a long time frame.

Table 1. Weather station (NOAA ISD) sites used.

Callsign	Station Name	State	Latitude	Longitude	Elevation	Start year
KSFO	SAN FRANCISCO INTERNATIONAL A	CA	+37.620	-122.365	2.4	1950
KFAT	FRESNO YOSEMITE INTERNATIONAL	CA	+36.780	-119.719	101.5	1950
KBUR	BURBANK-GLENDALE-PASA ARPT	CA	+34.201	-118.358	236.2	1973
KLAX	LOS ANGELES INTERNATIONAL AIR	CA	+33.938	-118.389	29.6	1950
KRIV	MARCH AIR RESERVE BASE	CA	+33.900	-117.250	468.2	1950
KSAN	SAN DIEGO INTERNATIONAL AIRPO	CA	+32.734	-117.183	4.6	1950
KJAX	JACKSONVILLE INTERNATIONAL A	FL	+30.495	-81.694	7.9	1950
KIAH	G BUSH INTERCONTINENTAL AP/HO	TX	+29.980	-95.360	29.0	1970
KMCO	ORLANDO INTERNATIONAL AIRPORT	FL	+28.434	-81.325	27.4	1973
KTPA	TAMPA INTERNATIONAL AIRPORT	FL	+27.962	-82.540	5.8	1950
KMIA	MIAMI INTERNATIONAL AIRPORT	FL	+25.791	-80.316	8.8	1950

Sites are referred to here by the last three letters of the callsign shown in table 1. For 8 sites (SFO, FAT, LAX, RIV, SAN, JAX, TPA, and MIA), temperature data starting on 1950-01-01 was used. The 3 other sites contained large (> 14 days) gaps or other problems in the early years of their data, so data starting on 1970-01-01 for IAH and 1973-01-01 for BUR and MCO was used. For all sites, temperature data from the start date through 2017-05-15 was used to generate PQLs for dates ranging from the start date to 2016-01-01.

Data was fetched and parsed using the `Fetching and parsing ISH.ipynb` program. Records for the same station callsign were merged, since identification, format, and precise location of stations has changed over the years. The data was then cleaned using the `Cleaning temperatures.ipynb` program by removing outliers, identifying large gaps (> 3 hours), resampling to every hour on the hour using linear interpolation, and filling the large gaps using day-over-day linear interpolation (interpolating using values for the same hour of day from previous and following days). The processing programs and resulting temperature datasets are provided in the Supplemental Materials.

Degree-Day Calculation

Degree-days were computed by the single-sine method[24] using a base development temperature of 12.39°C (53.3°F) and 345.56 degree-days Celsius (DDc; 622 DDf) per generation following the standard required by California Department of Food and Agriculture regulation 3406(b)[28, 22]. Since we have hourly temperature data, we also calculated degree-days by simple summation for comparison[29]. For each date, the number of days required for 3 generations of degree-day based life cycles was computed. These calculations are implemented in `Temperature functions.ipynb` in the Supplemental Materials.

Agent-based Simulations: MED-FOES

MED-FOES[25, 23] is an agent-based simulation explicitly modeling the eradication of a population of Medflies under inundative sterile male releases (aka: sterile insect technique or SIT) and other interventions such as increased trapping and foliar sprays. A MED-FOES simulation models a single non-spatial population starting from a given age distribution and tracks the number of individuals through the time the population experiences extirpation when the last fly (Agent) dies. The simulation is parameterized on the initial population, additional mortality induced by control efforts, the effectiveness of SIT, and a large number of biological parameters for which ranges are known from the literature, including temperature-dependent development and mortality. The simulations were performed using the same hourly timeseries of temperature values used for degree-day calculations and was updated in hourly time steps.

Due to the fact that ranges can be estimated for many of the parameters, 2500 individual MED-FOES simulations were run for each start date at each site evenly sampling different regions of parameter-space via Latin Hypercube Sampling[30] procedure. This set of simulations, encompassing a range of possible extirpation outcomes, is referred to as a ‘run’. Run data are translated into PQL by the number of days from the start date required for 95% of the simulations in a run to be eradicated, also referred to as pe_{95} .

Varying the start date for different simulations was achieved by simply starting at different points in the input temperature file; for this study a run starting every 7 days over the range of dates available for each site. Each set of runs for a single site over a range of starting dates is referred to as a ‘runset’. All runsets were conducted with the same input parameters aside from temperature. Initial population numbers were chosen as a “standard outbreak” based on seven real outbreaks modeled previously[23].

MED-FOES version 0.6.2 was run under OGS/Grid Engine 2011.11 on a CentOS 6.6 HPC cluster. The MED-FOES code, configuration files, and helper scripts are provided




Figure 2. Summary of normal quarantine length predictions for each site. Year range of input temperature data used is inclusive. All panels have identical limits except SFO quarantine.

in the Supplemental Materials. Overall, we created 11 runsets (one for each site), each containing runs starting every 7 days over the input temperature data range for that site, where each run contained 2500 individual simu-

lations sampling different regions of biologically plausible parameter space.

The MED-FOES data is summarized here by the number of days from the start date required for 95% of the simu-

lations in a run to be eradicated, referred to as the MED-FOES PQL (or pe95 within the programs/scripts).

Statistical analysis

The main results reported here are ‘normals’ in a meteorological sense of term, but without the typical running mean smoothing which would complicate interpretation of the results. For a variable of interest (eg. temperature or PQL), all values for the same calendar day irrespective of year (eg. 20-July) are aggregated and summary statistics such as mean, minimum, maximum, and standard deviation are computed for each aggregation. `Temperature functions.ipynb` contains the code used to perform normal calculations, and the code generating these figures is `Summary Figures.ipynb`. The results reported here are the normals of PQL computed using the full temperature time series as opposed to computing PQL from the normal of the temperature time-series. While the latter is fairly common practice, it is not mathematically proper since, as with means, the normal of a function of X is not generally equal to the function applied to the normal of X . Additionally, by computing the normals of the predicted quarantine durations, we can investigate properties of the distribution of values as shown in figures 4 and 5 and the “supernorm” supplemental figures S??, S??, and S??.

Results

Figure 2 shows the mean of the normal PQL based on 3 generation degree day accumulation and MED-FOES 95% extirpation along with the minimum and maximum of the normals for temperatures. Figures 4 and 5 show the standard deviations (σ) of the normals for the degree day and MED-FOES based PQL.

There is significant variation in PQL across both time and location. The temporal variation in PQL is dominated by a yearly cycle, characterized by the normal values shown in figure 2. Table 2 shows the percentage of variance in quarantine length predictions, captured by the mean of the normal yearly cycle (aka. R^2) for each site. At all but one site, greater than 75% of the variance in both degree day and MED-FOES based PQL is accounted for by the mean normal, and the majority exceed 90%. SFO is an exception to this common trend, with the mean normal accounting for only 9.1% of the variation in degree day based PQL and 28.0% of the MED-FOES based PQL. This is also shown in the respective ‘supernorm’ supplemental figures S??, and S??.

Seasonal dependence

Seasonal variation, evidenced by the general shape of the curves shown in figure 2, is doubtless familiar to anyone engaged in Medfly pest management. Outbreaks starting in the late summer, autumn, or early winter will extend through relatively cold periods, when thermal dependent development will be slow and therefore extend the duration of quarantine required for 3 generations of degree days to accumulate (referred to as DD PQL hereafter).

Table 2. Percentage of PQL variance captured by the mean of the normal. DD PQL is the 3 generation single sine degree day based prediction, and ABS PQL is the MED-FOES agent-based simulation predictions.

Site	R^2	
	DD PQL	ABS PQL
SFO	9.12%	28.01%
FAT	93.93%	75.68%
BUR	90.71%	90.88%
LAX	80.17%	83.07%
RIV	92.23%	81.89%
SAN	80.99%	80.91%
JAX	96.45%	94.78%
IAH	95.10%	91.80%
MCO	94.62%	95.77%
TPA	91.91%	94.40%
MIA	88.42%	92.00%

Similarly, outbreaks starting in the spring or early summer often lead to short quarantines due to the relatively high temperatures.

This familiar pattern is also seen in the MED-FOES ABS PQLs despite it being quite different in nature from simple degree day accumulation. However, the MED-FOES predictions (pe95) show a smaller seasonal swing. pe95 generally produces a smaller overall range of PQLs, with longer quarantines than DD PQL for spring and early summer outbreaks, and shorter quarantines for late summer through early winter in almost all cases.

A particular feature of interest, shown most dramatically at FAT in figure 2, is that ABS PQL often flattens out or even dips for quarantines starting in the late autumn or early winter. This can be due to relatively rare and brief cold-snaps, normally lasting only a few hours, which increase mortality. Since DD PQL does not account for mortality, it misses the effect of cold-snaps entirely. This cold-snap effect is most clearly seen at more northern and inland sites where cold-snaps are more likely: particularly FAT and RIV, but also BUR, LAX, JAX, and IAH.

Geographic dependence

PQL generally shows a positive correlation with latitude[?]. Sites are ordered by latitude in the figures and tables. As seen in figure 2, higher latitude sites tend to have longer PQL as well as larger seasonal swings for both degree day and MED-FOES based predictions.

Figure 3 shows the relationship between PQL and latitude. An ordinary least squares fit to the median PQL at each site shows a significant slope for both DD PQL ($F=14.08$, $p=0.005$) and ABS PQL ($F=10.55$, $p=0.010$), but the degree day based predictions are more sensitive to latitude than MED-FOES (coefficients of 17.39 and 4.78 respectively). Additionally, MED-FOES predictions are more stable for SFO, and to a lesser extent FAT, where the




Figure 3. Predicted quarantine length dependence on latitude. For each site, the mean, median, and inter-quartile range are shown (similar to a boxplot). An ordinary least-squares linear fit to the median values is shown by the green lines. The left pane is for single sine degree day predictions, and MED-FOES based predictions are in the right pane.

degree day model for Medfly produced PQLs that appear either unrealistically long (SFO) or are subject to rapid and extreme seasonal variation in the mid year (FAT).

In addition to the variation associated with latitude, large differences in PQLs computed for the same start date can exist between even relatively nearby sites. For example, the differences in both degree day and MED-FOES PQLs for the three sites in the Los Angeles region (LAX, BUR, RIV) (shown in the supplemental figure ??) show a strong seasonal component with a spike in July and/or August. The difference in DD PQL between LAX and BUR is normally about a month (overall median=35 days; overall 25% & 75% quantiles are 28 & 45 days), but the median difference of the normal exceeds 75 days in August with some PQL differences up to 142 days. Differences in MED-FOES PQLs are more seasonally stable, with the LAX minus BUR (MEAN?) difference not exceeding 42 days at its maximum.

Variance and uncertainty

Figures 4 and 5 report the standard deviation (σ) of the normal for DD PQL and MED-FOES PQL respectively. These indicate the year to year variability of the PQL for outbreaks starting at a given time of the year and can be used to gauge the uncertainty of quarantine length predictions relative to the actual quarantine length which will be required. Similar information is represented by the inter-quartile ranges shown in figure 3 and the 'supernorm' supplemental figures. Those 'supernorm' figures also show that the underlying distributions of PQL values are generally not highly skewed, making σ a relatively easy to interpret proxy for uncertainty.

Excluding SFO, the mean normal is a good predictor of DD PQL with σ values below 20 days except for the late summer and early autumn, where variance increases due to quarantines extending through the cold season. FAT and, to a lesser extent, RIV show this increase more dramatically, presumably due to their more arid/inland climates where both daily and seasonal temperature ranges

are larger (also see figure 2). The standard deviation generally decreases with decreasing latitude, together with reduced means. The standard deviation in DD PQL for SFO shows an inversion of the seasonal trend other sites exhibit. This is due to the colder temperatures leading to extremely long DD PQLs, frequently extending across two winter seasons.

The standard deviations of MED-FOES based PQL normals shown in figure 5 are generally about $1/2$ as large as for DD PQL. This indicates that MED-FOES PQL not only shows less dramatic seasonal swings, but is also produces more consistent predictions across years. Values again generally decrease with latitude, but less consistently than DD PQL σ of normals. Also, unlike with the DD PQL, the results for SFO appear consistent with other sites.

A notable feature is that BUR, LAX, and SAN all show an increase in the year to year variation in MED-FOES PQL starting in July and extending through November, while that increase for all other sites starts in July or August but extends to January or February. Additionally, results for FAT show a sharp increase in uncertainty starting in September, fitting with the more arid/inland climate. RIV shows a significant but more gradual increase.

Historical quarantines

34 Medfly quarantines in CA dating from 1975 to early 2017 were analyzed (supplemental table ??). All but two of these quarantines were declared in the latter half of the year (July through December) when DD PQLs are typically relatively long, with 68% (23/34) occurring in September through October (when DD PQLs are longest). August, the month where uncertainty in DD PQL often spikes (see figure 4), accounts for 30% (7/34) of historic quarantines.

DD PQLs and ABS PQLs were computed using the start date for each historic quarantine and the temperature data from the closest site of the 11 analyzed here (see supplemental table ??). For this set of hypothetical quarantines, MED-FOES produced significantly shorter quar-




Figure 4. Variation in quarantine length prediction based on 3 generations of single-sine degree day accumulation.




Figure 5. Variation in quarantine length prediction based on 95% of MED-FOES simulations showing extirpation.

antines (mean=169.7 days, $\sigma=21.8$ days) than simple 3 generation degree day accumulation (mean=234.2 days, $\sigma=79.2$ days) (t-statistic=6.01, $p < 10^{-5}$). Additionally, the variance in the difference between quarantine lengths using a specific date and the mean of the normal PQL for that day of year were smaller for MED-FOES ($\sigma=8.2$ days) than with degree day ($\sigma=25.9$ days) (F-statistic=9.92, $p < 10^{-8}$).

Discussion

The principal contributions of this work can be broken down into three categories: 1) Comparison of PQLs as determined by the degree day and ABS methods. 2) variation in average PQLs across time of year and space, and 3) variation in PQLs within a time of year and location. Consideration of all three of these by program managers, planners and other decision makers is likely to improve quarantines by informing resource allocation ahead of outbreaks, reducing quarantine costs in some cases, and reducing risk from premature quarantine suspension in others. The results presented cover most of the latitudinal range of Medfly suitability within the United States as well as many sites of probable introduction, and will hopefully find use as a general guide.

Extirpation models are extremely difficult to test for accuracy given the impracticality of experimental introductions and the sparse and idiosyncratic nature of historic outbreaks. However, analyzing locations of historic outbreaks suggests that quarantine lengths would generally be more consistent and shorter on average in California if estimated by ABS compared with degree day.

Requiring a fixed number (typically 3) generations of degree days to pass is a "tried and true" method, but not explicitly an extirpation model. It may overestimate required quarantine length through cold weather[?] and may underestimate length when growth conditions are very favorable, which somewhat paradoxically leads to quicker extirpation under SIT (CITE?). However, the simplicity of the degree day calculation is a point in its favor, together with its record of generally avoiding subsequent detections after quarantine[20].

ABS results may be used to inform and modulate responses and treatments which are under the discretion of managers. In situations where DD PQL greatly exceed those from MED-FOES, it is likely that degree day is missing important effects such as cold snaps which may justify shortening quarantine periods. On the other hand, in cases where MED-FOES predicts longer times to extirpa-

tion, it is plausible that the degree day indicated quarantine is short, and eradication treatments and SIT releases should be conducted aggressively.

A few specific results arising from overall comparisons of different locations are worth highlighting. In general, DD PQL for Medfly generated from San Francisco International Airport temperature data are almost certainly too long for the entire year. The ABS PQLs are flatter and seem more realistic at around 200 days for San Francisco compared with the 400-550 days of DD PQLs. For several other California locations (typified by Fresno and Riverside) DD PQLs are in close alignment with those from the ABS for the first half of the year but go significantly longer in the cooler months. For three of four Florida locations analyzed, DD PQLs are significantly shorter than the ABS results (Miami, Tampa, and Orlando). The extent of the difference in those Florida locations is smaller in the later months of the year, but the generality of this pattern suggests that the margin of safety for quarantines as calculated by degree day in those locations may be smaller than expected.

Other systematic comparisons of (PQLs? development?) include .. (PLUG Davis paper here)

As expected, there is significant variation in PQL depending on the location of the outbreak, with the extremes in our study sites represented by Miami and San Francisco. These geographic results could be compared directly to previous efforts to model climatic suitability of different parts of the US for Medfly, by equating longer PQLs with higher climatic suitability. One of the early studies on the subject focused on Medfly found higher climatic suitability in Florida locations (Fort Pierce and Orlando) compared with California sites [31]. Within California, however, those authors found a higher number of suitable months in coastal areas such as Oceanside compared with Riverside and Fresno, roughly paralleling our findings (compare Los Angeles or San Diego with Fresno or Riverside). A more recent analysis of climatic suitability likewise concludes that coastal S. California is the most favorable area of the state for Medfly, but favorability drops inland in the south due to desert conditions. Suitability in central and northern California is limited by cold temperatures and freezes [32].

Seasonal variation within locations revealed WORDS ON SEASONAL HERE [Anna's 2014 paper; refs she cites "It is well documented in regional studies from several areas of the world that Medfly has a highly seasonal pattern to its population dynamics"...: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4222914/>]

An important aspect of PQLs from the ABS is variation within particular times of years and locations. Rare events like cold snaps can increase mortality in the ABS, and thereby lead to shorter PQLs than expected based on historical averages, or DD PQLs. The specificity of the ABS is helpful for determining when quarantines might be safely suspended due to a rare event, something that might not be captured by the degree day model. The degree day model includes only development for generating PQLs, and development is halted at low temperatures, extend-

ing quarantine lengths. The ABS, however, also includes mortality for generating PQLs, which means that low temperatures can significantly reduce estimates.

In California, historically quarantines have most frequently occurred at times of year when degree day based quarantines are drawn out by cold weather and the MED-FOES ABS model predicts significantly shorter durations. Furthermore, 30% of those historic quarantines happened in August where there is a great deal of uncertainty in forward predictions of degree day quarantine durations based on normal values. If we assume those historic CA quarantines are a guide, the ABS model would very likely produced more predictable and shorter quarantine durations for future outbreaks.

The initial quarantine length estimate could be quickly produced based on the distribution of PQL values generated using historical temperatures. This would generate not just a single "typical" value as the current method of projecting using historical average/normal temperatures does. The median "most likely" value may be used for official estimates, while the variance and extremes would provide managers and affected parties additional information vital for planning.

Once the eradication program is underway, weekly simulations via the ABS could indicate the likelihood that extirpation has been achieved. If the threshold 95% extirpation is observed the decision to end quarantine early could be made, or in the case where the ABS has not reached the 95% threshold at the end of the DD PQL additional measures could be considered to reduce the risk of re-detection.

Author contributions

In order to give appropriate credit to each author of an article, the individual contributions of each author to the manuscript should be detailed in this section. We recommend using author initials and then stating briefly how they contributed.

Competing interests

None to declare.

Grant information

This research was funded by USDA-ARS, project number 2040-22430-025-00D and by the Headquarters Research Associate program (TCC).

Acknowledgements

We thank XX and YY for comments on an early draft of this paper, and N. Mullaly (USDA-APHIS) for data on historical outbreaks. This work was supported by the US Department of Agriculture, Agricultural Research Service. Opinions, findings, conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the USDA. USDA is an equal opportunity provider and employer.

References

- [1] Daniel Simberloff, Jean-Louis Martin, Piero Genovesi, Virginie Maris, David A. Wardle, James Aronson, Franck Courchamp, Bella Galil, Emili García-Berthou, Michel Pascal, Petr Pyšek, Ronaldo Sousa, Eric Tabacchi, and Montserrat Vilà. Impacts of biological invasions: what's what and the way forward. *Trends in Ecology & Evolution*, 28(1):58–66, January 2013. ISSN 0169-5347. doi: 10.1016/j.tree.2012.07.013. URL <http://www.sciencedirect.com/science/article/pii/S0169534712001747>.
- [2] David Pimentel, Marcia Pimentel, and Anne Wilson. *Plant, Animal, and Microbe Invasive Species in the United States and World*, pages 315–330. Springer Berlin Heidelberg, Berlin, Heidelberg, 2007. ISBN 978-3-540-36920-2. doi: 10.1007/978-3-540-36920-2_18. URL https://doi.org/10.1007/978-3-540-36920-2_18.
- [3] Dean R Paini, Andy W Sheppard, David C Cook, Paul J De Barro, Susan P Worner, and Matthew B Thomas. Global threat to agriculture from invasive species. *Proceedings of the National Academy of Sciences*, 113(27):7575–7579, 2016.
- [4] Andrew M. Liebhold and Patrick C. Tobin. Population Ecology of Insect Invasions and Their Management*. *Annual Review of Entomology*, 53(1):387–408, January 2008. ISSN 0066-4170, 1545-4487. doi: 10.1146/annurev.ento.52.110405.091401. URL <http://www.annualreviews.org/doi/abs/10.1146/annurev.ento.52.110405.091401>.
- [5] Judith H. Myers, Daniel Simberloff, Armand M. Kuris, and James R. Carey. Eradication revisited: dealing with exotic species. *Trends in Ecology & Evolution*, 15(8):316–320, August 2000. ISSN 0169-5347. doi: 10.1016/S0169-5347(00)01914-5. URL <http://www.sciencedirect.com/science/article/pii/S0169534700019145>.
- [6] F.L. Soper and D.B. Wilson. *Anopheles gambiae in Brazil, 1930 to 1940*. The Rockefeller Foundation, New York, 1943.
- [7] OR Causey, LM Deane, and MP Deane. Ecology of *Anopheles gambiae* in Brazil. *Am. J. Trop. Med.*, 23(1):73–94, 1943.
- [8] Gerry F Killeen, Ulrike Fillinger, Ibrahim Kiche, Louis C Gouagna, and Bart G J Knols. Eradication of *Anopheles gambiae* from Brazil: lessons for malaria control in Africa? *The Lancet Infectious Diseases*, 2(10):618–627, October 2002. ISSN 1473-3099. URL <http://www.ncbi.nlm.nih.gov/pubmed/12383612>.
- [9] David R. Gray. Hitchhikers on trade routes: A phenology model estimates the probabilities of gypsy moth introduction and establishment. *Ecological Applications*, 20(8):2300–2309, 2010. ISSN 1051-0761. doi: 10.1890/09-1540.1. URL <http://www.esajournals.org/doi/abs/10.1890/09-1540.1>.
- [10] A S Robinson, M J B Vreysen, J Hendrichs, and U Feldmann. Enabling technologies to improve area-wide integrated pest management programmes for the control of screwworms. *Medical and Veterinary Entomology*, 23 Suppl 1:1–7, June 2009. ISSN 1365-2915. doi: 10.1111/j.1365-2915.2008.00769.x. URL <http://www.ncbi.nlm.nih.gov/pubmed/19335824>.
- [11] Jason Matthews and Joe N Caudell. In the news spring 2017. *Human–Wildlife Interactions*, 11(1):3, 2017.
- [12] J.R. Carey. Establishment of the Mediterranean fruit fly in California. *Science*, 253(5026):1369, 1991.
- [13] Nikos T. Papadopoulos, Richard E. Plant, and James R. Carey. From trickle to flood: the large-scale, cryptic invasion of California by tropical fruit flies. *Proceedings of the Royal Society B: Biological Sciences*, 280(1768):20131466, October 2013. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2013.1466. URL <http://rspb.royalsocietypublishing.org/content/280/1768/20131466>.
- [14] James R Carey, Nikolas Papadopoulos, and Richard Plant. The 30-year debate on a multi-billion-dollar threat: Tephritid fruit fly establishment in California. *American Entomologist*, 63(2):100–113, 2017.
- [15] Neil Davies, Francis X Villablanca, and George K Roderick. Bioinvasions of the Medfly *Ceratitis capitata*: Source estimation using DNA sequences at multiple intron loci. *Genetics*, 153(1):351–360, September 1999. ISSN 0016-6731, 1943-2631.
- [16] D.S. Haymer, M. He, and D.O. McInnis. Genetic marker analysis of spatial and temporal relationships among existing populations and new infestations of the Mediterranean fruit fly (*Ceratitis capitata*). *Heredity*, 79(3):302–309, 1997.
- [17] M. Bonizzoni, L. Zheng, CR Guglielmino, DS Haymer, G. Gasperi, LM Gomulski, and AR Malacrida. Microsatellite analysis of medfly bioinfestations in California. *Molecular Ecology*, 10(10):2515–2524, 2001.
- [18] G. Gasperi, M. Bonizzoni, L.M. Gomulski, V. Murelli, C. Torti, A.R. Malacrida, and C.R. Guglielmino. Genetic Differentiation, Gene Flow and the Origin of Infestations of the Medfly, *Ceratitis Capitata*. *Genetica*, 116(1):125–135, 2002. ISSN 0016-6707. doi: 10.1023/A:1020971911612. URL <http://www.springerlink.com/content/m40626t713257367/abstract/>.
- [19] AM Sznyszewska, NC Leppla, Z Huang, and AJ Tatem. Analysis of seasonal risk for importation of the mediterranean fruit fly, *ceratitis capitata* (diptera: Tephritidae), via air passenger traffic arriving in florida and california. *Journal of economic entomology*, 109(6):2317–2328, 2016.
- [20] DO Mcinnis, J Hendrichs, T Shelly, N Barr, K Hoffman, R Rodriguez, DR Lance, K Bloem, DM Suckling, W Enkerlin, et al. Can polyphagous invasive tephritid pest populations escape detection for years under favorable climatic and host conditions? *American Entomologist*, 63(2):89–99, 2017.
- [21] AJ Gilbert, RR Bingham, MA Nicolas, and RA Clark. *Insect trapping guide, 13th edition*. CDEA., Sacramento CA, 2013.
- [22] California code of regulations, title 3, section 3406. URL <https://www.cdfe.ca.gov/plant/medfly/docs/regs/3406-TXT-medfly.pdf>. Last visited 2017-07-17.

- [23] Nicholas C. Manoukis and Kevin Hoffman. An agent-based simulation of extirpation of *Ceratitis capitata* applied to invasions in California. *Journal of Pest Science*, 87(1):39–51, March 2014. ISSN 1612-4758, 1612-4766. doi: 10.1007/s10340-013-0513-y. URL <https://link.springer.com/article/10.1007/s10340-013-0513-y>.
- [24] G. L. Baskerville and P Emin. Rapid estimation of heat accumulation from maximum and minimum temperatures. *Ecology*, 50(3):514–517, 1969. doi: 10.2307/1933912.
- [25] Nicholas C. Manoukis, Brian Hall, and Scott M. Geib. A computer model of insect traps in a landscape. *Scientific Reports*, 4:7015, November 2014. doi: 10.1038/srep07015. WOS:000344760700005.
- [26] Adam Smith, Neal Lott, and Russ Vose. The Integrated Surface Database: Recent Developments and Partnerships. *Bulletin of the American Meteorological Society*, 92(6):704–708, June 2011. doi: 10.1175/2011BAMS3015.1.
- [27] Integrated Surface Database (ISD) | National Centers for Environmental Information (NCEI) formerly known as National Climatic Data Center (NCDC). URL <https://www.ncdc.noaa.gov/isd>. Last visited 2017-07-05.
- [28] Mediterranean fruit fly: Regulation and quarantine boundaries. URL <https://www.cdfa.ca.gov/plant/medfly/regulation.html>. Last visited 2017-07-17.
- [29] William J. Roltsch, Frank G. Zalom, Ann J. Strawn, Joyce F. Strand, and Michael J. Pitcairn. Evaluation of several degree-day estimation methods in california climates. *International Journal of Biometeorology*, 42(4):169–176, Mar 1999. doi: 10.1007/s004840050101.
- [30] S. M. Blower and H. Dowlatabadi. Sensitivity and uncertainty analysis of complex models of disease transmission: An hiv model, as an example. *International Statistical Review / Revue Internationale de Statistique*, 62(2): 229–243, 1994. ISSN 03067734, 17515823. URL <http://www.jstor.org/stable/1403510>.
- [31] P. S. Messenger and N. E. Flitters. Bioclimatic Studies of Three Species of Fruit Flies in Hawaii. *Journal of Economic Entomology*, 47(5):756–765, October 1954. ISSN 0022-0493. doi: 10.1093/jee/47.5.756. URL <https://academic.oup.com/jee/article/47/5/756/2205684/Bioclimatic-Studies-of-Three-Species-of-Fruit>.
- [32] Andrew Paul Gutierrez and Luigi Ponti. Assessing the invasive potential of the Mediterranean fruit fly in California and Italy. *Biological Invasions*, 13(12):2661–2676, January 2011. ISSN 1387-3547, 1573-1464. doi: 10.1007/s10530-011-9937-6. URL http://www.springerimages.com/Images/LifeSciences/1-10.1007_s10530-011-9937-6-1.