

# FOURTH ORDER MODIFIED LAGUERRE'S METHOD

Thomas R. Cameron

Mathematics and Computer Science Department, Davidson College

## Abstract

We present a novel modification of Laguerre's method that results in a method for the concurrent approximation of all roots of a univariate polynomial. Our method has strong virtues including fourth-order convergence that is observed in practice and belonging to the class of embarrassingly parallel algorithms. A Fortran 90 implementation of our algorithm is available online and comparisons with several other software are provided.

## The Algorithm

Let  $p(\lambda) = a_0 + a_1\lambda + \dots + a_m\lambda^m$  be a polynomial with  $a_0a_m \neq 0$  and denote by  $(z_1, \dots, z_m)$  the current approximations to the roots of  $p(\lambda)$ . The  $j$ th approximation is updated via

$$\hat{z}_j = z_j - \frac{m}{G_j \pm \sqrt{(m-1)(mH_j - G_j^2)}}, \quad (1)$$

where

$$G_j = \frac{p'(z_j)}{p(z_j)} - \sum_{\substack{i=1 \\ i \neq j}}^m \frac{1}{(z_j - z_i)} \quad \text{and} \quad H_j = - \left( \frac{p'(z_j)}{p(z_j)} \right)' - \sum_{\substack{i=1 \\ i \neq j}}^m \frac{1}{(z_j - z_i)^2}. \quad (2)$$

On each iteration,  $z_j$  is updated for  $j = 1, \dots, m$ , unless it was accepted on a previous iteration.

**Initial Estimates.** In essence, we select complex numbers along circles of suitable radii. What constitutes suitable radii is formalized in [3] and can be computed via the upper envelope of the convex hull of the set  $\{(i, \log |a_i|), i = 0, 1, \dots, m\}$ . We compute the convex hull via Andrew's Monotone Chain algorithm [1].

**Backward Error.** The backward error of an approximate root  $\xi$  is given by

$$\eta(\xi) = \frac{|p(\xi)|}{\alpha(\xi)}, \quad (3)$$

where  $\alpha(\xi) = \sum_{i=0}^m |e_i| |\xi|^i$  and  $e_i$  represent tolerances against which perturbations are measured. We accept a root approximation  $\xi$  if  $\eta(\xi) < \mu$ , where  $\mu$  is machine precision.

**Condition.** The condition of a nonzero approximate root  $\xi$  is given by

$$\kappa(\xi) = \frac{\alpha(\xi)}{|\xi| |p'(\xi)|}. \quad (4)$$


If the root approximation  $\xi$  is accepted, then we also return its condition.

## Examples

**Initial Estimates.** Let

$$p(\lambda) = 1 + 3 \cdot 10^3 \lambda + 3 \cdot 10^6 \lambda^2 + 1 \cdot 10^9 \lambda^9 + \lambda^{10}.$$

The initial estimates and exact roots of  $p$  are below.



**Convergence.** Here we test the convergence of the roots of three polynomials. The first polynomial is  $z^5 - 1$ , the second is the degree 10 Chebyshev polynomial, and the third is  $z^{10} + \dots + z + 1$ . The error is measured as the maximum relative forward error. For each polynomial, the error after each iteration is recorded in the table below.

Iteration	Error-1	Error-2	Error-3
1	0.55	3.57	0.37
2	0.14	2.27	1.32
3	$1.91 \cdot 10^{-4}$	0.22	$2.55 \cdot 10^{-2}$
4	$3.33 \cdot 10^{-16}$	0.16	$5.93 \cdot 10^{-8}$
5	$3.33 \cdot 10^{-16}$	$1.49 \cdot 10^{-3}$	$1.96 \cdot 10^{-15}$
6	0	$2.39 \cdot 10^{-13}$	$1.96 \cdot 10^{-15}$
7	0	$1.02 \cdot 10^{-14}$	0
8	0	$1.02 \cdot 10^{-14}$	0
9	0	0	0
10	0	0	0

## Numerical Experiments

Comparisons between FPML, Polzeros [3], and the singleshift version of AMVW [2].




Fig. 1: Random Polynomials




Fig. 2: Roots of Unity

Poly No.	Description	Deg.	Roots
1	Wilkinson polynomial	10	1, ..., 10
2	Wilkinson polynomial	15	1, ..., 15
3	Wilkinson polynomial	20	1, ..., 20
4	scale and shifted Wilkinson polynomial	20	-2.1, -1.9, ..., 1.7
5	reverse Wilkinson polynomial	10	1.1/2, ..., 1/10
6	reverse Wilkinson polynomial	15	1.1/2, ..., 1/15
7	reverse Wilkinson polynomial	20	1.1/2, ..., 1/20
8	prescribed roots of varying scale	20	$2^{-10}, 2^{-9}, \dots, 2^9$
9	prescribed roots of varying scale -3	20	$2^{-10} - 3, 2^{-9} - 3, \dots, 2^9 - 3$
10	Chebyshev polynomial	20	$\cos(\frac{2j-1}{20}\pi)$
11	$z^{20} + z^{19} + \dots + z + 1$	20	$e^{i\frac{2\pi j}{20}}$
12	C. Traverso	24	known
13	Mandelbrot	31	known
14	Mandelbrot	63	known

Fig. 3: Special Polynomials

## Conclusion

Fortran 90 code along with installation instructions and additional experiment results and references are provided at <https://github.com/trcameron/FPML>.

## References

- [1] A. M. Andrew, *Another efficient algorithm for convex hulls in two dimensions*, Info. Proc. Letters **9** (1979), no. 15, 216–219.
- [2] J. L. Aurentz, T. Mach, R. Vandebril, and D. S. Watkins, *Fast and backward stable computation of roots of polynomials*, SIAM J. Matrix Anal. Appl. **36** (2015), no. 3, 942–973.
- [3] D. A. Bini, *Numerical computation of polynomial zeros by means of Aberth's method*, Numer. Algorithms **13** (1996), 179–200.