

---



# Linear Algebra

주재걸


고려대학교 컴퓨터학과




# ONTO and ONE-TO-ONE



- **Definition:** A mapping  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be **onto**  $\mathbb{R}^m$  if each  $b \in \mathbb{R}^m$  is the image of **at least** one  $x \in \mathbb{R}^n$ . That is, the range is equal to the co-domain.



$T$  is NOT onto  $\mathbb{R}^m$





$T$  is onto  $\mathbb{R}^m$

# ONTO and ONE-TO-ONE

2    - - > 3    one - to - one


3    - - > 2

- **Definition:** A mapping  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be **one-to-one** if each  $b \in \mathbb{R}^m$  is the image of **at most** one  $x \in \mathbb{R}^n$ . That is, each output vector in the range is mapped by only one input vector, no more than that.




# Neural Network Example

- Fully-connected layers



# Neural Network Example: ONE-TO-ONE


- Will there be many (or unique) people mapped to the same (over\_weighted, tall\_and\_smoking)?





# Neural Network Example: ONTO

- Is there any (over\_weighted, tall\_and\_smoking) that does not exist at all?





# ONTO and ONE-TO-ONE

- Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, i.e.,

$$T(\mathbf{x}) = A\mathbf{x} \text{ for all } \mathbf{x} \in \mathbb{R}^n.$$

- $T$  is **one-to-one** if and only if the columns of  $A$  are **linearly independent**.
- $T$  maps  $\mathbb{R}^n$  **onto**  $\mathbb{R}^m$  if and only if the columns of  $A$  **span**  $\mathbb{R}^m$ .



# ONTO and ONE-TO-ONE

- **Example:**

$$\text{Let } T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 2 & 0 \\ -1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

- Is  $T$  one-to-one?
- Does  $T$  map  $\mathbb{R}^2$  onto  $\mathbb{R}^3$ ?



# ONTO and ONE-TO-ONE

- **Example:**

$$\text{Let } T(\mathbf{x}) = A\mathbf{x} = \begin{bmatrix} 1 & 4 & 5 \\ 2 & 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- Is  $T$  one-to-one?
- Does  $T$  map  $\mathbb{R}^3$  onto  $\mathbb{R}^2$ ?



# Further Study

- Gaussian elimination, row reduction, echelon form
  - Lay Ch1.2,
- LU factorization: efficiently solving linear systems
  - Lay Ch2.5
- Computing invertible matrices
  - Lay Ch2.2
- Invertible matrix theorem for square matrices
  - Lay Ch2.3, Ch2.9