

# The Two Set Relations Generating Geometry

George. M. Van Treeck

ABSTRACT. Where each domain set has a corresponding range set, distance is the cardinal of the union of the range sets and volume is the cardinal of the Cartesian product of disjoint range sets. A ruler (measuring stick) partitions intervals approximately into sets of size  $c$  subintervals and sums the sizes. The distance and volume set operations on the sets of subintervals converge to the properties of metric space, Manhattan distance, Euclidean distance, and Euclidean volume as  $c$  goes to 0. The volume proof allows simpler derivations of Coulomb's charge force and Newton's gravity force equations without using other laws of physics or Gauss's divergence theorem. Spacetime limits physical distance and volume to 3 dimensions. All proofs are verified in Coq.

## CONTENTS

1. Introduction	1
2. Ruler measure and convergence	2
3. Distance	2
4. Euclidean Volume	6
5. Applications to physics	7
6. Insights and implications	10
References	12

## 1. Introduction

Metric space, the Euclidean distance as a metric/vector norm, and Euclidean area/volume in the Lebesgue, Borel, and Hausdorff measures have been definitions [Gol76] [Rud76]. In this article, where each disjoint domain set has a corresponding range set, distance is the cardinal of the union of the range sets and volume is the cardinal of the Cartesian product of disjoint range sets. A ruler (measuring stick) measures interval lengths by partitioning all domain and range intervals approximately into sets of size  $c$  subintervals and summing the sizes. The distance and volume set operations on the sets of subintervals converges to the properties of

metric space, intersecting range sets converge to Euclidean distance, disjoint range sets converge to Manhattan distance, and Euclidean volume as  $c \rightarrow 0$ .

The derivations of metric space, distance equations, and volume from the set operations provide some insights into geometry and physics, for example: the constraint on the union operation generating the symmetry property of metric space; the domain-to-range set mapping that makes Euclidean distance the smallest possible distance between two distinct points in  $\mathbb{R}^n$ ; the constraints on domain and range sets that generate both Manhattan distance and Euclidean volume; how the Euclidean volume proof allows deriving Coulomb's charge force and Newton's gravity force equations without using other laws of physics or Gauss's divergence theorem; how the proportionate interval principle generates the inverse square law; how spacetime places an additional constraint on physical sets of intervals, which limits physical distance and volume to 3 dimensions.

All the proofs in this article have been formally verified using using the Coq proof verification system [Coq15]. The formal proofs are located in the Coq files, “euclidrelations.v” and “threed.v,” at: <https://github.com/treeck/RASRGeometry>.

## 2. Ruler measure and convergence

**DEFINITION 2.1.** Ruler measure: A ruler measures the size,  $M$ , of a closed, open, or semi-open interval as the sum of the sizes of the nearest integer number of whole subintervals,  $p$ , each subinterval having the same size,  $c$ . Notionally:

$$(2.1) \quad \forall c, s \in \mathbb{R}, [a, b] \subset \mathbb{R}, s = |a - b| \wedge c > 0 \wedge \\ (p = \text{floor}(s/c) \vee p = \text{ceiling}(s/c)) \wedge M = \sum_{i=1}^p c = pc.$$

**THEOREM 2.2.** *Ruler convergence:*

$$\forall [a, b] \subset \mathbb{R}, s = |a - b| \Rightarrow M = \lim_{c \rightarrow 0} pc = s.$$

The theorem, “limit\_c.0.M.eq\_exact\_size,” and formal proof is in the Coq file, euclidrelations.v.

**PROOF.** (epsilon-delta proof)

By definition of the floor function,  $\text{floor}(x) = \max(\{y : y \leq x, y \in \mathbb{Z}, x \in \mathbb{R}\})$ :

$$(2.2) \quad \forall c > 0, p = \text{floor}(s/c) \wedge 0 \leq |\text{floor}(s/c) - s/c| < 1 \Rightarrow 0 \leq |p - s/c| < 1.$$

Multiply all sides of inequality 2.2 by  $|c|$ :

$$(2.3) \quad \forall c > 0, 0 \leq |p - s/c| < 1 \Rightarrow 0 \leq |pc - s| < |c|.$$

$$(2.4) \quad \forall \delta : |pc - s| < |c| = |c - 0| < \delta \\ \Rightarrow \forall \epsilon = \delta : |c - 0| < \delta \wedge |pc - s| < \epsilon := M = \lim_{c \rightarrow 0} pc = s. \quad \square$$

The proof steps using the ceiling function (the outer measure) are the same as the steps in the previous proof using the floor function (the inner measure). The following is an example of ruler convergence for the interval,  $[0, \pi]$ :  $s = |0 - \pi|$ , and  $p = \text{floor}(s/c) \Rightarrow p \cdot c = 3.1_{c=10^{-1}}, 3.14_{c=10^{-2}}, 3.141_{c=10^{-3}}, \dots, \pi_{\lim_{c \rightarrow 0}}$ .

## 3. Distance

**Notation convention:** Vertical bars around a set or list,  $|\dots|$ , indicates the cardinal (number of members in the set or list).

**3.1. Countable distance.** A concrete example of a countable distance is the number of same-sized steps walked in a range (distance) set,  $y$ , which equals the number of same-sized pieces of land in a corresponding domain set,  $x$ :  $|x| = |y|$ . Generalizing, each disjoint domain set,  $x_i$ , has a corresponding range (distance) set,  $y_i$ . The countable distance spanning the disjoint domain sets is the number of members,  $d_c$ , in the union range set:

DEFINITION 3.1. Countable distance,  $d_c$  :

$$d_c = |\bigcup_{i=1}^n y_i|, \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i|.$$

THEOREM 3.2. *Inclusion-exclusion Inequality:*  $|\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i|$ .

The inclusion-exclusion inequality follows from the inclusion-exclusion principle [CG15]. But, a more intuitive and simple proof follows from the associative law of addition where the sum of set sizes is equal to the size of all the set members appended into a list and the commutative law of addition that allows sorting that list into a list of unique members (the *union* set) and a list of duplicates. The duplicates being  $\geq 0$  implies the union size is always  $\leq$  the sum of set sizes.

A formal proof, `inclusion_exclusion_inequality`, using sorting into a set of unique members (*union* set) and a list of duplicates, is in the file `euclidrelations.v`.

PROOF.

$$(3.1) \quad \begin{aligned} \sum_{i=1}^n |y_i| &= |\text{append}_{i=1}^n y_i| = |\text{sort}(\text{append}_{i=1}^n y_i)| \\ &= |\bigcup_{i=1}^n y_i| + |\text{duplicates}_{i=1}^n y_i|. \end{aligned}$$

$$(3.2) \quad \begin{aligned} |\bigcup_{i=1}^n y_i| + |\text{duplicates}_{i=1}^n y_i| &= \sum_{i=1}^n |y_i| \quad \wedge \quad |\text{duplicates}_{i=1}^n y_i| \geq 0 \\ \Rightarrow \quad |\bigcup_{i=1}^n y_i| &\leq \sum_{i=1}^n |y_i|. \quad \square \end{aligned}$$

**3.2. Metric Space.** All function range intervals,  $d(u, w)$ , satisfying the countable distance definition,  $d_c = |\bigcup_{i=1}^n y_i|$ , where the ruler is applied, generates three of the four metric space properties: triangle inequality, non-negativity, and identity of indiscernibles. The set-based reason for the fourth property of metric space, symmetry [ $d(u, v) = d(v, u)$ ], will be identified in the last section of this article. The formal proofs: `triangle_inequality`, `non_negativity`, and `identity_of_indiscernibles` are in the Coq file, `euclidrelations.v`.

THEOREM 3.3. *Triangle Inequality:*  $d_c = |y_1 \cup y_2| \Rightarrow d(u, w) \leq d(u, v) + d(v, w)$ .

PROOF. Apply the ruler measure (2.1), the countable distance condition (3.1), inclusion-exclusion inequality (3.2), and then ruler convergence (2.2).

$$(3.3) \quad \begin{aligned} \forall c > 0, \quad d(u, w), \quad d(u, v), \quad d(v, w) : \\ |y_1| &= \text{floor}(d(u, v)/c) \quad \wedge \quad |y_2| = \text{floor}(d(v, w)/c) \quad \wedge \\ d_c &= \text{floor}(d(u, w)/c) \quad \wedge \quad d_c = |y_1 \cup y_2| \leq |y_1| + |y_2| \\ \Rightarrow \quad \text{floor}(d(u, w)/c) &\leq \text{floor}(d(u, v)/c) + \text{floor}(d(v, w)/c). \\ \Rightarrow \quad \text{floor}(d(u, w)/c) \cdot c &\leq \text{floor}(d(u, v)/c) \cdot c + \text{floor}(d(v, w)/c) \cdot c \\ \Rightarrow \quad \lim_{c \rightarrow 0} \text{floor}(d(u, w)/c) \cdot c &\leq \lim_{c \rightarrow 0} \text{floor}(d(u, v)/c) \cdot c + \lim_{c \rightarrow 0} \text{floor}(d(v, w)/c) \cdot c \\ &\Rightarrow \quad d(u, w) \leq d(u, v) + d(v, w). \quad \square \end{aligned}$$

**THEOREM 3.4.** *Non-negativity:*  $d_c = |y_1 \cup y_2| \Rightarrow d(u, w) \geq 0$ .

**PROOF.** By definition, a set always has a size (cardinal)  $\geq 0$ :

$$(3.4) \quad \forall c > 0, d(u, w) : \quad \text{floor}(d(u, w)/c) = d_c \quad \wedge \quad d_c = |y_1 \cup y_2| \geq 0 \\ \Rightarrow \quad \text{floor}(d(u, w)/c) = d_c \geq 0 \quad \Rightarrow \quad d(u, w) = \lim_{c \rightarrow 0} d_c \cdot c \geq 0. \quad \square$$

**THEOREM 3.5.** *Identity of Indiscernibles:*  $d(w, w) = 0$ .

**PROOF.** Apply the triangle inequality property (3.3):

$$(3.5) \quad \forall d(u, v) = d(v, w) = 0 \quad \wedge \quad d(u, w) \leq d(u, v) + d(v, w) \quad \Rightarrow \quad d(u, w) \leq 0.$$

Combine the non-negativity property (3.4) and the previous inequality (3.5):

$$(3.6) \quad d(u, w) \geq 0 \quad \wedge \quad d(u, w) \leq 0 \quad \Leftrightarrow \quad 0 \leq d(u, w) \leq 0 \quad \Rightarrow \quad d(u, w) = 0.$$

Combine the result of step 3.6 and the condition,  $d(u, v) = 0$ , in step 3.5.

$$(3.7) \quad d(u, w) = 0 \quad \wedge \quad d(u, v) = 0 \quad \Rightarrow \quad w = v.$$

Combine the condition,  $d(v, w) = 0$ , in step 3.5 and the result of step 3.7.

$$(3.8) \quad d(v, w) = 0 \quad \wedge \quad w = v \quad \Rightarrow \quad d(w, w) = 0. \quad \square$$

**3.3. Countable distance range.** From the countable distance definition,  $d_c = |\bigcup_{i=1}^n y_i|$ , as the amount of intersection increases, more domain set members can map to a single range set member. Therefore, the countable distance,  $d_c$ , is a function of the number of domain-to-range set member mappings.

From the countable distance property (3.1), where  $|x_i| = |y_i| = p_i$ , the range of possible domain-to-range set member mappings is the one-to-one (bijective),  $1 \cdot |y_i| = p_i$ , number of mappings, to the many-to-many,  $|x_i| \cdot |y_i| = p_i^2$ , number of mappings. Therefore, the total number of domain-to-range set mappings varies from  $\sum_{i=1}^n (1 \cdot |y_i|) = \sum_{i=1}^n p_i$  to  $\sum_{i=1}^n (|y_i| \cdot |x_i|) = \sum_{i=1}^n p_i^2$  mappings. Applying the ruler (2.1) and ruler convergence theorem (2.2) to the smallest and largest total number of domain-to-range set mapping cases converges to the real-valued Manhattan and Euclidean distance relations.

### 3.4. Manhattan distance.

**THEOREM 3.6.** *Manhattan (largest) distance,  $d$ , is the size of the range interval,  $[d_0, d_m]$ , corresponding to a set of disjoint domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d = \sum_{i=1}^n s_i, \quad d = |d_0 - d_m|, \quad s_i = |a_i - b_i|.$$

The theorem, “taxicab\_distance,” and formal proof is in the Coq file, euclidrelations.v.

**PROOF.**

From the countable distance definition (3.1) and the inclusion-exclusion inequality (3.2), the largest possible countable distance,  $d_c$ , is the equality case:

$$(3.9) \quad d_c = |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i| \quad \wedge \quad |x_i| = |y_i| = p_i \\ \Rightarrow \quad d_c \leq \sum_{i=1}^n |y_i| = \sum_{i=1}^n p_i \quad \Rightarrow \quad \exists p_i, d_c : d_c = \sum_{i=1}^n p_i.$$

Multiply both sides of equation 3.11 by  $c$  and take the limit:

$$(3.10) \quad d_c = \sum_{i=1}^n p_i \Rightarrow d_c \cdot c = \sum_{i=1}^n (p_i \cdot c) \Rightarrow \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c).$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to the definition of  $d$ :

$$(3.11) \quad d = |d_0 - d_m| \Rightarrow \exists c \, d : \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to each domain interval:

$$(3.12) \quad s_i = |a_i - b_i| \quad \wedge \quad \text{floor}(s_i/c) = |x_i| = |y_i| = p_i \Rightarrow \lim_{c \rightarrow 0} p_i \cdot c = s_i.$$

Combine equations 3.11, 3.10, 3.12:

$$(3.13) \quad d = \lim_{c \rightarrow 0} d_c \cdot c \quad \wedge \quad \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) \quad \wedge \\ \lim_{c \rightarrow 0} (p_i \cdot c) = s_i \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \sum_{i=1}^n s_i. \quad \square$$

### 3.5. Euclidean distance.

**THEOREM 3.7.** *Euclidean (smallest) distance,  $d$ , is the size of the range interval,  $[d_0, d_m]$ , corresponding to a set of disjoint domain intervals,*

*$\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d^2 = \sum_{i=1}^n s_i^2, \quad d = |d_0 - d_m|, \quad s_i = |a_i - b_i|.$$

The theorem, “Euclidean\_distance,” and formal proof is in the Coq file, euclidrelations.v.

**PROOF.**

Apply the rule of product to the largest number of domain-to-range set mappings, where all  $p_i$  number of range set members,  $y_i$ , map to each of the  $p_i$  number of members in the domain set,  $x_i$ , which is the Cartesian product,  $|y_i| \cdot |x_i|$ :

$$(3.14) \quad |x_i| = |y_i| = p_i \Rightarrow \sum_{i=1}^n |y_i| \cdot |x_i| = \sum_{i=1}^n p_i^2.$$

From the countable distance definition (3.1) and the inclusion-exclusion inequality (3.2), choose the equality case:

$$(3.15) \quad d_c = |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i| \quad \wedge \quad |x_i| = |y_i| = p_i \\ \Rightarrow d_c \leq \sum_{i=1}^n |y_i| = \sum_{i=1}^n p_i \Rightarrow \exists p_i, d_c : d_c = \sum_{i=1}^n p_i.$$

Square both sides of equation 3.15 ( $x = y \Leftrightarrow f(x) = f(y)$ ):

$$(3.16) \quad \exists p_i, d_c : d_c = \sum_{i=1}^n p_i \Leftrightarrow \exists p_i, d_c : d_c^2 = (\sum_{i=1}^n p_i)^2.$$

Apply the square of sum inequality,  $(\sum_{i=1}^n p_i)^2 \geq \sum_{i=1}^n p_i^2$ , to equation 3.16 and select the smallest area (the equality) case:

$$(3.17) \quad d_c^2 = (\sum_{i=1}^n p_i)^2 = \sum_{i=1}^n p_i \sum_{j=1}^n p_j \\ = \sum_{i=1}^n p_i^2 + \sum_{i=1}^n p_i \sum_{j=1, j \neq i}^n p_j \geq \sum_{i=1}^n p_i^2 \Rightarrow \exists p_i : d_c^2 = \sum_{i=1}^n p_i^2.$$

Multiply both sides of equation 3.17 by  $c^2$ , simplify, and take the limit.

$$(3.18) \quad d_c^2 = \sum_{i=1}^n p_i^2 \Rightarrow d_c^2 \cdot c^2 = \sum_{i=1}^n p_i^2 \cdot c^2 \Leftrightarrow (d_c \cdot c)^2 = \sum_{i=1}^n (p_i \cdot c)^2 \\ \Rightarrow \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) and square both sides:

$$(3.19) \quad \exists c \, d \in \mathbb{R} : \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c \Rightarrow d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to each domain interval:

$$(3.20) \quad s_i = |a_i - b_i| \quad \wedge \quad \text{floor}(s_i/c) = |x_i| = |y_i| = p_i \Rightarrow \lim_{c \rightarrow 0} p_i \cdot c = s_i.$$

Combine equations 3.19, 3.18, 3.20:

$$(3.21) \quad d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 \quad \wedge \quad \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2 \quad \wedge \\ \lim_{c \rightarrow 0} (p_i \cdot c) = s_i \quad \Rightarrow \quad d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2 = \sum_{i=1}^n s_i^2. \quad \square$$

#### 4. Euclidean Volume

The case of a 1-1 correspondence of domain-to-range set mappings and disjoint range sets,  $d_c = |\bigcup_{i=1}^n y_i|$ ,  $\bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset$ , converges to Manhattan distance (3.6). The sum of same-sized sub-Manhattan distances is the sum of ordered n-tuples of subintervals, where each n-tuple is an ordered set containing one same-sized subinterval from each range set,  $y_i$ . The number of all possible ordered sub-Manhattan distance n-tuples,  $v_c$ , is the number of ordered n-tuples generated by Cartesian product of the range sets:

DEFINITION 4.1. Countable Volume,  $v_c$ :

$$v_c = |\times_{i=1}^n y_i|, \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset.$$

THEOREM 4.2. *Euclidean volume,  $v$ , is size of the range interval,  $[v_0, v_m]$ , corresponding to the Cartesian product of all the members of the domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ . Notionally:*

$$v = \prod_{i=1}^n s_i, \quad v = |v_0 - v_m|, \quad s_i = |a_i - b_i|.$$

The theorem, “Euclidean\_volume,” and formal proof is in the Coq file, euclidrelations.v.

PROOF.

Use the ruler (2.1) to partition each of the domain intervals,  $[a_i, b_i]$ , into a set,  $x_i$ , containing  $p_i$  number of subintervals.

$$(4.1) \quad \forall i \ n \in \mathbb{N}, \quad i \in [1, n], \quad c > 0 \quad \wedge \quad \text{floor}(s_i/c) = p_i = |x_i| = |y_i|.$$

Apply the ruler convergence theorem (2.2) to equation 4.1:

$$(4.2) \quad \text{floor}(s_i/c) = p_i \quad \Rightarrow \quad \lim_{c \rightarrow 0} (p_i \cdot c) = s_i.$$

Apply the associative law of multiplication to derive the countable volume (4.1) in terms of  $p_i$ :

$$(4.3) \quad v_c = |\times_{i=1}^n y_i| \quad \wedge \quad |y_i| = p_i \quad \Rightarrow \quad v_c = |\times_{i=1}^n y_i| = \prod_{i=1}^n |y_i| = \prod_{i=1}^n p_i.$$

Multiply both sides of equation 4.3 by  $c^n$ :

$$(4.4) \quad v_c \cdot c^n = (\prod_{i=1}^n p_i) \cdot c^n = \prod_{i=1}^n (p_i \cdot c).$$

Use those cases, where  $v_c$  has an integer  $n^{\text{th}}$  root.

$$(4.5) \quad \forall n, p, v_c \in \mathbb{N} : \quad p^n = v_c \quad \Rightarrow \quad v_c \cdot c^n = p^n \cdot c^n = (p \cdot c)^n = \prod_{i=1}^n (p_i \cdot c).$$

Apply the ruler (2.1) and ruler convergence (2.2) to the range interval,  $[v_0, v_m]$  (where  $v = |v_0 - v_m|$ ), and then combine with equations 4.5 and 4.2:

$$(4.6) \quad \text{floor}(v/c^n) = p^n \Rightarrow v = \lim_{c \rightarrow 0} (p \cdot c)^n = \prod_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \prod_{i=1}^n s_i. \quad \square$$

## 5. Applications to physics

**5.1. Coulomb's charge force.** The sizes,  $q_1$  and  $q_2$ , of two charges are independent domain variables, where each infinitesimal size,  $c$ , component of a charge exerts a force on each infinitesimal size,  $c$ , component of the other charge. The total force,  $F$ , is equal to the total number of forces (the Cartesian product of the infinitesimal size,  $c$ , components) multiplied times a quantum charge force,  $m_C a_C$ . From the volume proof (4.2), the Cartesian product converges to  $q_1 q_2$ :

$$(5.1) \quad F = (m_C a_C) \left( \lim_{c \rightarrow 0} \text{floor}(q_1/c) \cdot c \right) \left( \lim_{c \rightarrow 0} \text{floor}(q_2/c) \cdot c \right) = (m_C a_C) \int_0^{q_1} \int_0^{q_2} dq^2 = (m_C a_C) (q_1 q_2).$$

From equation 5.1, a change in charge,  $q$ , causes a proportionate change in force,  $F$ . Solving for the quantum force,  $F = m_C a_C$ , requires a proportionate variable,  $r$ , to offset the effect of a change in  $q$ :  $r \propto q \Rightarrow \exists q_C/r_C \in \mathbb{R} : r(q_C/r_C) = q$ :

$$(5.2) \quad \forall q_1, q_2 \geq 0 \exists q \in \mathbb{R} : q^2 = q_1 q_2 \quad \wedge \quad r(q_C/r_C) = q \Rightarrow (r(q_C/r_C))^2 = q_1 q_2.$$

$$(5.3) \quad \begin{aligned} (r(q_C/r_C))^2 &= q_1 q_2 \quad \wedge \quad F = (m_C a_C) (q_1 q_2) \\ &\Rightarrow F = (m_C a_C) (r(q_C/r_C))^2 = (m_C a_C) (q_1 q_2) \\ &\Rightarrow F = m_C a_C = (m_C a_C r_C^2 / q_C^2) q_1 q_2 / r^2 = k_C q_1 q_2 / r^2. \end{aligned}$$

where  $k_C = m_C a_C r_C^2 / q_C^2$  corresponds to the SI units:  $N m^2 C^{-2}$ .

**5.2. Newton's gravity force equation.** The sizes,  $m_1$  and  $m_2$ , of two masses are independent domain variables, where each infinitesimal size,  $c$ , component of a mass exerts a force on each infinitesimal size,  $c$ , component of the other mass. The total force,  $F$ , is equal to the total number of forces (the Cartesian product of the size,  $c$ , components) multiplied times a quantum gravity force,  $m_G a_G$ . From the volume proof (4.2), the Cartesian product converges to  $m_1 m_2$ :

$$(5.4) \quad F = (m_G a_G) \left( \lim_{c \rightarrow 0} \text{floor}(m_1/c) \cdot c \right) \left( \lim_{c \rightarrow 0} \text{floor}(m_2/c) \cdot c \right) = (m_G a_G) \int_0^{m_1} \int_0^{m_2} dm^2 = (m_G a_G) (m_1 m_2).$$

From equation 5.4, an change in mass,  $m$ , causes a proportionate change in force,  $F$ . Solving for the quantum force,  $F = m_G a_G$ , requires a proportionate variable,  $r$ , to offset the effect of a change in  $m$ :  $r \propto m \Rightarrow \exists m_G/r_G \in \mathbb{R} : r(m_G/r_G) = m$ :

$$(5.5) \quad \forall m_1, m_2 \geq 0 \exists m \in \mathbb{R} : \quad m^2 = m_1 m_2 \quad \wedge \quad r(m_G/r_G) = m \\ \Rightarrow (r(m_G/r_G))^2 = m_1 m_2.$$

$$(5.6) \quad \begin{aligned} (r(m_G/r_G))^2 &= m_1 m_2 \quad \wedge \quad F = (m_G a_G) (m_1 m_2) \\ &\Rightarrow F = (m_G a_G) (r(m_G/r_G))^2 = (m_G a_G) (m_1 m_2) \\ &\Rightarrow F = m_G a_G = (a_G r_G^2 / m_G) m_1 m_2 / r^2. \end{aligned}$$

$$(5.7) \quad \begin{aligned} \exists t_G \in \mathbb{R} : r_G/t_G^2 &= a_G \quad \wedge \quad F = m_G a_G = (a_G r_G^2 / m_G) m_1 m_2 / r^2 \\ &\Rightarrow F = m_G a_G = (r_G^3 / m_G t_G^2) m_1 m_2 / r^2 = G m_1 m_2 / r^2, \end{aligned}$$

where  $G = r_G^3 / m_G t_G^2$  corresponds to the SI units:  $m^3 k g^{-1} s^{-2}$ .

**5.3. Spacetime equations.** The charge (5.3) and gravity (5.7) force equations were derived from the principle that charge and mass are proportionate to distance:  $r = (r_C/q_C)q = (r_G/m_G)m$ . Suppose that time is also proportionate to distance,  $r = (r_c/t_c)t = ct$ .

Applying the ruler to two intervals,  $[0, d_1]$  and  $[0, d_2]$ , in two inertial (independent, non-accelerating) frames of reference, the distance (and time) spanning the two domain intervals converges to a range of distances (and times) from Manhattan (3.6) to Euclidean distance (3.7).

$$(5.8) \quad r^2 = d_1^2 + d_2^2 \quad \wedge \quad r = (r_c/t_c)t = ct$$

$$\Rightarrow \quad (ct)^2 = d_1^2 + d_2^2 \quad \Rightarrow \quad d_2 = \sqrt{(ct)^2 - d_1^2}.$$

$$(5.9) \quad d_2 = \sqrt{(ct)^2 - d_1^2} \quad \wedge \quad d = d_2 \quad \wedge \quad d_1 = vt$$

$$\Rightarrow \quad d = \sqrt{(ct)^2 - (vt)^2} = ct\sqrt{1 - (v^2/c^2)},$$

which is the spacetime dilation equation. [Bru17].

$$(5.10) \quad d_2^2 = (ct)^2 - d_1^2 \quad \wedge \quad s = d_2 \quad \wedge \quad d_1^2 = x^2 + y^2 + z^2$$

$$\Rightarrow \quad s^2 = (ct)^2 - (x^2 + y^2 + z^2),$$

which is one form of the spacetime interval equation [Bru17].

**5.4. 3 dimensions of physical geometry.** The case of every distance interval having a proportionate time interval generates the spacetime equations (5.3). *Deterministic* sequencing through each Manhattan distance interval, where it can be verified that all distance intervals were sequenced and none over-sequenced, requires that the intervals can be assigned a total order that does not change with respect to the corresponding Manhattan time interval.

By the commutative law of addition, the Manhattan distance (and total time) is the same for every possible ordered sum of interval sizes. Deterministic sequencing in all possible orders requires a static total order during the *time* of sequencing, where every interval is sequentially adjacent (either a successor or predecessor) to every other interval, herein referred to as a symmetric geometry.

Ordered and symmetric sets with one and two members are the trivial cases. It will now be proved that the largest ordered and symmetric set is a cyclic set that contains at most 3 members, in this case, 3 dimensions of physical distance and volume.

DEFINITION 5.1. Ordered geometry:

$$\forall i \ n \in \mathbb{N}, \ i \in [1, n-1], \ \forall x_i \in \{x_1, \dots, x_n\},$$

$$\text{successor } x_i = x_{i+1} \quad \wedge \quad \text{predecessor } x_{i+1} = x_i.$$

DEFINITION 5.2. Symmetric geometry (every set member is sequentially adjacent to every other member):

$$\forall i \ j \ n \in \mathbb{N}, \ \forall x_i \ x_j \in \{x_1, \dots, x_n\}, \ \text{successor } x_i = x_j \Leftrightarrow \text{predecessor } x_j = x_i.$$

THEOREM 5.3. An ordered and symmetric set is a cyclic set.

$$i = n \quad \wedge \quad j = 1 \quad \Rightarrow \quad \text{successor } x_n = x_1 \quad \wedge \quad \text{predecessor } x_1 = x_n.$$



The theorem, “ordered\_symmetric\_is\_cyclic,” and formal proof is in the Coq file, `threed.v`.

PROOF. A total order (5.1) defines unique successors and predecessors for all set members except for the successor of  $x_n$  and the predecessor of  $x_1$ . Therefore, the only member that can be a successor of  $x_n$ , without creating a contradiction, is  $x_1$ . And the only member that can be a predecessor of  $x_1$ , without creating a contradiction, is  $x_n$ . From the properties of a symmetric geometry (5.2):

$$(5.11) \quad i = n \wedge j = 1 \Rightarrow \text{successor } x_i = x_j = \text{successor } x_n = x_1.$$

Applying the definition of a symmetric geometry (5.2) to conclusion 5.11:

$$(5.12) \quad \text{successor } x_i = x_j \Rightarrow \text{predecessor } x_j = x_i \Rightarrow \text{predecessor } x_1 = x_n. \quad \square$$

THEOREM 5.4. *An ordered and symmetric set is limited to at most 3 members.*

The lemmas and formal proofs in the Coq file `threed.v` are:

Lemmas: `adj111`, `adj122`, `adj212`, `adj123`, `adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and `not_all_mutually_adjacent_gt_3`.

The following proof uses Horn clauses (a subset of first-order logic) that uses unification and resolution. Horn clauses make it clear which facts satisfy a proof goal.

PROOF.

It was proved that an ordered and symmetric set is a cyclic set (5.3). In other words, the successors and predecessors of an ordered and symmetric set are cyclic:

DEFINITION 5.5. Cyclic successor of  $m$  is  $n$ :

$$(5.13) \quad \text{Successor}(m, n, \text{setsize}) \leftarrow (m = \text{setsize} \wedge n = 1) \vee (n = m + 1 \leq \text{setsize}).$$

DEFINITION 5.6. Cyclic predecessor of  $m$  is  $n$ :

$$(5.14) \quad \text{Predecessor}(m, n, \text{setsize}) \leftarrow (m = 1 \wedge n = \text{setsize}) \vee (n = m - q \geq 1).$$

DEFINITION 5.7. Adjacent: member  $m$  is sequentially adjacent to member  $n$  if the cyclic successor of  $m$  is  $n$  or the cyclic predecessor of  $m$  is  $n$ . Notionally:

$$(5.15) \quad \text{Adjacent}(m, n, \text{setsize}) \leftarrow \text{Successor}(m, n, \text{setsize}) \vee \text{Predecessor}(m, n, \text{setsize}).$$

Prove that every member is adjacent to every other member, where  $\text{setsize} \in \{1, 2, 3\}$ :

$$(5.16) \quad \text{Adjacent}(1, 1, 1) \leftarrow \text{Successor}(1, 1, 1) \leftarrow (m = \text{setsize} \wedge n = 1).$$

$$(5.17) \quad \text{Adjacent}(1, 2, 2) \leftarrow \text{Successor}(1, 2, 2) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.18) \quad \text{Adjacent}(2, 1, 2) \leftarrow \text{Successor}(2, 1, 2) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(5.19) \quad \text{Adjacent}(1, 2, 3) \leftarrow \text{Successor}(1, 2, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.20) \quad \text{Adjacent}(2, 1, 3) \leftarrow \text{Predecessor}(2, 1, 3) \leftarrow (n = m - q \geq 1).$$

$$(5.21) \quad \text{Adjacent}(3, 1, 3) \leftarrow \text{Successor}(3, 1, 3) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(5.22) \quad \text{Adjacent}(1, 3, 3) \leftarrow \text{Predecessor}(1, 3, 3) \leftarrow (m = 1 \wedge n = \text{setsize}).$$

$$(5.23) \quad \text{Adjacent}(2, 3, 3) \leftarrow \text{Successor}(2, 3, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.24) \quad \text{Adjacent}(3, 2, 3) \leftarrow \text{Predecessor}(3, 2, 3) \leftarrow (n = m - q \geq 1).$$

Must prove that for all  $\text{setsize} > 3$ , there exist non-adjacent members. For example, the first and third members are not ( $\neg$ ) adjacent:

$$(5.25) \quad \forall \text{setsize} > 3 : \quad \neg \text{Successor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Successor}(1, 2, \text{setsize} > 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

That is, member 2 is the only successor of member 1 for all  $\text{setsize} > 3$ , which implies member 3 is not a successor of member 1 for all  $\text{setsize} > 3$ .

$$(5.26) \quad \forall \text{setsize} > 3 : \quad \neg \text{Predecessor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Predecessor}(1, \text{setsize}, \text{setsize} > 3) \leftarrow (m = 1 \wedge n = \text{setsize} > 3).$$

That is, member  $n = \text{setsize} > 3$  is the only predecessor of member 1, which implies member 3 is not a predecessor of member 1 for all  $\text{setsize} > 3$ .

$$(5.27) \quad \forall \text{setsize} > 3 : \quad \neg \text{Adjacent}(1, 3, \text{setsize} > 3) \\ \leftarrow \neg \text{Successor}(1, 3, \text{setsize} > 3) \wedge \neg \text{Predecessor}(1, 3, \text{setsize} > 3). \quad \square$$

That is, for all  $\text{setsize} > 3$ , some elements are not sequentially adjacent to every other element (not symmetric).

## 6. Insights and implications

Applying the ruler measure (2.1) and ruler convergence (2.2) to the set relations, countable distance (3.1) and countable volume (4.1) yields the following insights and implications:

- (1) The properties of metric space, Euclidean distance and area/volume are derived from two set operations without using the notions of Euclidean geometry like plane, side, angle, perpendicular, congruence, intersection, etc. [Joy98].
- (2) Making the measures, countable distance (3.1) and volume (4.1), independent of the measure of intervals (2.1) allows deriving geometric relations and insights into geometry that metric space, the Lebesgue, Borel, and Hausdorff measures are incapable of providing.

- (a) The number of all possible ordered sub-Manhattan distance n-tuples,  $v_c$ , is the number of ordered n-tuples generated by Cartesian product of the range sets, which is a set-based pattern relating volume to distance:

$$d_c = |\bigcup_{i=1}^n y_i|, \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset \Rightarrow \\ v_c = |\times_{i=1}^n y_i|, \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset.$$

- (b) The largest intersection (smallest distance),  $d_c = |\bigcup_{i=1}^n y_i|$ , has the most domain set members mapping to each range set member. The equality constraint,  $|x_i| = |y_i| = p_i$ , on countable distance (3.1) limits the largest total number of domain-to-range set mappings to

$\sum_{i=1}^n |x_i| \cdot |y_i| = \sum_{i=1}^n p_i^2$ , which is the set-based reason Euclidean distance (3.7) is the smallest possible distance between two distinct points in  $\mathbb{R}^n$  (flat space, where  $|x_i| = |y_i|$ ).

- (c) Generalizing the countable distance constraint,  $|x_i| = |y_i|$ , (3.1) to  $|x_i| = |y_i|^q$ ,  $q \geq 0$  generates all the  $L^p$  norms,  $\|L\|_p = (\sum_{i=1}^n s_i^p)^{1/p}$ . For example, using the same proof pattern as for Euclidean distance (3.7):  $p_i = |y_i| \Rightarrow |x_i| = p_i^q \Rightarrow \sum_{i=1}^n |x_i| \cdot |y_i| = \sum_{i=1}^n p_i^{q+1} \leq d_c^{q+1} \dots d = (\sum_{i=1}^n s_i^{q+1})^{1/(q+1)}$ .
  - (d) It was shown that the countable distance set operation,  $d_c = |\bigcup_{i=1}^n y_i|$ , (3.1) generates three of the metric space properties (3.2). The countable distance constraint,  $|x_i| = |y_i|$ , (3.1) is the reason for the fourth property of metric space, symmetry,  $d(u, v) = d(v, u)$ , which causes the same combinatorial domain-to-range set mapping for every domain-range set pair that manifests as the same value,  $p$ , in each term of an  $L^p$  norm.
  - (e) The Euclidean volume proof was used to derive the Coulomb's charge force (5.1) and Newton's gravity force (5.4) without using other laws of physics or Gauss's divergence theorem.
- (3) **The Proportionate Interval Principle:** The derivations of the charge force, gravity force, and spacetime equations using the ruler-based Euclidean volume and distance derivations shows that all Euclidean distance range intervals having a size,  $r$ , have proportionately sized intervals of other types, for example:  $r = (r_C/q_C)q = (r_G/m_G)m = (r_c/t_c)t$ .
- (a) The derivations of charge and gravity forces requiring the ratios,  $r = (r_C/q_C)q = (r_G/m_G)m$ , implies that if there are quantum values of charge,  $q_C$ , and mass,  $m_G$ , then there are quantum distances,  $r_C$  and  $r_G$ , where the charge and gravity forces do not exist (are not defined) at smaller distances, which might have implications for the strong force, fine-structure constant, particle collisions, and the density of black holes.
  - (b) A countable set of values has measure 0 and thus no proportion relationship to distance. Therefore, a countable set of state value changes with respect to time are independent of distance (for example, the change in the spin values of two quantum coupled particles).
- (4) Relativity theory assumes only 3 dimensions of physical space [Bru17]. The proof in this article shows that spacetime constrains physical distance and volume to at most three dimensions (5.4). Higher dimensions of physical distance and volume would have sequential adjacency contradictions.
- (a) The proof of at most 3 members in any ordered and symmetric set (5.4), implies that each *physical* infinitesimal volume (ball) can have at most 3 ordered and symmetric dimensions of discrete *physical* states of the same type. And each dimension of discrete physical states can have at most 3 ordered and symmetric discrete state values, which allows  $3 \cdot 3 \cdot 3 = 27$  possible combinations of discrete values per infinitesimal ball.
  - (b) If each of the three possible ordered and symmetric dimensions of discrete physical states contained unordered sets of discrete state values, for example, unordered binary values, then there would be

$2 \cdot 2 \cdot 2 = 8$  possible combinations of values. Unordered sets (states) are non-deterministic. For example, every sequential point in time an unordered state is physically measured, there is a 50 percent chance of having one of the binary values.

- (c) Where infinitesimal balls intersect, an algebra of the interactions of the discrete states with respect to time needs to be developed. The interaction of the discrete states associated with overlapping infinitesimal balls with respect to time might result in what we perceive as motion, waves, particles, work, force, mass, charge, etc.

## References

- [Bru17] P. Bruskiewich, *A very simple introduction to special relativity: Part two - four vectors, the lorentz transformation and group velocity (the new mathematics for the millions book 38)*, Pythagoras Publishing, 2017. ↑8, 11
- [CG15] W. Conradie and V. Goranko, *Logic and discrete mathematics*, Wiley, 2015. ↑3
- [Coq15] Coq, *Coq proof assistant*, 2015. <https://coq.inria.fr/documentation>. ↑2
- [Gol76] R. R. Goldberg, *Methods of real analysis*, John Wiley and Sons, 1976. ↑1
- [Joy98] D. E. Joyce, *Euclid's elements*, 1998. <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>. ↑10
- [Rud76] W. Rudin, *Principles of mathematical analysis*, McGraw Hill Education, 1976. ↑1

GEORGE VAN TREECK, 668 WESTLINE DR., ALAMEDA, CA 94501