

# The Real Analysis and Combinatorics of Geometry

George. M. Van Treeck

ABSTRACT. The range of combinatorial relations between members of each disjoint domain set and members of a corresponding distance set containing the same number of members, where the distance sets sometimes intersect and the set members are the same-sized subintervals of intervals, converges to the triangle inequality, Manhattan distance at the upper boundary, and Euclidean distance at the lower boundary, which provides set-based definitions of: metric space, longest, and shortest distances spanning disjoint sets. The Cartesian product of the same-sized subintervals of intervals converges to the product of interval sizes used in the Lebesgue measure and Euclidean integrals. A cyclic set of 3 dimensions emerges from the same combinatorial relations generating distance and volume. All ordered and symmetric, higher-dimensional geometries, like the spacetime four-vector, collapse into hierarchical 2 or 3-dimensional geometries. Proofs are verified in Coq.

## CONTENTS

1. Introduction	1
2. Ruler measure and convergence	3
3. Distance	3
4. Size (length/area/volume)	6
5. Ordered and symmetric geometries	7
6. Summary	9
References	11

## 1. Introduction

The triangle inequality of a metric space, the Manhattan and Euclidean distance metrics, and the volume equation (product of interval sizes) of the Lebesgue measure and Euclidean integrals (for example, Riemann and Lebesgue integrals) are imported as definitions [Gol76] rather than derived from set-based relations. Using geometric relations as primitives can not expose the set-based relationships that generate those geometric relations and other fundamental, geometric properties.

Euclidean distance and volume are derived from many-to-many, set-based relationships. But, a function only allows each domain value to map to one image value, which can not describe many-to-many relationships, where each domain value can map to many image values. Therefore, Euclidean integrals and measures, like the Lebesgue and Hausdorff measures, that use distance, area, and volume functions, and use those functions as primitives, are not capable of deriving the set-based relationships that converge to continuous, bijective distance and volume functions.

A method of measurement that allows the full range of mappings from a one-to-one correspondence (bijective) to a many-to-many mapping is required. A ruler (measuring stick) measures a real-valued interval as the nearest integer number of same-sized subintervals (units), where the partial subintervals are ignored. The ruler measure allows defining combinatorial relations, for example a many-to-many relation, between the same-sized subintervals in one interval and the same-sized subintervals in another interval. The discrete, combinatorial relations converge to continuous, bijective functions as the subinterval size converges to zero.

This article defines a set-based countable distance range having two properties: 1) For each disjoint domain set there exists a corresponding distance (image) set containing the same number of members; and 2) the distance sets sometimes intersect. The first property limits the range of intersection of the distance sets from no intersection (an inverse, one-to-one correspondence between each distance set member to a corresponding domain set member) to the maximum allowed intersection (each distance set member inversely maps to all the corresponding domain set members, which is a many-to-many domain-to-distance mapping).

When the ruler is used to divide some disjoint domain intervals and a distance interval into sets of same-sized subintervals: the no-intersection case converges to Manhattan distance, the maximum-allowed-intersection case converges to Euclidean distance, and the full-range converges to the triangle inequality. The countable distance range provides set-based definitions of: metric space, longest, and shortest distances spanning disjoint domain sets.

Using the ruler measure, the Cartesian product of the same-sized subintervals of intervals converges to the product of interval sizes (length/area/volume functions) used in the Lebesgue measure and Euclidean integrals.

The commutative property of the set-based definitions from which distance and volume are derived: 1) Allow a sequential (total) ordering of the domain sets (dimensions), herein referred to as an ordered geometry; and 2) Allow every disjoint domain set (dimension) to be sequentially adjacent to any other domain set (dimension), herein referred to as a symmetric geometry. It is proved in this article that satisfying both a total ordering and symmetry results in a cyclic set of at most three dimensions, which explains why we live in a three dimensional world.

All ordered and symmetric, higher-dimensional geometries collapse into hierarchical 2 or 3-dimensional geometries. As shown in the summary, the four-vector lengths common in physics, like the spacetime four-vector length, are 2-dimensional Euclidean distance equations that have been "flattened."

The proofs in this article are verified formally using the Coq Proof Assistant [Coq15] version 8.7.0. The Coq-based definitions, theorems, and proofs are in the files "euclidrelations.v" and "threed.v" located at:

<https://github.com/treeck/CombinatorialGeometry>.

## 2. Ruler measure and convergence

DEFINITION 2.1. Ruler measure: A ruler measures the size,  $M$ , of a closed, open, or semi-open interval as the sum of the sizes of the nearest integer number of whole subintervals,  $p$ , each subinterval having the same size,  $c$ . Notionally:

$$(2.1) \quad \forall c \, s \in \mathbb{R}, \, [a, b] \subset \mathbb{R}, \, s = |a - b| \wedge c > 0 \wedge \\ (p = \text{floor}(s/c) \vee p = \text{ceiling}(s/c)) \wedge M = \sum_{i=1}^p c = pc.$$

THEOREM 2.2. *Ruler convergence:*

$$\forall [a, b] \subset \mathbb{R}, \, s = |a - b| \Rightarrow M = \lim_{c \rightarrow 0} pc = s.$$

The Coq-based theorem and proof in the file euclidrelations.v is “limit.c\_0\_M.eq.exact\_size.”

PROOF. (epsilon-delta proof)

By definition of the floor function,  $\text{floor}(x) = \max(\{y : y \leq x, y \in \mathbb{Z}, x \in \mathbb{R}\})$ :

$$(2.2) \quad \forall c > 0, \, p = \text{floor}(s/c) \Rightarrow 0 \leq |p - s/c| < 1.$$

Multiply all sides of inequality 2.2 by  $|c|$ :

$$(2.3) \quad \forall c > 0, \, 0 \leq |p - s/c| < 1 \Rightarrow 0 \leq |pc - s| < |c|.$$

$$(2.4) \quad \forall \delta : |pc - s| < |c| = |c - 0| < \delta \\ \Rightarrow \forall \epsilon = \delta : |c - 0| < \delta \wedge |pc - s| < \epsilon := M = \lim_{c \rightarrow 0} pc = s. \quad \square$$

The proof steps using the ceiling function (the outer measure) are the same as the steps in the previous proof using the floor function (the inner measure). The following is an example of ruler convergence, where:  $[0, \pi]$ ,  $s = |\pi - 0|$ ,  $c = 10^{-i}$ , and  $p = \text{floor}(s/c) \Rightarrow p \cdot c = 3.1_{i=1}, 3.14_{i=2}, 3.141_{i=3}, \dots, \pi$ .

## 3. Distance

A simple countable distance measure is that an image (distance) set has the same number of members as a corresponding domain set. For example, the number of steps walked in a distance set must equal the number pieces of land traversed. Generalizing, for each distance set,  $y_i$ , containing  $p_i$  number of members there exists a corresponding domain set,  $x_i$ , with the same  $p_i$  number of members.

**Notation conventions:** The vertical bars around a set is the standard notation for indicating the cardinal (number of members in the set). To prevent over use of the vertical bar, the symbol for “such that” is the colon.

If the domain sets are disjoint ( $\sum_{i=1}^n |x_i| = |\bigcup_{i=1}^n x_i|$ ) and the distance sets intersect ( $\sum_{i=1}^n |y_i| > |\bigcup_{i=1}^n y_i|$ ), then a distance set member can inversely map to (be combined with) multiple domain set members. Therefore, the size of the union of the distance sets,  $d_c$ , is a function of the number of mappings to (combinations with) domain set members. Notionally:

DEFINITION 3.1. Countable distance range,  $d_c$ :

$$\forall i \, n \in \mathbb{N}, \, x_i \subseteq X, \, \sum_{i=1}^n |x_i| = |\bigcup_{i=1}^n x_i|, \, \forall x_i \exists y_i \subseteq Y : \\ |x_i| = |y_i| = p_i \wedge d_c = |Y| = |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i|.$$

The countable distance range principle (3.1),  $|x_i| = |y_i| = p_i$ , constrains the range of combinatorial relations from each member of  $y_i$  combining with (mapping to) one member of  $x_i$  to each member of  $y_i$  combining with as many as  $p_i$  number of members of  $x_i$ . More than  $p_i$  number of mappings to a member of  $x_i$  would be over-counting combinations, because a member of  $y_i$  would map to the same member of  $x_i$  more than once.

Using the rule of product, there is a range from  $|y_i| \cdot 1 = p_i$  to  $|y_i| \cdot p_i = p_i^2$  number of domain-to-distance combinations per distance set. Therefore,  $d_c = f(\sum_{i=1}^n p_i)$  is largest possible distance because it is the case of the smallest number of combinations ( $p_i$ ) per distance set (no intersection of the distance sets). And  $d_c = f(\sum_{i=1}^n p_i^2)$  is the smallest possible distance because it is the case of the largest number of combinations ( $p_i^2$ ) per distance set (largest allowed intersection of distance sets).

It will now be proved that using the ruler (2.1) to divide a set of real-valued domain intervals and a distance interval into sets of same-sized subintervals, and applying the ruler convergence theorem (2.2) to the longest and shortest distance cases converge to the real-valued, Manhattan and Euclidean distance equations.

**THEOREM 3.2.** *Manhattan (longest) distance,  $d$ , is the size of the distance interval,  $[d_0, d_m]$ , mapping to a set of disjoint domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d = \sum_{i=1}^n s_i, \quad d = |d_0 - d_m|, \quad s_i = |a_i - b_i|.$$

The formal Coq-based theorem and proof in file euclidrelations.v is “taxicab.distance.”

**PROOF.**

Use the ruler (2.1) to divide the exact size,  $s_i = |a_i - b_i|$ , of each of the domain intervals,  $[a_i, b_i]$ , into a set,  $x_i$ , containing  $p_i$  number of subintervals and apply the definition of the countable distance range (3.1), where each domain set has a corresponding distance set containing the same  $p_i$  number of members.

$$(3.1) \quad \forall i \ n \in \mathbb{N}, \quad i \in [1, n], \quad s_i \in \mathbb{R}, \quad \exists c > 0 : \quad floor(s_i/c) = p_i = |x_i| = |y_i|.$$

Next, apply the rule of product to the case of one domain set member per distance set member:

$$(3.2) \quad |y_i| = p_i \quad \Rightarrow \quad \sum_{i=1}^n |y_i| \cdot 1 = \sum_{i=1}^n p_i.$$

Apply the countable distance range definition (3.1) to equation 3.2:

$$(3.3) \quad \sum_{i=1}^n |y_i| \cdot 1 = \sum_{i=1}^n p_i \quad \wedge \quad \sum_{i=1}^n |y_i| \geq d_c \\ \Rightarrow \quad \sum_{i=1}^n p_i \geq d_c \quad \Rightarrow \quad \exists p_i, d_c : \sum_{i=1}^n p_i = d_c.$$

Multiply both sides of 3.3 by  $c$  and apply the ruler convergence theorem (2.2):

$$(3.4) \quad s_i = \lim_{c \rightarrow 0} p_i \cdot c \quad \wedge \quad \sum_{i=1}^n (p_i \cdot c) = d_c \cdot c \\ \Rightarrow \quad \sum_{i=1}^n s_i = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \lim_{c \rightarrow 0} d_c \cdot c.$$

Use the ruler to divide the exact size,  $d = |d_0 - d_m|$ , of the image interval,  $[d_0, d_m]$ , into a set,  $Y$ , containing  $d_c$  number of members:

$$(3.5) \quad \forall d_c \in \mathbb{N}, \ c > 0 \exists d \in \mathbb{R} : \quad floor(d/c) = d_c.$$

Apply the ruler convergence theorem (2.2):

$$(3.6) \quad \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c.$$

Combine equations 3.6 and 3.4:

$$(3.7) \quad d = \lim_{c \rightarrow 0} d_c \cdot c \quad \wedge \quad \sum_{i=1}^n s_i = \lim_{c \rightarrow 0} d_c \cdot c \quad \Rightarrow \quad d = \sum_{i=1}^n s_i. \quad \square$$

**THEOREM 3.3.** *Euclidean (shortest) distance,  $d$ , is the size of the distance interval,  $[d_0, d_m]$ , mapping to a set of disjoint domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d^2 = \sum_{i=1}^n s_i^2, \quad d = |d_0 - d_m|, \quad s_i = |a_i - b_i|.$$

The formal Coq-based theorem and proof in the file euclidrelations.v is “Euclidean\_distance.”

**PROOF.**

Use the ruler (2.1) to divide the exact size,  $s_i = |a_i - b_i|$ , of each of the domain intervals,  $[a_i, b_i]$ , into a set,  $x_i$ , containing  $p_i$  number of subintervals and apply the definition of the countable distance range (3.1), where each domain set has a corresponding distance set containing the same  $p_i$  number of members.

$$(3.8) \quad \forall i \ n \in \mathbb{N}, \quad i \in [1, n], \quad s_i \in \mathbb{R}, \quad \exists c > 0 : \quad \text{floor}(s_i/c) = p_i = |x_i| = |y_i|.$$

Apply the rule of product to the largest number of domain-to-distance set combinations, where all  $p_i$  number of domain set members,  $x_i$ , combine with (map to) each of the  $p_i$  number of members in the distance set,  $y_i$ :

$$(3.9) \quad \sum_{i=1}^n |y_i| \cdot |x_i| = \sum_{i=1}^n p_i^2.$$

Choose the equality case of the Cauchy-Schwartz inequality:

$$(3.10) \quad \sum_{i=1}^n p_i^2 \leq (\sum_{i=1}^n p_i)^2 \Rightarrow \exists p_i : \sum_{i=1}^n p_i^2 = (\sum_{i=1}^n p_i)^2$$

Choose the equality case of the countable distance range definition (3.1) and square both sides ( $x = y \Rightarrow f(x) = f(y)$ ):

$$(3.11) \quad \sum_{i=1}^n |y_i| = \sum_{i=1}^n p_i \geq d_c \Rightarrow \exists p_i, d_c : \sum_{i=1}^n p_i = d_c \\ \Rightarrow \exists p_i, d_c : (\sum_{i=1}^n p_i)^2 = d_c^2.$$

Combine equations 3.10 and 3.11:

$$(3.12) \quad \exists p_i : \sum_{i=1}^n p_i^2 = (\sum_{i=1}^n p_i)^2 \quad \wedge \quad \exists p_i, d_c : (\sum_{i=1}^n p_i)^2 = d_c^2 \\ \Rightarrow \exists p_i, d_c : \sum_{i=1}^n p_i^2 = d_c^2.$$

Multiply both sides of equation 3.12 by  $c^2$  and apply the ruler convergence theorem:

$$(3.13) \quad s_i = \lim_{c \rightarrow 0} p_i \cdot c \quad \wedge \quad \sum_{i=1}^n (p_i \cdot c)^2 = (d_c \cdot c)^2 \\ \Rightarrow \sum_{i=1}^n s_i^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2.$$

Use the ruler to divide the exact size,  $d = |d_0 - d_m|$ , of the image interval,  $[d_0, d_m]$  into a set,  $Y$ , containing  $d_c$  number of members:

$$(3.14) \quad \forall d_c \in \mathbb{N}, \quad c > 0 \exists d \in \mathbb{R} : \text{floor}(d/c) = d_c.$$

Apply the ruler convergence theorem (2.2) and then square both sides:

$$(3.15) \quad \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c \Rightarrow d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2.$$

Combine equations 3.15 and 3.13:

$$(3.16) \quad d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 \wedge \sum_{i=1}^n s_i^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 \Rightarrow d^2 = \sum_{i=1}^n s_i^2. \quad \square$$

**3.1. Triangle inequality.** The definition of a metric in real analysis is based on the triangle inequality,  $\mathbf{d}(\mathbf{u}, \mathbf{w}) \leq \mathbf{d}(\mathbf{u}, \mathbf{v}) + \mathbf{d}(\mathbf{v}, \mathbf{w})$ , that has been intuitively motivated by the triangle [Gol76]. Applying the ruler (2.1), and ruler convergence theorem (2.2) to the definition of a countable distance range (3.1) yields the real-valued triangle inequality:

$$(3.17) \quad d_c = |Y| = |\bigcup_{i=1}^2 y_i| \leq \sum_{i=1}^2 |y_i| \wedge \\ d_c = \text{floor}(\mathbf{d}(\mathbf{u}, \mathbf{w})/c) \wedge |y_1| = \text{floor}(\mathbf{d}(\mathbf{u}, \mathbf{v})/c) \wedge |y_2| = \text{floor}(\mathbf{d}(\mathbf{v}, \mathbf{w})/c) \\ \Rightarrow \mathbf{d}(\mathbf{u}, \mathbf{w}) = \lim_{c \rightarrow 0} d_c \cdot c \leq \sum_{i=1}^2 \lim_{c \rightarrow 0} |y_i| \cdot c = \mathbf{d}(\mathbf{u}, \mathbf{v}) + \mathbf{d}(\mathbf{v}, \mathbf{w}).$$

The other metric space properties:  $\mathbf{d}(\mathbf{u}, \mathbf{w}) = 0 \Leftrightarrow u = w$ ,  $\mathbf{d}(\mathbf{u}, \mathbf{w}) = \mathbf{d}(\mathbf{w}, \mathbf{u})$ , and  $\mathbf{d}(\mathbf{u}, \mathbf{w}) \geq 0$  also follow from the countable distance range definition.

#### 4. Size (length/area/volume)

The combinatorial relations between all members in set  $x_1$  to each member of set  $x_2$  results in the Cartesian product of  $|x_1| \cdot |x_2|$  number of combinations. This section will use the ruler (2.1) and ruler convergence theorem (2.2) to prove that the Cartesian product of the same-sized subintervals of intervals converges to the product of interval sizes as the subinterval converges to zero. The first step is to define a set-based, countable size measure as the Cartesian product of disjoint domain set members.

DEFINITION 4.1. Countable size (length/area/volume) measure,  $S_c$ :

$$\forall i \ n \in \mathbb{N}, \quad i \in [1, n], \quad x_i \subseteq X, \quad \sum_{i=1}^n |x_i| = |\bigcup_{i=1}^n x_i|, \quad \wedge \quad S_c = \prod_{i=1}^n |x_i|.$$

THEOREM 4.2. *Euclidean size (length/area/volume),  $S$ , is the size of an image interval,  $[v_0, v_m]$ , corresponding to a set of disjoint intervals:  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$S = \prod_{i=1}^n s_i, \quad S = |v_0 - v_m|, \quad s_i = |a_i - b_i|, \quad i \in [1, n], \quad i, n \in \mathbb{N}.$$

The Coq-based theorem and proof in the file euclidrelations.v is “Euclidean\_size.”

PROOF.

Use the ruler (2.1) to divide the exact size,  $s_i = |a_i - b_i|$ , of each of the domain intervals,  $[a_i, b_i]$ , into a set,  $x_i$  of  $p_i$  number of subintervals.

$$(4.1) \quad \forall i \ n \in \mathbb{N}, \quad i \in [1, n], \quad c > 0 \quad \wedge \quad \text{floor}(s_i/c) = p_i = |x_i|.$$

Use the ruler (2.1) to divide the exact size,  $S = |v_0 - v_m|$ , of the image interval,  $[v_0, v_m]$ , into  $p_S^n$  subintervals. Every integer number,  $S_c$ , does **not** have an integer  $n^{\text{th}}$  root. However, for those cases where  $S_c$  does have an integer  $n^{\text{th}}$  root, there is a  $p_S^n$  that satisfies the definition a countable size measure,  $S_c$  (4.1). Notionally:

$$(4.2) \quad \forall p_S^n = S_c \in \mathbb{N}, \exists S \in \mathbb{R}, x_i : \text{floor}(S/c) = p_S^n = S_c = \prod_{i=1}^n |x_i| = \prod_{i=1}^n p_i.$$

Multiply both sides of equation 4.2 by  $c^n$  to get the ruler measures:

$$(4.3) \quad p_S^n = \prod_{i=1}^n p_i \quad \Rightarrow \quad (p_S \cdot c)^n = \prod_{i=1}^n (p_i \cdot c).$$

Apply the ruler convergence theorem (2.2) to equation 4.3:

$$(4.4) \quad S = \lim_{c \rightarrow 0} (p_S \cdot c)^n \quad \wedge \quad (p_S \cdot c)^n = \prod_{i=1}^n (p_i \cdot c) \quad \wedge \quad \lim_{c \rightarrow 0} (p_i \cdot c) = s_i \\ \Rightarrow \quad S = \lim_{c \rightarrow 0} (p_S \cdot c)^n = \prod_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \prod_{i=1}^n s_i. \quad \square$$

## 5. Ordered and symmetric geometries

The commutative property of the union and addition operations in the countable distance range principle (3.1) that generates the triangle inequality, Manhattan and Euclidean distances and the commutative property of the union and multiplication operations in the countable size principle (4.1) that generates length/area/volume allow a sequential (total) ordering of the disjoint domain sets (dimensions) to exist. And the commutative property also allows every dimension to be sequentially adjacent to any other dimension (herein, referred to as a symmetric geometry).

It will now be proved that satisfying both the total ordering and symmetry properties simultaneously limits distance and volume to a cyclic set of at most three dimensions.

DEFINITION 5.1. Ordered geometry:

$$\forall i \ n \in \mathbb{N}, \ i \in [1, n-1], \ \forall x_i \in \{x_1, \dots, x_n\}, \\ \text{successor } x_i = x_{i+1} \ \wedge \ \text{predecessor } x_{i+1} = x_i.$$

where each  $x_i \in \{x_1, \dots, x_n\}$  is a set of subintervals of a real-valued domain interval (dimension).

DEFINITION 5.2. Symmetric geometry (every member is sequentially adjacent to every other member):

$$\forall i \ j \ n \in \mathbb{N}, \ \forall x_i \ x_j \in \{x_1, \dots, x_n\}, \ \text{successor } x_i = x_j \ \wedge \ \text{predecessor } x_j = x_i.$$

THEOREM 5.3. An ordered and symmetric geometry is a cyclic set.

$$\text{successor } x_n = x_1 \ \wedge \ \text{predecessor } x_1 = x_n.$$

The theorem and formal Coq-based proof is “ordered\_symmetric\_is\_cyclic,” which is located in the file threed.v.

PROOF. The property of order (5.1) defines unique successors and predecessors for all members except for the successor of  $x_n$  and the predecessor of  $x_1$ . From the properties of a symmetric geometry (5.2):

$$(5.1) \quad i = n \ \wedge \ j = 1 \ \wedge \ \text{successor } x_i = x_j \Rightarrow \text{successor } x_n = x_1.$$

$$(5.2) \quad i = n \ \wedge \ j = 1 \ \wedge \ \text{predecessor } x_j = x_i \Rightarrow \text{predecessor } x_1 = x_n. \quad \square$$

THEOREM 5.4. An ordered and symmetric geometry is limited to at most 3 members.

The Coq-based lemmas and proofs in the file threed.v are:

**Lemmas:** `adj111`, `adj122`, `adj212`, `adj123`, `adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and `not_all_mutually_adjacent_gt_3`.

The following proof uses Horn clauses (a subset of first-order logic) that uses unification and resolution. Horn clauses make it clear which facts satisfy a proof goal.

**PROOF.**

Because a symmetric and ordered set is a cyclic set (5.3), the successors and predecessors are cyclic:

**DEFINITION 5.5.** Successor of  $m$  is  $n$ :

$$(5.3) \quad \text{Successor}(m, n, \text{setsize}) \leftarrow (m = \text{setsize} \wedge n = 1) \vee (m + 1 \leq \text{setsize}).$$

**DEFINITION 5.6.** Predecessor of  $m$  is  $n$ :

$$(5.4) \quad \text{Predecessor}(m, n, \text{setsize}) \leftarrow (m = 1 \wedge n = \text{setsize}) \vee (m - 1 \geq 1).$$

**DEFINITION 5.7.** Adjacent: member  $m$  is sequentially adjacent to member  $n$  (required for a "symmetric" set (5.2)), if the cyclic successor of  $m$  is  $n$  or the cyclic predecessor of  $m$  is  $n$ . Notionally:

$$(5.5) \quad \text{Adjacent}(m, n, \text{setsize}) \leftarrow \text{Successor}(m, n, \text{setsize}) \vee \text{Predecessor}(m, n, \text{setsize}).$$

Every member is adjacent to every other member, where  $\text{setsize} \in \{1, 2, 3\}$ :

$$(5.6) \quad \text{Adjacent}(1, 1, 1) \leftarrow \text{Successor}(1, 1, 1) \leftarrow (1 = 1 \wedge 1 = 1).$$

$$(5.7) \quad \text{Adjacent}(1, 2, 2) \leftarrow \text{Successor}(1, 2, 2) \leftarrow (1 + 1 \leq 2).$$

$$(5.8) \quad \text{Adjacent}(2, 1, 2) \leftarrow \text{Successor}(2, 1, 2) \leftarrow (2 = 2 \wedge 1 = 1).$$

$$(5.9) \quad \text{Adjacent}(1, 2, 3) \leftarrow \text{Successor}(1, 2, 3) \leftarrow (1 + 1 \leq 2).$$

$$(5.10) \quad \text{Adjacent}(2, 1, 3) \leftarrow \text{Predecessor}(2, 1, 3) \leftarrow (2 - 1 \geq 1).$$

$$(5.11) \quad \text{Adjacent}(3, 1, 3) \leftarrow \text{Successor}(3, 1, 3) \leftarrow (3 = 3 \wedge 1 = 1).$$

$$(5.12) \quad \text{Adjacent}(1, 3, 3) \leftarrow \text{Predecessor}(1, 3, 3) \leftarrow (1 = 1 \wedge 3 = 3).$$

$$(5.13) \quad \text{Adjacent}(2, 3, 3) \leftarrow \text{Successor}(2, 3, 3) \leftarrow (2 + 1 \leq 3).$$

$$(5.14) \quad \text{Adjacent}(3, 2, 3) \leftarrow \text{Predecessor}(3, 2, 3) \leftarrow (3 - 1 \geq 1).$$

Must prove that for all  $\text{setsize} > 3$ , there exist non-adjacent members. For example, the first and third members are not adjacent:

$$(5.15) \quad \forall \text{setsize} > 3: \quad \neg \text{Successor}(1, 3, \text{setsize}) \\ \leftarrow \text{Successor}(1, 2, \text{setsize}) \leftarrow (1 + 1 \leq \text{setsize}).$$

That is, 2 is the only successor of 1 for all  $\text{setsize} > 3$ , which implies 3 is not a successor of 1 for all  $\text{setsize} > 3$ .

$$(5.16) \quad \forall \text{setsize} > 3: \quad \neg \text{Predecessor}(1, 3, \text{setsize}) \\ \leftarrow \text{Predecessor}(1, n, \text{setsize}) \leftarrow (1 = 1 \wedge n = \text{setsize}).$$



That is,  $n = \text{setsize}$  is the only predecessor of 1 for all  $\text{setsize} > 3$ , which implies 3 is not a predecessor of 1 for all  $\text{setsize} > 3$ .

$$(5.17) \quad \forall \text{setsize} > 3: \quad \neg \text{Adjacent}(1, 3, \text{setsize}) \\ \leftarrow \neg \text{Successor}(1, 3, \text{setsize}) \wedge \neg \text{Predecessor}(1, 3, \text{setsize}). \quad \square$$

## 6. Summary

Applying some very simple real analysis, in the form of the ruler measure (2.1) and ruler convergence proof (2.2), to a set of real-valued domain intervals and an image interval yields some new insights into geometry and physics.

- (1) Discrete, combinatorial relations converge to the continuous, bijective relations: triangle inequality, Manhattan distance, Euclidean distance and volume.
- (2) The ruler measure is used to derive Euclidean distance and volume from many-to-many relationships between the elements of countable sets, where one domain element can map to many image elements. But, a function only allows each domain value to map to one image value, which can not describe many-to-many relationships, where each domain value can map to many image values. Therefore, Euclidean integrals and measures, like the Lebesgue and Hausdorff measures, that use distance, area, and volume functions, and use those functions as primitives, are **not** capable of deriving the set-based relationships that converge to continuous, bijective distance and volume functions.
- (3) Ruler measure-based proofs expose the difference between distance and size (length/area/volume) measures: Distance is the combinatorial relation between the members of each disjoint domain set and members of a corresponding image (distance) set. In contrast, volume is a combinatorial relation between the members of disjoint domain sets.
- (4) Applying the ruler measure to the countable distance range (3.1) provides the insight that all notions of distance are based on the principle that for each disjoint domain set there exists a corresponding distance set containing the same number of members, where the distance sets sometimes intersect:
  - (a) The countable distance range principle converges to the real-valued triangle inequality (3.1), which is the basis for the definition of metric space. The other properties of metric space also come from the countable distance range principle. Therefore, a function is not a distance metric unless it satisfies the more fundamental countable distance range (3.1).
  - (b) The upper bound of the countable distance range converging to Manhattan distance (3.2) provides the insight that the largest (longest) monotonic distance path is the case of the minimum number of combinations (mappings) per distance set, where each member in the  $i^{th}$  distance set combines with (maps to) only one member in the  $i^{th}$  domain set, which is the case of disjoint distance sets.
  - (c) The lower bound of the countable distance range converging to Euclidean distance (3.3) provides the insight that the smallest (shortest)

possible monotonic distance path is the case of the maximum number of combinations (mappings) per distance set, where each of the  $p_i$  number of members in the  $i^{th}$  distance set combine with (map to) all  $p_i$  number of members in the  $i^{th}$  domain set, which is the case of the maximum allowed intersection of the distance sets.

- (d) All  $L^{p>2}$  norms generated from the countable distance range principle would require each member of the  $i^{th}$  distance set to map to (form a combination with) a member of the  $i^{th}$  domain set more than once, which would be over-counting the number of possible mappings (combinations). Therefore,  $L^{p>2}$  norms are not valid distance measures. Measure theories have not provided this over-counting insight into  $L^{p>2}$  norms.
- (e) Euclidean distance (3.3) was derived from a combinatorial relation without any notions of side, angle, or shape. A parametric variable relating the sizes of two domain intervals can be easily derived using calculus (Taylor series) and the Euclidean distance equation to generate the arc sine and arc cosine functions of the parametric variable (arc angle). In other words, the notions of side and angle are derived from Euclidean distance, which is the reverse perspective of classical geometry [Joy98] [Loo68] [Ber88] and axiomatic foundations for geometry [Bir32] [Hil80] [TG99].
- (5) Applying the ruler measure and ruler convergence proof to the countable size definition (4.1) allows a proof that the Cartesian product of same-sized subintervals of real-valued intervals converges to the product of the interval sizes (Euclidean volume):
  - (a) Euclidean size (length/area/volume) was derived from a combinatorial relation without notions of sides, angles, and shape.
- (6) The set-based definitions of countable distance range converging to the triangle inequality, Manhattan distance, Euclidean distance and countable size converging to Euclidean volume provides a self-contained foundation under real analysis, calculus, and measure theory by not having to import those equations from Euclidean geometry as definitions and not having to rely on external geometry notions of side, angle, shape, side-angle-side relation, etc.
- (7) The set-based relations of countable distance range (3.1) and countable size (4.1) that generate the real-valued triangle inequality, Manhattan distance, Euclidean distance, and volume equations have a symmetry property (5.2) that limits distance and volume to a cyclic set (5.3) of three dimensions (5.4). This symmetry property explains why only three dimensions of physical space can be observed.
- (8) All ordered and symmetric, higher-dimensional geometries collapse into hierarchical 2 or 3-dimensional geometries. The four-vector lengths common in physics are 2-dimensional Euclidean lengths that have been "flattened." For example, the spacetime four-vector length,  $d = \sqrt{(ct)^2 - (x^2 + y^2 + z^2)}$ , is sometimes expressed in a form like,  $d_2 = \sqrt{(ct)^2 - d_1^2}$ , where  $d_1 = \sqrt{x^2 + y^2 + z^2}$  and  $d_2 = d$ .

Applying the Euclidean distance proof (3.3) to the 2-dimensional Poincaré form (where  $c = 1$ ),  $t^2 = d_1^2 + d_2^2$ , provides the perspective

that  $d_1$  and  $d_2$  are lengths in two frames of reference (the lengths of two domain intervals) and the size of each time subinterval is the same size (same speed of light) in both frames of reference.

## References

- [Ber88] B. C. Berndt, *Ramanujan-100 years old (fashioned) or 100 years new (fangled)?*, The Mathematical Intelligencer **10** (1988), no. 3. ↑10
- [Bir32] G. D. Birkhoff, *A set of postulates for plane geometry (based on scale and protractors)*, Annals of Mathematics **33** (1932). ↑10
- [Coq15] Coq, *Coq proof assistant*, 2015. <https://coq.inria.fr/documentation>. ↑2
- [Gol76] R. R. Goldberg, *Methods of real analysis*, John Wiley and Sons, 1976. ↑1, 6
- [Hil80] D. Hilbert, *The foundations of geometry (2cd ed)*, Chicago: Open Court, 1980. <http://www.gutenberg.org/ebooks/17384>. ↑10
- [Joy98] D. E. Joyce, *Euclid's elements*, 1998. <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>. ↑10
- [Loo68] E. S. Loomis, *The pythagorean proposition*, NCTM, 1968. ↑10
- [TG99] A. Tarski and S. Givant, *Tarski's system of geometry*, The Bulletin of Symbolic Logic **5** (1999), no. 2, 175–214. ↑10

GEORGE VAN TREECK, 668 WESTLINE DR., ALAMEDA, CA 94501