

# The Two Set Relations Generating Geometry

George. M. Van Treeck

ABSTRACT. Where each countable domain set has a corresponding range set, countable distance is defined as the cardinal of the union of the range sets and countable volume as the cardinal of the set of  $n$ -tuples of members from the disjoint range sets. The countable distance and volume set operations applied to sets of size  $c$  subintervals of domain and range intervals generate the properties of metric space, all  $L_p$  norms (for example, Manhattan and Euclidean distance), and the volume equation as  $c$  goes to 0. The volume proof is used to derive Coulomb's charge force and Newton's gravity force equations without using other laws of physics or Gauss's divergence theorem. A symmetry constraint on a totally ordered set limits the set to at most 3 members, for example, 3 dimensions. All proofs are verified in Coq.

## CONTENTS

1. Introduction	1
2. Ruler measure and convergence	2
3. Distance	2
4. Euclidean Volume	6
5. Applications to physics	7
6. Insights and implications	10
References	12

## 1. Introduction

The definitions of metric space, Euclidean distance, and Euclidean area/volume in mathematical analysis [Gol76] [Rud76] are motivated by Euclidean geometry [Joy98] rather than derived from an abstract set and limit-based foundation. An abstract set and limit-based foundation exposes: the constraint between countable domain and range sets that makes a space flat or curved; the countable domain-to-range set mapping that makes Euclidean distance the smallest possible distance between two distinct points in flat space; the set operation and constraint generating the properties of metric space; etc.

Where each disjoint, countable domain set has a corresponding range set, countable distance is defined as the cardinal of the union of the range sets and countable volume as the cardinal of the set of  $n$ -tuples of members from disjoint range sets. The countable distance and volume set operations applied to the sets of same-sized, size  $c$ , subintervals of domain and range intervals generate the properties of metric space, the absolute value metric in  $\mathbb{R}$ , all  $L_p$  norms (Minkowski distances, for example, Manhattan and Euclidean distance) in  $\mathbb{R}^n: n > 1$ , and the volume equation as  $c \rightarrow 0$ . Some applications to physics are also shown.

All the proofs in this article have been verified using the Coq proof verification system [Coq15]. The formal proofs are in the Coq files, “euclidrelations.v” and “threed.v,” at: <https://github.com/treeck/RASRGeometry>.

## 2. Ruler measure and convergence

In this article, geometric relations are derived from the number of correspondences (mappings) between the set of size  $c$  subintervals in each interval (as  $c \rightarrow 0$ ). Derivatives and integrals in  $\mathbb{R}^n$  are bijective functions, which precludes, for example, a surjective mapping from a size  $c$  subinterval in a domain interval to multiple size  $c$  subintervals in a range interval. Therefore, derivatives and integrals are not well suited for deriving geometric relations from the full range of possible mappings between sets of size  $c$  subintervals.

A ruler (measuring stick) measures the size of each domain and a range interval *approximately* as the sum of the nearest integer number,  $p$ , of whole subintervals, where each subinterval has the *same* size,  $c$ . The ruler makes it easy to derive geometric relations from the number of possible mappings between the  $p_x$  number of subintervals in one interval and the  $p_y$  number of subintervals in another interval.

**DEFINITION 2.1.** Ruler measure,  $M$ :  $\forall c, s \in \mathbb{R}, [a, b] \subset \mathbb{R}, s = b - a \wedge c > 0 \wedge (p = \text{floor}(s/c) \vee p = \text{ceiling}(s/c)) \wedge M = \sum_{i=1}^p c = pc$ .

**THEOREM 2.2.** *Ruler convergence:*  $M = \lim_{c \rightarrow 0} pc = s$ .

The theorem, “limit\_c\_0\_M\_eq\_exact\_size,” and formal proof is in the Coq file, euclidrelations.v. The proof is trivial proof, but is included here for completeness.

**PROOF.** (epsilon-delta proof)

By definition of the floor function,  $\text{floor}(x) = \max(\{y : y \leq x, y \in \mathbb{Z}, x \in \mathbb{R}\})$ :

$$(2.1) \quad \forall c > 0, p = \text{floor}(s/c) \wedge 0 \leq |\text{floor}(s/c) - s/c| < 1 \Rightarrow 0 \leq |p - s/c| < 1.$$

Multiply all sides of inequality 2.1 by  $c$ :

$$(2.2) \quad \forall c > 0, \quad 0 \leq |p - s/c| < 1 \Rightarrow 0 \leq |pc - s| < |c|.$$

$$(2.3) \quad \forall \delta : |pc - s| < |c| = |c - 0| < \delta \\ \Rightarrow \quad \forall \epsilon = \delta : |c - 0| < \delta \wedge |pc - s| < \epsilon := M = \lim_{c \rightarrow 0} pc = s. \quad \square$$

The following is an example of ruler convergence for the interval,  $[0, \pi]$ :  $s = \pi - 0$ , and  $p = \text{floor}(s/c) \Rightarrow p \cdot c = 3.1_{c=10^{-1}}, 3.14_{c=10^{-2}}, 3.141_{c=10^{-3}}, \dots, \pi_{\lim_{c \rightarrow 0}}$ .

## 3. Distance

**Notation conventions:** Vertical bars around a set,  $|\{\dots\}|$ , or list,  $|\llbracket \dots \rrbracket|$ , indicates the cardinal (the number of members in the set or list).

**3.1. Countable distance.** Distance in one dimension is independent of distance in every other other dimension. Therefore, each disjoint domain set,  $x_i$ , has its own independent range set,  $y_i$ . And the space is flat, where  $|x_i| = |y_i|$ . The countable distance spanning the disjoint domain sets is the number of members,  $d_c$ , in the union range (distance) set:

DEFINITION 3.1. Countable distance,  $d_c$ , in flat space:

$$d_c = |\bigcup_{i=1}^n y_i| : \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i|.$$

It will be shown in the next subsections that the constraint,  $|x_i| = |y_i|$ , generates Manhattan and Euclidean distance at the boundaries (generates flat space/rectilinear distances). Generalizing distance and volume beyond flat space is shown in the last section of this article.

**3.2. Union-Sum Inequality.** The inequality,  $|\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i|$ , is used often in this article. The proof is trivial proof, but is included here for completeness.

The proof follows from the associative law of addition where the sum of set sizes is equal to the size of all the set members appended into a list and the commutative law of addition that allows sorting that list into a list of unique members (the *union* set) and a list of duplicates. For example,  $y_1 = \{a, b, c\}$  and  $y_2 = \{c, d, e\} \Rightarrow |\bigcup_{i=1}^2 y_i| = |\{a, b, c, d, e\}| = 5 < \sum_{i=1}^2 |y_i| = |[a, b, c, c, d, e]| = 6$ .

LEMMA 3.2. *Union-Sum Inequality:*  $|\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i|$ .

PROOF. A formal proof, `union_sum_inequality`, using sorting into a set of unique members (*union* set) and a list of duplicates, is in the file `euclidrelations.v`.

$$(3.1) \quad \begin{aligned} \sum_{i=1}^n |y_i| &= |\text{append}_{i=1}^n y_i| = |\text{sort}(\text{append}_{i=1}^n y_i)| \\ &= |\bigcup_{i=1}^n y_i| + |\text{duplicates}_{i=1}^n y_i|. \end{aligned}$$

$$(3.2) \quad \begin{aligned} |\bigcup_{i=1}^n y_i| + |\text{duplicates}_{i=1}^n y_i| &= \sum_{i=1}^n |y_i| \quad \wedge \quad |\text{duplicates}_{i=1}^n y_i| \geq 0 \\ &\Rightarrow |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i|. \quad \square \end{aligned}$$

**3.3. Countable distance range.** From the countable distance definition (3.1),  $d_c = |\bigcup_{i=1}^n y_i|$ , as the amount of intersection increases, more domain set members can map to a single range set member. Therefore, the countable distance,  $d_c$ , is a function of the total number of domain-to-range set member mappings.

Each domain set,  $x_i$  has its own independent range set,  $y_i$ . From the countable distance constraint (3.1), where  $|x_i| = |y_i| = p_i$ , the countable distance,  $d_c$ , ranges from a function of the sum of 1-1 correspondence mappings,  $d_c = f(\sum_{i=1}^n (1 \cdot |y_i|)) = f(\sum_{i=1}^n p_i)$ , to a function of the sum of all-to-each (Cartesian product) mappings,  $d_c = f(\sum_{i=1}^n (|x_i| \cdot |y_i|)) = f(\sum_{i=1}^n p_i^2)$ .

Applying the ruler (2.1) and ruler convergence theorem (2.2) to the smallest and largest total number of domain-to-range set mapping cases converges to the real-valued Manhattan and Euclidean distance relations.

### 3.4. Manhattan distance.

THEOREM 3.3. *Manhattan (largest) distance,  $d$ , is the size of the range interval,  $[d_0, d_m]$ , corresponding to a set of disjoint domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d = \sum_{i=1}^n s_i, \quad d = d_m - d_0, \quad s_i = b_i - a_i.$$

The formal proof, “taxicab\_distance,” is in the Coq file, euclidrelations.v.

PROOF.

From the countable distance definition (3.1) and the union-sum inequality (3.2), the largest possible countable distance,  $d_c$ , is the equality case:

$$(3.3) \quad d_c = |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i| \quad \wedge \quad |x_i| = |y_i| = p_i \quad \Rightarrow \quad d_c \leq \sum_{i=1}^n p_i \\ \Rightarrow \quad \exists p_i, d_c : d_c = \sum_{i=1}^n p_i.$$

Multiply both sides of equation 3.3 by  $c$  and take the limit:

$$(3.4) \quad d_c = \sum_{i=1}^n p_i \Rightarrow d_c \cdot c = \sum_{i=1}^n (p_i \cdot c) \Rightarrow \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c).$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to the definition of  $d$ :

$$(3.5) \quad d = d_m - d_0 \Rightarrow \exists c d : \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to each domain interval:

$$(3.6) \quad s_i = b_i - a_i \quad \wedge \quad \text{floor}(s_i/c) = |x_i| = |y_i| = p_i \Rightarrow \lim_{c \rightarrow 0} p_i \cdot c = s_i.$$

Combine equations 3.5, 3.4, 3.6:

$$(3.7) \quad d = \lim_{c \rightarrow 0} d_c \cdot c \quad \wedge \quad \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) \quad \wedge \\ \lim_{c \rightarrow 0} (p_i \cdot c) = s_i \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \sum_{i=1}^n s_i. \quad \square$$

### 3.5. Euclidean distance.

**THEOREM 3.4.** *Euclidean (smallest) distance,  $d$ , is the size of the range interval,  $[d_0, d_m]$ , corresponding to a set of disjoint domain intervals,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ , where:*

$$d^2 = \sum_{i=1}^n s_i^2, \quad d = d_m - d_0, \quad s_i = b_i - a_i.$$

The formal proof, “Euclidean\_distance,” is in the Coq file, euclidrelations.v.

PROOF.

Apply the rule of product to the largest number of domain-to-range set mappings, where all  $p_i$  number of range set members,  $y_i$ , map to each of the  $p_i$  number of members in the domain set,  $x_i$ , which is the Cartesian product,  $|y_i| \cdot |x_i|$ :

$$(3.8) \quad |x_i| = |y_i| = p_i \Rightarrow \sum_{i=1}^n |y_i| \cdot |x_i| = \sum_{i=1}^n p_i^2.$$

From the countable distance definition (3.1) and the union-sum inequality (3.2), choose the equality case:

$$(3.9) \quad d_c = |\bigcup_{i=1}^n y_i| \leq \sum_{i=1}^n |y_i| \quad \wedge \quad |x_i| = |y_i| = p_i \Rightarrow d_c \leq \sum_{i=1}^n p_i \\ \Rightarrow \quad \exists p_i, d_c : d_c = \sum_{i=1}^n p_i.$$

Square both sides of equation 3.9 ( $x = y \Leftrightarrow f(x) = f(y)$ ):

$$(3.10) \quad \exists p_i, d_c : d_c = \sum_{i=1}^n p_i \Leftrightarrow \exists p_i, d_c : d_c^2 = (\sum_{i=1}^n p_i)^2.$$

Apply the square of sum inequality,  $(\sum_{i=1}^n p_i)^2 \geq \sum_{i=1}^n p_i^2$ , to equation 3.10 and select the smallest area (the equality) case:

$$(3.11) \quad d_c^2 = (\sum_{i=1}^n p_i)^2 = \sum_{i=1}^n p_i \sum_{j=1}^n p_j \\ = \sum_{i=1}^n p_i^2 + \sum_{i=1}^n p_i \sum_{j=1, j \neq i}^n p_j \geq \sum_{i=1}^n p_i^2 \Rightarrow \exists p_i : d_c^2 = \sum_{i=1}^n p_i^2.$$

Multiply both sides of equation 3.11 by  $c^2$ , simplify, and take the limit.

$$(3.12) \quad d_c^2 = \sum_{i=1}^n p_i^2 \Rightarrow d_c^2 \cdot c^2 = \sum_{i=1}^n p_i^2 \cdot c^2 \Leftrightarrow (d_c \cdot c)^2 = \sum_{i=1}^n (p_i \cdot c)^2 \\ \Rightarrow \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) and square both sides:

$$(3.13) \quad \exists c d \in \mathbb{R} : \text{floor}(d/c) = d_c \Rightarrow d = \lim_{c \rightarrow 0} d_c \cdot c \Rightarrow d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2.$$

Apply the ruler (2.1) and ruler convergence theorem (2.2) to each domain interval:

$$(3.14) \quad s_i = b_i - a_i \wedge \text{floor}(s_i/c) = |x_i| = |y_i| = p_i \Rightarrow \lim_{c \rightarrow 0} p_i \cdot c = s_i.$$

Combine equations 3.13, 3.12, 3.14:

$$(3.15) \quad d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 \wedge \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2 \wedge \\ \lim_{c \rightarrow 0} (p_i \cdot c) = s_i \Rightarrow d^2 = \lim_{c \rightarrow 0} (d_c \cdot c)^2 = \sum_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c)^2 = \sum_{i=1}^n s_i^2. \quad \square$$

**3.6. Metric Space.** All distances,  $d(u, w)$ , satisfying the countable distance definition (3.1), where the ruler is applied, generates the properties of metric space. The formal proofs: triangle\_inequality, non\_negativity, identity\_of\_indiscernibles, and symmetry are in the Coq file, euclidrelations.v.

**THEOREM 3.5. Triangle Inequality:**  $d_c = |y_1 \cup y_2| \Rightarrow d(u, w) \leq d(u, v) + d(v, w)$ .

**PROOF.** Apply the ruler measure (2.1), the countable distance condition (3.1), union-sum inequality (3.2), and then ruler convergence (2.2).

$$(3.16) \quad \forall c > 0, d(u, w), d(u, v), d(v, w) : \\ |y_1| = \text{floor}(d(u, v)/c) \wedge |y_2| = \text{floor}(d(v, w)/c) \wedge \\ d_c = \text{floor}(d(u, w)/c) \wedge d_c = |y_1 \cup y_2| \leq |y_1| + |y_2| \\ \Rightarrow \text{floor}(d(u, w)/c) \leq \text{floor}(d(u, v)/c) + \text{floor}(d(v, w)/c) \\ \Rightarrow \text{floor}(d(u, w)/c) \cdot c \leq \text{floor}(d(u, v)/c) \cdot c + \text{floor}(d(v, w)/c) \cdot c \\ \Rightarrow \lim_{c \rightarrow 0} \text{floor}(d(u, w)/c) \cdot c \leq \lim_{c \rightarrow 0} \text{floor}(d(u, v)/c) \cdot c + \lim_{c \rightarrow 0} \text{floor}(d(v, w)/c) \cdot c \\ \Rightarrow d(u, w) \leq d(u, v) + d(v, w). \quad \square$$

**THEOREM 3.6. Non-negativity:**  $d_c = |y_1 \cup y_2| \Rightarrow d(u, w) \geq 0$ .

**PROOF.** By definition, a set always has a size (cardinal)  $\geq 0$ :

$$(3.17) \quad \forall c > 0, d(u, w) : \text{floor}(d(u, w)/c) = d_c \wedge d_c = |y_1 \cup y_2| \geq 0 \\ \Rightarrow \text{floor}(d(u, w)/c) = d_c \geq 0 \Rightarrow d(u, w) = \lim_{c \rightarrow 0} d_c \cdot c \geq 0. \quad \square$$

**THEOREM 3.7. Identity of Indiscernibles:**  $d(u, w) = 0$ .

**PROOF.** Apply the triangle inequality property (3.5):

$$(3.18) \quad \forall d(u, v) = d(v, w) = 0 \wedge d(u, w) \leq d(u, v) + d(v, w) \Rightarrow d(u, w) \leq 0.$$

Combine the non-negativity property (3.6) and the previous inequality (3.18):

$$(3.19) \quad d(u, w) \geq 0 \wedge d(u, w) \leq 0 \Leftrightarrow 0 \leq d(u, w) \leq 0 \Rightarrow d(u, w) = 0.$$

Combine the result of step 3.19 and the condition,  $d(u, v) = 0$ , in step 3.18.

$$(3.20) \quad d(u, w) = 0 \wedge d(u, v) = 0 \Rightarrow w = v.$$

Combine the condition,  $d(v, w) = 0$ , in step 3.18 and the result of step 3.20.

$$(3.21) \quad d(v, w) = 0 \wedge w = v \Rightarrow d(w, w) = 0. \quad \square$$

THEOREM 3.8. *Symmetry:*  $|x_i| = |y_i| \Rightarrow d(u, v) = d(v, u)$ .

PROOF.

The range of countable distances (3.3) is a function of domain-to-range set members, under the constraint,  $|x_i| = |y_i|$ :

$$(3.22) \quad |x_i| = |y_i| = p_i \quad \Rightarrow \quad d_c = f(\sum_{i=1}^n |x_i| \cdot |y_i|^q) = f(\sum_{i=1}^n p_i^{1+q}), \quad 0 \leq q \leq 1.$$

Applying the ruler to real-valued domain and range intervals, where  $s_i$  is the size of domain interval,  $[a_i, b_i]$ , generates the range of distances from Manhattan distance (3.3),  $d(x, y) = f(\sum_{i=1}^n s_i^1)$ , to Euclidean distance (3.4),  $d(x, y) = f(\sum_{i=1}^n s_i^2)$ . Generalizing:

$$(3.23) \quad \forall p : p \geq 1, \quad d(x, y) = f(\sum_{i=1}^n s_i^p).$$

Distance is a function of domain interval sizes,  $s_i$ . For all  $x, y \in \mathbb{R}$ , there are two domain interval cases:

*Case #1:  $n = 1$  domain interval:* In this case,  $d(x, y)$  is the distance from domain value  $x$  to domain value  $y$  in  $\mathbb{R}$ , which make  $x$  and  $y$  the boundary values of a domain interval having size,  $s : s = |x - y|$ . And applying equation 3.23, yields the absolute value metric:

$$(3.24) \quad d(x, y) = f(s^p) = f(|x - y|^p) \Rightarrow d(u, v) = f(|u - v|^p) = f(|v - u|^p) = d(v, u).$$

*Case #2:  $n = 2$  domain intervals:* In this case,  $d(x, y)$  is the distance in  $\mathbb{R}^2$ , where  $x$  and  $y$  are the sizes of two domain intervals,  $[a_1, b_1]$  and  $[a_2, b_2]$ , where  $s_1 = x = |a_1 - b_1|$  and  $s_2 = y = |a_2 - b_2|$ :

$$(3.25) \quad d(x, y) = f(s_1^p + s_2^p) = f(x^p + y^p) \\ \Rightarrow \quad d(u, v) = f(u^p + v^p) = f(v^p + u^p) = d(v, u). \quad \square$$

#### 4. Euclidean Volume

$\mathbb{R}^n$ , the Lebesgue measure, Riemann integral, and Lebesgue integral define (assume) area/volume to be the product of domain interval lengths. The goal here is to derive the area/volume equation from an abstract, set-based definition of volume (a product  $\sigma$ -algebra), without assuming the product of interval lengths.

DEFINITION 4.1. Countable Volume,  $v_c$ , is the cardinal of the set of n-tuples of members from countable, disjoint, range sets:

$$v_c = |\times_{i=1}^n y_i| : \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset.$$

THEOREM 4.2. *Euclidean volume,  $v$ , is length of the range interval,  $[v_0, v_m]$ , equal to product of domain interval lengths,  $\{[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]\}$ :*

$$v = \prod_{i=1}^n s_i, \quad v = v_m - v_0, \quad s_i = b_i - a_i.$$

The formal proof, “Euclidean\_volume,” is in the Coq file, euclidrelations.v.

PROOF.

Use the ruler (2.1) to partition each of the domain intervals,  $[a_i, b_i]$ , into a set,  $x_i$ , containing  $p_i$  number of subintervals.

$$(4.1) \quad \forall i \, n \in \mathbb{N}, \quad i \in [1, n], \quad c > 0 \quad \wedge \quad \text{floor}(s_i/c) = p_i = |x_i| = |y_i|.$$

Apply the ruler convergence theorem (2.2) to equation 4.1:

$$(4.2) \quad \text{floor}(s_i/c) = p_i \quad \Rightarrow \quad \lim_{c \rightarrow 0} (p_i \cdot c) = s_i.$$

Apply the associative law of multiplication to derive the countable volume (4.1) in terms of  $p_i$ :

$$(4.3) \quad v_c = |\times_{i=1}^n y_i| = \prod_{i=1}^n |y_i| \quad \wedge \quad |y_i| = p_i \quad \Rightarrow \quad v_c = \prod_{i=1}^n p_i.$$

Multiply both sides of equation 4.3 by  $c^n$ :

$$(4.4) \quad v_c \cdot c^n = (\prod_{i=1}^n p_i) \cdot c^n = \prod_{i=1}^n (p_i \cdot c).$$

$$(4.5) \quad \forall n, v_c \in \mathbb{N} \exists p \in \mathbb{R} : p^n = v_c \Rightarrow v_c \cdot c^n = p^n \cdot c^n = (p \cdot c)^n = \prod_{i=1}^n (p_i \cdot c).$$

Apply the ruler (2.1) and ruler convergence (2.2) to the range interval,  $[v_0, v_m]$  (where  $v = v_m - v_0$ ), and then combine with equations 4.5 and 4.2:

$$(4.6) \quad floor(v/c^n) = p^n \Rightarrow v = \lim_{c \rightarrow 0} (p \cdot c)^n = \prod_{i=1}^n \lim_{c \rightarrow 0} (p_i \cdot c) = \prod_{i=1}^n s_i. \quad \square$$

## 5. Applications to physics

**5.1. Coulomb's charge force.**  $q_1$  and  $q_2$ , are the sizes of two independent charge intervals, where each infinitesimal size  $c$  subinterval of a charge interval exerts a quantum force,  $m_{CA}c$ , on each size  $c$  subinterval of the other charge interval. The total force,  $F$ , is proportionate to the total number of forces (the Cartesian product of the infinitesimal size  $c$  components) multiplied times the quantum charge force,  $m_{CA}c$ . Applying the ruler,  $p_1 = floor(q_1/c)$  and  $p_2 = floor(q_2/c)$ , and the Cartesian product,  $p_1 \times p_2$ , of size  $c$  components yields:

$$(5.1) \quad F \propto m_{CA}c (\lim_{c \rightarrow 0} p_1 c \cdot \lim_{c \rightarrow 0} p_2 c) = m_{CA}c \int_0^{q_2} \int_0^{q_1} d^2c = m_{CA}c (q_1 q_2).$$

For every size,  $q$ , of a charge interval, another interval having size,  $r$ , can be defined, where  $r : q = (q_C/r_C)r$  and  $q_C/r_C$  is a unit-factoring conversion ratio.

$$(5.2) \quad \forall q_1, q_2 \geq 0 \exists q \in \mathbb{R} : q^2 = q_1 q_2 \quad \wedge \quad (q_C/r_C)r = q \Rightarrow ((q_C/r_C)r)^2 = q_1 q_2.$$

$$(5.3) \quad ((q_C/r_C)r)^2 = q_1 q_2 \quad \wedge \quad F \propto m_{CA}c (q_1 q_2) \\ \Rightarrow F \propto m_{CA}c ((q_C/r_C)r)^2 = m_{CA}c (q_1 q_2) \\ \Rightarrow F = m_{CA}c = (m_{CA}c r_C^2 / q_C^2) q_1 q_2 / r^2 = k_C q_1 q_2 / r^2.$$

where  $k_C = m_{CA}c r_C^2 / q_C^2$  corresponds to the SI units:  $Nm^2C^{-2}$ . And multiplying both sides of equation 5.3 by  $xy$ :

$$(5.4) \quad \forall x, y \geq 1, m_0 = x \cdot m_C, a = y \cdot a_C, d = r/(x \cdot y) \Rightarrow F = m_0 a = k_C q_1 q_2 / d^2.$$

**5.2. Newton's gravity force equation.**  $m_1$  and  $m_2$ , are the sizes of two independent mass intervals, where each infinitesimal size  $c$  subinterval of a mass interval exerts a quantum force,  $m_{GA}c$ , on each size  $c$  subinterval of the other mass interval. The total force,  $F$ , is proportionate to the total number of forces (the Cartesian product of the size  $c$  components) multiplied times the quantum gravity force,  $m_{GA}c$ . Applying the ruler,  $p_1 = floor(m_1/c)$  and  $p_2 = floor(m_2/c)$ , and the Cartesian product,  $p_1 \times p_2$ , of size  $c$  components yields:

$$(5.5) \quad F \propto m_{GA}c (\lim_{c \rightarrow 0} p_1 c \cdot \lim_{c \rightarrow 0} p_2 c) = m_{GA}c \int_0^{m_2} \int_0^{m_1} d^2c = m_{GA}c (m_1 m_2).$$

For every size,  $m$ , of a mass interval, another interval having size,  $r$ , can be defined, where  $r : m = (m_G/r_G)r$  and  $m_G/r_G$  is a unit-factoring conversion ratio.

$$(5.6) \quad \forall m_1, m_2 \geq 0 \exists m \in \mathbb{R} : m^2 = m_1 m_2 \quad \wedge \quad (m_G/r_G)r = m \\ \Rightarrow ((m_G/r_G)r)^2 = m_1 m_2.$$

$$\begin{aligned}
(5.7) \quad ((m_G/r_G)r)^2 &= m_1 m_2 \quad \wedge \quad F \propto m_G a_G (m_1 m_2) \\
&\Rightarrow \quad F \propto m_G a_G ((m_G/r_G)r)^2 = m_G a_G (m_1 m_2) \\
&\Rightarrow \quad F = m_G a_G = (a_G r_G^2 / m_G) m_1 m_2 / r^2.
\end{aligned}$$

$$\begin{aligned}
(5.8) \quad \exists t_G \in \mathbb{R} : r_G / t_G^2 &= a_G \quad \wedge \quad F = (a_G r_G^2 / m_G) m_1 m_2 / r^2 \\
&\Rightarrow \quad F = (r_G^3 / m_G t_G^2) m_1 m_2 / r^2 = G m_1 m_2 / r^2,
\end{aligned}$$

where  $G = r_G^3 / m_G t_G^2$  corresponds to the SI units:  $m^3 kg^{-1} s^{-2}$ . And multiplying both sides of equation 5.8 by  $xy$ :

$$(5.9) \quad \forall x, y \geq 1, m_0 = x \cdot m_G, a = y \cdot a_G, d = r / (x \cdot y) \Rightarrow F = m_0 a = G m_1 m_2 / d^2.$$

**5.3. Spacetime equations.** For any Euclidean distance interval having size,  $r$ , a interval having size,  $t$ , can be defined, where  $r = (r_c/t_c)t = ct$ , and  $r_c/t_c = c$  is a unit-factoring conversion ratio.

Applying the ruler to two intervals,  $[0, d_1]$  and  $[0, d_2]$ , in two inertial (independent, non-accelerating) frames of reference, the distance (and time) spanning the two domain intervals converges to a range of distances (and times) from Manhattan (3.3) to Euclidean distance (3.4).

$$\begin{aligned}
(5.10) \quad r^2 &= d_1^2 + d_2^2 \quad \wedge \quad r = (r_c/t_c)t = ct \quad \Rightarrow \quad (ct)^2 = d_1^2 + d_2^2 \\
&\Leftrightarrow \quad d_1^2 = (ct)^2 - (x^2 + y^2 + z^2),
\end{aligned}$$

where  $d_2^2 = x^2 + y^2 + z^2$ , which is one form of Minkowski's well-known flat spacetime interval equation [Bru17]. And, the time dilation and length contraction equations also follow directly by dividing both sides of  $(ct)^2 = d_1^2 + d_2^2$  by  $t^2$  and using  $v = d/t$ .

**5.4. 3 dimensional balls.** Countable distance,  $d_c = |\bigcup_{i=1}^n y_i|$ , (3.1), countable volume,  $d_c = |\times_{i=1}^n y_i|$ , (4.1), Manhattan distance (3.3), Euclidean distance (3.4), and volume (4.2) requires a *strict total order* ( $i = 1$  to  $n$ ) of a set of intervals/dimensions. And the commutative properties of union, addition, and multiplication allow sequencing through each interval (dimension) in every possible order. **Note** that “jumping” from member 1 to member  $m$  of a set requires calculating an offset that is an implicit traversal of successor/predecessor relations. Therefore, “strict” sequencing (no jumping over other members) via the successor and predecessor relations of a strict totally ordered set in every possible order requires each set member to be sequentially adjacent (either a successor or predecessor) to every other member, herein referred to as a symmetric geometry.

It will now be proved that the constraint (coexistence) of symmetry on a sequentially ordered set defines a cyclic set that contains at most 3 members, in this case, 3 dimensions of ordered and symmetric distance and volume. If there are higher dimension of space, then the cyclic property prevents sequencing from the 3 lower, cyclic set of dimensions to any higher dimensions.

**DEFINITION 5.1.** Ordered geometry:

$$\begin{aligned}
\forall i \in \mathbb{N}, \quad i \in [1, n-1], \quad \forall x_i \in \{x_1, \dots, x_n\}, \\
\text{successor } x_i = x_{i+1} \quad \wedge \quad \text{predecessor } x_{i+1} = x_i.
\end{aligned}$$



DEFINITION 5.2. Symmetric geometry (every set member is sequentially adjacent to every other member):

$$\forall i \, j \, n \in \mathbb{N}, \, \forall x_i \, x_j \in \{x_1, \dots, x_n\}, \, \text{successor } x_i = x_j \Leftrightarrow \text{predecessor } x_j = x_i.$$

THEOREM 5.3. *An ordered and symmetric set is a cyclic set.*

$$i = n \wedge j = 1 \Rightarrow \text{successor } x_n = x_1 \wedge \text{predecessor } x_1 = x_n.$$

The formal proof, “ordered\_symmetric\_is\_cyclic,” is in the Coq file, threaded.v.

PROOF. A total order (5.1) defines unique successors and predecessors for all set members except for the successor of  $x_n$  and the predecessor of  $x_1$ . Therefore, the only member that can be a successor of  $x_n$ , without creating a contradiction, is  $x_1$ . And the only member that can be a predecessor of  $x_1$ , without creating a contradiction, is  $x_n$ . From the properties of a symmetric geometry (5.2):

$$(5.11) \quad i = n \wedge j = 1 \wedge \text{successor } x_i = x_j \Rightarrow \text{successor } x_n = x_1.$$

Applying the definition of a symmetric geometry (5.2) to conclusion 5.11:

$$(5.12) \quad \text{successor } x_i = x_j \Rightarrow \text{predecessor } x_j = x_i \Rightarrow \text{predecessor } x_1 = x_n. \quad \square$$

THEOREM 5.4. *An ordered and symmetric set is limited to at most 3 members.*

The lemmas and formal proofs in the Coq file threaded.v are:

Lemmas: adj111, adj122, adj212, adj123, adj133, adj233, adj213, adj313, adj323, and not\_all\_mutually\_adjacent\_gt\_3.

The following proof uses Horn clauses (a subset of first-order logic) that uses unification and resolution. Horn clauses make it clear which facts satisfy a proof goal.

PROOF.

It was proved that an ordered and symmetric set is a cyclic set (5.3). In other words, the successors and predecessors of an ordered and symmetric set are cyclic:

DEFINITION 5.5. Cyclic successor of  $m$  is  $n$ :

$$(5.13) \quad \text{Successor}(m, n, \text{setsize}) \leftarrow (m = \text{setsize} \wedge n = 1) \vee (n = m + 1 \leq \text{setsize}).$$

DEFINITION 5.6. Cyclic predecessor of  $m$  is  $n$ :

$$(5.14) \quad \text{Predecessor}(m, n, \text{setsize}) \leftarrow (m = 1 \wedge n = \text{setsize}) \vee (n = m - q \geq 1).$$

DEFINITION 5.7. Adjacent: member  $m$  is sequentially adjacent to member  $n$  if the cyclic successor of  $m$  is  $n$  or the cyclic predecessor of  $m$  is  $n$ . Notionally:

$$(5.15) \quad \text{Adjacent}(m, n, \text{setsize}) \leftarrow \text{Successor}(m, n, \text{setsize}) \vee \text{Predecessor}(m, n, \text{setsize}).$$

Prove that every member is adjacent to every other member, where  $\text{setsize} \in \{1, 2, 3\}$ :

$$(5.16) \quad \text{Adjacent}(1, 1, 1) \leftarrow \text{Successor}(1, 1, 1) \leftarrow (m = \text{setsize} \wedge n = 1).$$

$$(5.17) \quad \text{Adjacent}(1, 2, 2) \leftarrow \text{Successor}(1, 2, 2) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.18) \quad \text{Adjacent}(2, 1, 2) \leftarrow \text{Successor}(2, 1, 2) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(5.19) \quad \text{Adjacent}(1, 2, 3) \leftarrow \text{Successor}(1, 2, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.20) \quad \text{Adjacent}(2, 1, 3) \leftarrow \text{Predecessor}(2, 1, 3) \leftarrow (n = m - q \geq 1).$$

$$(5.21) \quad \text{Adjacent}(3, 1, 3) \leftarrow \text{Successor}(3, 1, 3) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(5.22) \quad \text{Adjacent}(1, 3, 3) \leftarrow \text{Predecessor}(1, 3, 3) \leftarrow (m = 1 \wedge n = \text{setsize}).$$

$$(5.23) \quad \text{Adjacent}(2, 3, 3) \leftarrow \text{Successor}(2, 3, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(5.24) \quad \text{Adjacent}(3, 2, 3) \leftarrow \text{Predecessor}(3, 2, 3) \leftarrow (n = m - q \geq 1).$$

Must prove that for all  $\text{setsize} > 3$ , there exist non-adjacent members. For example, the first and third members are not ( $\neg$ ) adjacent:

$$(5.25) \quad \forall \text{setsize} > 3 : \quad \neg \text{Successor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Successor}(1, 2, \text{setsize} > 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

That is, member 2 is the only successor of member 1 for all  $\text{setsize} > 3$ , which implies member 3 is not a successor of member 1 for all  $\text{setsize} > 3$ .

$$(5.26) \quad \forall \text{setsize} > 3 : \quad \neg \text{Predecessor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Predecessor}(1, \text{setsize}, \text{setsize} > 3) \leftarrow (m = 1 \wedge n = \text{setsize} > 3).$$

That is, member  $n = \text{setsize} > 3$  is the only predecessor of member 1, which implies member 3 is not a predecessor of member 1 for all  $\text{setsize} > 3$ .

$$(5.27) \quad \forall \text{setsize} > 3 : \quad \neg \text{Adjacent}(1, 3, \text{setsize} > 3) \\ \leftarrow \neg \text{Successor}(1, 3, \text{setsize} > 3) \wedge \neg \text{Predecessor}(1, 3, \text{setsize} > 3). \quad \square$$

That is, for all  $\text{setsize} > 3$ , some elements are not sequentially adjacent to every other element (not symmetric).

## 6. Insights and implications

- (1) In flat space, where  $|x_i| = |y_i| = p_i$ , the largest possible number of domain-to-range set mappings is where each domain set,  $x_i$ , member maps to every member of the corresponding range set,  $y_i$ , which is the Cartesian product of the number of domain and range set members:  $d_c = f(\sum_{i=1}^n |x_i| \cdot |y_i|) = f(\sum_{i=1}^n p_i^2)$ . The largest possible number of domain-to-range set mappings is the set-based reason that Euclidean distance (3.4) is the smallest possible distance between two distinct points (in flat space).
- (2) Generalizing the countable distance and volume constraint,  $|x_i| = |y_i|$ , to  $|x_i| = |y_i|^q$ ,  $q \geq 0$ , generates all the  $L^p$  norms (Minkowski distances),  $\|L\|_p = (\sum_{i=1}^n s_i^p)^{1/p}$ . For example, using the same proof pattern as for Euclidean distance (3.4):  $|y_i| = p_i \Rightarrow |x_i| = p_i^q \Rightarrow \sum_{i=1}^n |x_i| \cdot |y_i| = \sum_{i=1}^n p_i^q \cdot p_i = \sum_{i=1}^n p_i^{q+1} \leq d_c^{q+1} \dots d = (\sum_{i=1}^n s_i^{q+1})^{1/(q+1)}$ .
- (3) The curvature of a space around a point is typically measured in terms of second order differential equations, e.g., the Laplacian. A set-based measure of the amount of curvature is how far  $q$  deviates from the value, 1, in the countable distance and volume constraint,  $|x_i| = |y_i|^q$ .
- (4)  $-1 \leq q < 0$  would generate  $L^p$  norms that do not satisfy the metric space triangle inequality property. But, such values of  $q$  might occur in probability and physics, for example, where there are more nondeterministic outcomes than domain set members.

- (5) Manhattan distance and Euclidean volume are both cases of disjoint range sets,  $\bigcap_{i=1}^n y_i = \emptyset$ , in flat space (where  $|x_i| = |y_i|$ ):

$$d_c = |\bigcup_{i=1}^n y_i| : \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset.$$

$$v_c = |\times_{i=1}^n y_i| : \quad \bigcap_{i=1}^n x_i = \emptyset \quad \wedge \quad |x_i| = |y_i| \quad \wedge \quad \bigcap_{i=1}^n y_i = \emptyset.$$

- (6) There are combinatorial relationships between countable sets of subintervals of intervals in statistics, probability, physics, etc., where the ruler is an applicable tool. For example, applying the ruler (2.1) and ruler convergence (2.2) to the Cartesian product of same-sized, infinitesimal charge forces and mass forces allowed deriving Coulomb's charge force (5.1) and Newton's gravity force (5.5) equations in a few steps each, without using other laws of physics or Gauss's divergence theorem.
- (7) **The Proportionate Interval Principle:** The derivations of the charge force, gravity force, and spacetime equations shows that all Euclidean distance intervals having a size,  $r$ , have proportionately sized intervals of other types:  $r = (r_C/q_C)q = (r_G/m_G)m = (r_c/t_c)t = ct$ , where the conversion ratios are for unit-factor analysis.
- (a) Some versions of the charge constant, vacuum magnetic permeability constant, fine structure constant, etc. contain the value  $4\pi$  because the creators assumed flux divergence on the surface of a sphere,  $4\pi r^2$ . Using Occam's razor, the mapping of rectangular geometric area to rectangular charge and mass areas provides more parsimonious derivations of the inverse square law, charge, and gravity force equations than flux divergence. Therefore, those versions of the constants containing the value  $4\pi$  might be incorrect.
- (b)  $(r_G/m_G)m \cdot ct = r^2 \Rightarrow m = (m_G/r_G c)r^2/t$ . For a constant mass,  $m$ , as the distance,  $r$ , to the mass decreases, then time,  $t$ , must also decrease (time slows down), which agrees with relativity theory and observation. Also,  $(r_G/m_G)m \cdot (ct)^2 = r^3 \Rightarrow E = mc^2 = (m_G/r_G)r^3/t^2 = (m_G/r_G)rv^2$ .  
Likewise, for charge,  $q = (q_C/r_C c)r^2/t \dots$
- (c) If there are quantum values of charge,  $q_C$ , and mass,  $m_G$ , then there are quantum distances (wavelengths),  $r_C$  and  $r_G$ , where the charge and gravity forces do not exist (are not defined) at smaller distances.
- (d) A countable set of values has measure 0. Therefore, there is no proportion relationship of a countable set of values to distance. Therefore, a countable set of state value changes with respect to time are independent of distance. For example, the change in the spin values of two quantum entangled electrons and the change in polarization of two quantum entangled photons are independent of the distance between the entangled particles.
- (8) Any higher dimensions of space not being sequentially reachable from the lower 3 dimensions because the lower 3 dimensions are a cyclic set is a more parsimonious explanation of not seeing any higher dimensions than the higher dimensions being rolled into infinitesimal balls, which requires an additional explanation of what causes the higher dimensions to be rolled up and additional equations describing the rollups.

- (9) If there are higher dimensions of space, then there is most likely an ordered and symmetric set of three members, each member being an ordered and symmetric set of 3 dimensions (three boxes), yielding a total of 9 ordered and symmetric dimensions of space. For example, suppose charge, mass, and time are Euclidean distances in the three boxes that correspond to a geometric Euclidean distance:  $r = f(q, m, t)$ .
- (10) Each ordered and symmetric ball can have at most 3 ordered and symmetric dimensions of discrete states of the same type, for example, a set of 3 binary values, 1 and -1, indicating vector orientation.
- (11) Each dimension of discrete physical states can have at most 3 ordered and symmetric discrete state values, which allows  $3 \cdot 3 \cdot 3 = 27$  possible combinations of discrete values of the same type per ball, for example, spin values: -1, 0, 1 per orthogonal plane in the ball.
- (12) Each of the 3 possible ordered and symmetric dimensions of discrete physical states could contain an unordered collection (bag) of discrete state values. Bags (of states) are non-deterministic. For example, every time an unordered binary state is measured, there is a 50 percent chance of having one of the binary values.
- (13) Where balls having a quantum radius intersect, an arithmetic of the interactions of the discrete states with respect to time needs to be developed. The interaction of the discrete states associated with intersecting balls with respect to time might result in what we perceive as motion, waves, particles, spin, polarization, work, force, mass, charge, etc.

## References

- [Bru17] P. Bruskiewich, *A very simple introduction to special relativity: Part two - four vectors, the lorentz transformation and group velocity (the new mathematics for the millions book 38)*, Pythagoras Publishing, 2017. ↑8
- [CG15] W. Conradie and V. Goranko, *Logic and discrete mathematics*, Wiley, 2015. ↑
- [Coq15] Coq, *Coq proof assistant*, 2015. <https://coq.inria.fr/documentation>. ↑2
- [Gol76] R. R. Goldberg, *Methods of real analysis*, John Wiley and Sons, 1976. ↑1
- [Joy98] D. E. Joyce, *Euclid's elements*, 1998. <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>. ↑1
- [Rud76] W. Rudin, *Principles of mathematical analysis*, McGraw Hill Education, 1976. ↑1

GEORGE VAN TREECK, 668 WESTLINE DR., ALAMEDA, CA 94501