

# The Set Properties Generating Geometry and Physics

George. M. Van Treeck

ABSTRACT. Volume and the Minkowski distances/Lp norms (e.g., Manhattan and Euclidean distance) are derived from a set and limit-based foundation without referencing the primitives and relations of geometry. Sequencing a strict linearly ordered set in all n-at-a-time permutations via successor/predecessor relations is a cyclic set of at most 3 members. Therefore, all other interval lengths have different types from a cyclic set of 3 distance interval lengths. Unit-factoring ratios between different types of interval lengths and the set proofs provide simpler derivations of the spacetime, Newton's gravity, Coulomb's charge force, Planck-Einstein, quantum-relativity gravity equations and corresponding constants. All proofs are verified in Coq.

## CONTENTS

1. Introduction	1
2. Ruler measure and convergence	2
3. Volume	3
4. Distance	4
5. Applications to physics	8
6. Insights and implications	10
References	12

## 1. Introduction

Mathematical analysis can construct differential calculus from a set and limit-based foundation without referencing the primitives and relations of Euclidean geometry, like straight line, angle, shape, etc., which provides a more rigorous foundation to calculus. But volume in the Riemann integral, Lebesgue integral, measure theory and distance in the vector magnitude and metric space criteria are definitions motivated by Euclidean geometry. [Gol76] [Rud76] Here, volume and distance are motivated and derived from a set and limit-based foundation.

A well-known set-based motivation of Euclidean volume is the set of Cartesian product n-tuples:  $v_c = \prod_{i=1}^n |x_i|$ , where  $|x_i|$  is the cardinal of the countable, disjoint

set,  $x_i$ . But, where each  $x_i$  is a set of subintervals of an interval,  $[a_i, b_i] \subset \mathbb{R}$ , and  $s_i = b_i - a_i$ , there have been no proofs that  $v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i$ , hence, volume has been defined rather than derived. In this article,  $v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i$  is proved. Non-Euclidean volumes have the forms,  $v_c = \prod_{i=1}^n \alpha_i |x_i|$  and  $v = \prod_{i=1}^n \beta_i s_i$ , where  $\alpha_i$  and  $\beta_i$  are functions returning scalar values  $\neq 1$ .

N-volumes of the same class,  $n$ , can be measured as a function of the average domain set size:  $\forall \beta_i, s_i \exists d : d = (\sum_{i=1}^n \beta_i s_i) / n \Rightarrow v = d^n = \prod_{i=1}^n \beta_i s_i$ . Where  $\alpha_i = \beta_i = 1$ , it will be proved that  $v_c = \sum_{j=1}^m v_{c_i} \Rightarrow d^n = \sum_{i=1}^m d_i^n$ .  $d$  is the  $L_p$  norm (Minkowski distance), which will be proved to imply the metric space criteria.

Sequencing a set,  $\{x_1 \cdots x_n\}$ , from  $i = 1$  to  $n$ , is a strict linear (total) order that set theory defines in terms of successor and predecessor functions. If a set can be sequenced in all n-at-a-time orders, then a “symmetry” constraint is required, where every set member is either a successor or predecessor to every other set member. A strict linearly ordered and symmetric set will be proved to be a cyclic set, where  $n \leq 3$ .

Therefore, if  $\{x, y, z\}$  is a strict linearly ordered and symmetric set of 3 “distance” dimensions, then a fourth dimension,  $t$ , must have a different type (is a member of different set). Definite integrals divide domain intervals into the same number of subintervals, where each subinterval of a distance domain interval maps to a proportionate-sized subinterval of some other type of domain interval, which is expressed by a unit-factoring ratio, for example, *meters/second*.

Simpler and shorter derivations of the: spacetime, Newton’s gravity, Coulomb’s charge force, Planck-Einstein, quantum-relativity gravity equations, and their corresponding constants are provided using the ratios,  $r = (r_c/t_c)t = (r_c/m_c)m = (r_c/q_c)q$ , combined with the results of the volume, distance, and 3D proofs. Impacts on Einstein’s field equations are also discussed.

All the proofs in this article have been verified using the Coq proof verification system [Coq23]. The formal proofs are in the Coq files, “euclidrelations.v” and “thread.v,” at: <https://github.com/treeck/RASRGeometry>.

## 2. Ruler measure and convergence

Derivatives and integrals divide each domain interval into the same number of subintervals, where the size of the subintervals are proportionate to the size of the containing domain interval, which precludes using derivatives and integrals to directly express many-to-many (Cartesian product mappings) between same-sized, size  $\kappa$ , subintervals in different-sized domain intervals. Further, derivatives and integrals that define Euclidean volume and distance cannot be used to derive Euclidean volume and distance. Therefore, thinking outside of the derivative/integral box is required.

A ruler (measuring stick) measures the size of each interval *approximately* as the sum of the nearest integer number,  $p$ , of size  $\kappa$  subintervals. The ruler is both an inner and outer measure of an interval.

**DEFINITION 2.1.** Ruler measure,  $M = \sum_{i=1}^p \kappa = p\kappa$ , where  $\forall [a, b] \subset \mathbb{R}$ ,  $s = b - a \wedge 0 < \kappa \leq 1 \wedge (p = \text{floor}(s/\kappa) \vee p = \text{ceiling}(s/\kappa))$ .

**THEOREM 2.2.** *Ruler convergence:*  $M = \lim_{\kappa \rightarrow 0} p\kappa = s$ .

The formal proof, “limit\_c\_0\_M.eq\_exact\_size,” is in the file, euclidrelations.v.

PROOF. (epsilon-delta proof)

By definition of the floor function,  $\text{floor}(x) = \max(\{y : y \leq x, y \in \mathbb{Z}, x \in \mathbb{R}\})$ :

$$(2.1) \quad \forall \kappa > 0, p = \text{floor}(s/\kappa) \wedge 0 \leq |\text{floor}(s/\kappa) - s/\kappa| < 1 \Rightarrow |p - s/\kappa| < 1.$$

Multiply both sides of inequality 2.1 by  $\kappa$ :

$$(2.2) \quad \forall \kappa > 0, |p - s/\kappa| < 1 \Rightarrow |p\kappa - s| < |\kappa| = |\kappa - 0|.$$

$$(2.3) \quad \begin{aligned} \forall \epsilon = \delta \quad \wedge \quad |p\kappa - s| < |\kappa - 0| < \delta \\ \Rightarrow \quad |\kappa - 0| < \delta \quad \wedge \quad |p\kappa - s| < \epsilon \quad := \quad M = \lim_{\kappa \rightarrow 0} p\kappa = s. \quad \square \end{aligned}$$

The following is an example of ruler convergence for the interval,  $[0, \pi]$ :  $s = \pi - 0$ , and  $p = \text{floor}(s/\kappa) \Rightarrow p \cdot \kappa = 3.1_{\kappa=10^{-1}}, 3.14_{\kappa=10^{-2}}, 3.141_{\kappa=10^{-3}}, \dots, \pi_{\lim_{\kappa \rightarrow 0}}$ .

LEMMA 2.3.  $\forall n \geq 1, 0 < \kappa < 1 \Rightarrow \lim_{\kappa \rightarrow 0} \kappa^n = \lim_{\kappa \rightarrow 0} \kappa$ .

PROOF. The formal proof, “lim.c.to.n.eq.lim.c,” is in the Coq file, euclidrelations.v.

$$(2.4) \quad n \geq 1 \quad \wedge \quad 0 < \kappa < 1 \Rightarrow 0 < \kappa^n < \kappa \Rightarrow |\kappa - \kappa^n| < |\kappa| = |\kappa - 0|.$$

$$(2.5) \quad \begin{aligned} \forall \epsilon = \delta \quad \wedge \quad |\kappa - \kappa^n| < |\kappa - 0| < \delta \\ \Rightarrow \quad |\kappa - 0| < \delta \quad \wedge \quad |\kappa - \kappa^n| < \delta = \epsilon \quad := \quad \lim_{\kappa \rightarrow 0} \kappa^n = 0. \end{aligned}$$

$$(2.6) \quad \lim_{\kappa \rightarrow 0} \kappa^n = 0 \quad \wedge \quad \lim_{\kappa \rightarrow 0} \kappa = 0 \Rightarrow \lim_{\kappa \rightarrow 0} \kappa^n = \lim_{\kappa \rightarrow 0} \kappa. \quad \square$$

### 3. Volume

DEFINITION 3.1. A countable n-volume is the number of ordered combinations (n-tuples),  $v_c$ , of the members of  $n$  number of disjoint, countable domain sets,  $x_i$ :

$$(3.1) \quad \exists n \in \mathbb{N}, v_c \in \{0, \mathbb{N}\}, x_i \in \{x_1, \dots, x_n\} : \bigcap_{i=1}^n x_i = \emptyset \wedge v_c = \prod_{i=1}^n |x_i|.$$

THEOREM 3.2. *Euclidean volume*,

$$(3.2) \quad \begin{aligned} \forall [a_i, b_i] \in \{[a_1, b_1], \dots, [a_n, b_n]\}, [v_a, v_b] \subset \mathbb{R}, s_i = b_i - a_i, v = v_b - v_a : \\ v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i. \end{aligned}$$

The formal proof, “Euclidean.volume,” is in the Coq file, euclidrelations.v.

PROOF.

$$(3.3) \quad v_c = \prod_{i=1}^n |x_i| \Leftrightarrow v_c \kappa = (\prod_{i=1}^n |x_i|) \kappa \Leftrightarrow \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i|) \kappa.$$

Apply the ruler (2.1) and ruler convergence (2.2) to equation 3.3:

$$(3.4) \quad \exists v, \kappa \in \mathbb{R} : v_c = \text{floor}(v/\kappa) \Rightarrow v = \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i|) \kappa.$$

Apply lemma 2.3 to equation 3.4:

$$(3.5) \quad v = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i|) \kappa = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i|) \kappa^n = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i| \kappa).$$

Apply the ruler (2.1) and ruler convergence (2.2) to  $s_i$ :

$$(3.6) \quad \exists s_i, \kappa \in \mathbb{R} : \text{floor}(s_i/\kappa) = |x_i| \Rightarrow \lim_{\kappa \rightarrow 0} (|x_i| \kappa) = s_i.$$

$$(3.7) \quad v = \lim_{\kappa \rightarrow 0} (\prod_{i=1}^n |x_i| \kappa) \wedge \lim_{\kappa \rightarrow 0} (|x_i| \kappa) = s_i \Rightarrow v = \prod_{i=1}^n s_i \quad \square$$

**THEOREM 3.3.** *Sum of volumes:*

$$(3.8) \quad \forall x_{i,j} \in \{x_{i_1}, \dots, x_{i_m}\} = x_i : v_c = \prod_{i=1}^n |x_i| \quad \wedge \quad v_{c_j} = \prod_{i=1}^n |x_{i,j}| \quad \wedge \\ v_c = \sum_{j=1}^m v_{c_j} \quad \Rightarrow \quad \exists s_i, s_{i,j} \in \mathbb{R} : \prod_{i=1}^n s_i = \sum_{j=1}^m (\prod_{i=1}^n s_{i,j}).$$

The formal proof, “sum\_of\_volumes,” is in the Coq file, euclidrelations.v.

**PROOF.** From the Euclidean volume theorem (3.2):

$$(3.9) \quad v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i \quad \wedge \quad v_{c_j} = \prod_{i=1}^n |x_{i,j}| \Rightarrow v_j = \prod_{i=1}^n s_{i,j}.$$

Apply the ruler (2.1) and ruler convergence (2.2):

$$(3.10) \quad \exists v, v_j, \kappa \in R : \quad v_c = \text{floor}(v/\kappa) \quad \wedge \quad v_{c_j} = \text{floor}(v_j/\kappa) \\ \Rightarrow \quad v = \lim_{\kappa \rightarrow 0} v_c \kappa \quad \wedge \quad v_i = \lim_{\kappa \rightarrow 0} v_{c_j} \kappa.$$

$$(3.11) \quad v_c = \sum_{j=1}^m v_{c_j} \quad \Leftrightarrow \quad v = \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} (\sum_{j=1}^m v_{c_j}) \kappa.$$

Apply lemma 2.3 to equation 3.11:

$$(3.12) \quad \lim_{\kappa \rightarrow 0} \kappa^n = \lim_{\kappa \rightarrow 0} \kappa \quad \wedge \quad v = \lim_{\kappa \rightarrow 0} (\sum_{j=1}^m v_{c_j}) \kappa \quad \wedge \quad v_i = \lim_{\kappa \rightarrow 0} v_{c_j} \kappa \\ \Rightarrow \quad v = \lim_{\kappa \rightarrow 0} (\sum_{j=1}^m v_{c_j}) \kappa^n = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_j} \kappa) = \sum_{j=1}^m v_j.$$

$$(3.13) \quad v = \prod_{i=1}^n s_i \quad \wedge \quad v_j = \prod_{i=1}^n s_{i,j} \quad \wedge \quad v = \sum_{j=1}^m v_j \\ \Rightarrow \quad \prod_{i=1}^n s_i = \sum_{j=1}^m \prod_{i=1}^n s_{i,j}. \quad \square$$

## 4. Distance

**DEFINITION 4.1.** The average countable domain set (distance) size,  $\alpha d_c$ , of a countable n-volume, n-volume,  $v_c = (\alpha d_c)^n = \sum_{i=1}^m (\alpha d_{c_i})^n = \sum_{j=1}^m v_{c_i}$ , where  $\alpha$  is a function returning a scalar value.

### 4.1. Minkowski distance ( $L_p$ norm).

**THEOREM 4.2.** *Minkowski distance ( $L_p$  norm):*

$$v_c = \prod_{i=1}^n |x_i| = \sum_{j=1}^m (\prod_{i=1}^n |x_{i,j}|) = \sum_{j=1}^m v_{c_i} \quad \Rightarrow \quad d^n = \sum_{i=1}^m d_i^n.$$

The formal proof, “Minkowski\_distance,” is in the Coq file, euclidrelations.v.

**PROOF.** From the sum of volumes proof (3.3), where all subintervals of all intervals are the same size,  $\kappa$ :

$$(4.1) \quad \prod_{i=1}^n |x_i| = \sum_{j=1}^m (\prod_{i=1}^n |x_{i,j}|) \quad \Rightarrow \quad \prod_{i=1}^n s_i = \sum_{j=1}^m (\prod_{i=1}^n s_{i,j})$$

$$(4.2) \quad \forall s_i, s_{i,j} \in \mathbb{R} \exists d, d_i \in \mathbb{R} : d = (\sum_{i=1}^n s_i)/n \quad \wedge \quad d_i = (\sum_{j=1}^m s_{i,j})/n \\ \Rightarrow \quad d^n = \prod_{i=1}^n s_i = \sum_{j=1}^m (\prod_{i=1}^n s_{i,j}) = \sum_{i=1}^m d_i^n. \quad \square$$

**4.2. Distance inequality.** The formal proof, distance\_inequality, is in the Coq file, euclidrelations.v.

**THEOREM 4.3.** *Distance inequality*

$$\forall n \in \mathbb{N}, v_a, v_b \geq 0 : (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}.$$

**PROOF.** Expand  $(v_a^{1/n} + v_b^{1/n})^n$  using the binomial expansion:

$$(4.3) \quad \forall v_a, v_b \geq 0 : v_a + v_b \leq v_a + v_b + \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^{n-i} (v_b^{1/n})^i + \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^i (v_b^{1/n})^{n-i} = (v_a^{1/n} + v_b^{1/n})^n.$$

Take the  $n^{th}$  of both sides of the inequality 4.3:

$$(4.4) \quad \forall v_a, v_b \geq 0, n \in \mathbb{N} : v_a + v_b \leq (v_a^{1/n} + v_b^{1/n})^n \Rightarrow (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}. \quad \square$$

**4.3. Distance sum inequality.** The formal proof, distance\_sum\_inequality, is in the Coq file, euclidrelations.v.

**THEOREM 4.4.** *Distance sum inequality*

$$\forall m, n \in \mathbb{N}, a_i, b_i \geq 0 : (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}.$$

**PROOF.** Apply the distance inequality (4.3):

$$(4.5) \quad \forall m, n \in \mathbb{N}, v_a, v_b \geq 0 : v_a = \sum_{i=1}^m a_i^n \quad \wedge \quad v_b = \sum_{i=1}^m b_i^n \quad \wedge \\ (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \quad \Rightarrow \quad ((\sum_{i=1}^m a_i^n) + (\sum_{i=1}^m b_i^n))^{1/n} = \\ (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad \square$$

**4.4. Metric Space.** All Minkowski distances ( $L_p$  norms) have the properties of metric space.

The formal proofs: triangle\_inequality, symmetry, non\_negativity, and identity\_of\_indiscernibles are in the Coq file, euclidrelations.v.

**THEOREM 4.5.** *Triangle Inequality:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, w) \leq d(u, v) + d(v, w).$$

**PROOF.**  $\forall p \geq 1, k > 1, u = s_1, w = s_2, v = w/k:$

$$(4.6) \quad (u^p + w^p)^{1/p} \leq ((u^p + w^p) + 2v^p)^{1/p} = ((u^p + v^p) + (v^p + w^p))^{1/p}.$$

Apply the distance inequality (4.3) to the inequality 4.6:

$$(4.7) \quad (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\ \wedge \quad v_a = u^p + v^p \quad \wedge \quad v_b = v^p + w^p \\ \Rightarrow \quad (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} \\ \Rightarrow \quad d(u, w) = (u^p + w^p)^{1/p} \leq \\ (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \quad \square$$

**THEOREM 4.6.** *Symmetry:*  $d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, v) = d(v, u).$

**PROOF.** By the commutative law of addition:

$$(4.8) \quad \forall p : p \geq 1, d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} = (s_1^p + s_2^p)^{1/p} \\ \Rightarrow d(u, v) = (u^p + v^p)^{1/p} = (v^p + u^p)^{1/p} = d(v, u). \quad \square$$

THEOREM 4.7. *Non-negativity:*  $d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, w) \geq 0$ .

PROOF. By definition, the length of an interval is always  $\geq 0$ :

$$(4.9) \quad \forall [a_1, b_1], [a_2, b_2], \quad u = b_1 - a_1, v = b_2 - a_2, \quad \Rightarrow \quad u \geq 0, v \geq 0.$$

$$(4.10) \quad p \geq 1, \quad u, v \geq 0 \quad \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} \geq 0. \quad \square$$

THEOREM 4.8. *Identity of Indiscernibles:*  $d(u, u) = 0$ .

PROOF. From the non-negativity property (4.7):

$$(4.11) \quad d(u, w) \geq 0 \quad \wedge \quad d(u, v) \geq 0 \quad \wedge \quad d(v, w) \geq 0 \\ \Rightarrow \quad \exists d(u, w) = d(u, v) = d(v, w) = 0.$$

$$(4.12) \quad d(u, w) = d(v, w) = 0 \quad \Rightarrow \quad u = v.$$

$$(4.13) \quad d(u, v) = 0 \quad \wedge \quad u = v \quad \Rightarrow \quad d(u, u) = 0. \quad \square$$

#### 4.5. The properties limiting a set to at most 3 members.

DEFINITION 4.9. Totally ordered set:

$$\forall i \in \mathbb{N}, \quad i \in [1, n-1], \quad \forall x_i \in \{x_1, \dots, x_n\}, \\ \text{successor } x_i = x_{i+1} \quad \wedge \quad \text{predecessor } x_{i+1} = x_i.$$

DEFINITION 4.10. Symmetry (every set member is sequentially adjacent to every other member):

$$\forall i, j, n \in \mathbb{N}, \quad \forall x_i, x_j \in \{x_1, \dots, x_n\}, \quad \text{successor } x_i = x_j \Leftrightarrow \text{predecessor } x_j = x_i.$$

THEOREM 4.11. *A strict linearly ordered and symmetric set is a cyclic set.*

$$i = n \quad \wedge \quad j = 1 \quad \Rightarrow \quad \text{successor } x_n = x_1 \quad \wedge \quad \text{predecessor } x_1 = x_n.$$

The formal proof, “ordered\_symmetric\_is\_cyclic,” is in the Coq file, `threed.v`.

PROOF. A total order (4.9) assigns a unique label to each set member and assigns unique successors and predecessors for all set members except for the successor of  $x_n$  and the predecessor of  $x_1$ . Therefore, the only member that can be a successor of  $x_n$ , without creating a contradiction, is  $x_1$ . And the only member that can be a predecessor of  $x_1$ , without creating a contradiction, is  $x_n$ . Applying the symmetry property (4.10):

$$(4.14) \quad i = n \quad \wedge \quad j = 1 \quad \wedge \quad \text{successor } x_i = x_j \quad \Rightarrow \quad \text{successor } x_n = x_1.$$

Applying the definition of the symmetry property (4.10) to conclusion 4.14:

$$(4.15) \quad \text{successor } x_i = x_j \quad \Rightarrow \quad \text{predecessor } x_j = x_i \quad \Rightarrow \quad \text{predecessor } x_1 = x_n. \quad \square$$

THEOREM 4.12. *An ordered and symmetric set is limited to at most 3 members.*

The formal proofs in the Coq file `threed.v` are:

**Lemmas:** `adj111`, `adj122`, `adj212`, `adj123`, `adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and `not_all_mutually_adjacent_gt_3`.

The following proof uses Horn clauses (a subset of first order logic), which makes it clear which facts satisfy a proof goal.

PROOF.

It was proved that an ordered and symmetric set is a cyclic set (4.11).

DEFINITION 4.13. (Cyclic) Successor of  $m$  is  $n$ :

$$(4.16) \quad \text{Successor}(m, n, \text{setsize}) \leftarrow (m = \text{setsize} \wedge n = 1) \vee (n = m + 1 \leq \text{setsize}).$$

DEFINITION 4.14. (Cyclic) Predecessor of  $m$  is  $n$ :

$$(4.17) \quad \text{Predecessor}(m, n, \text{setsize}) \leftarrow (m = 1 \wedge n = \text{setsize}) \vee (n = m - 1 \geq 1).$$

DEFINITION 4.15. Adjacent: member  $m$  is sequentially adjacent to member  $n$  if the successor of  $m$  is  $n$  or the predecessor of  $m$  is  $n$ . Notionally:

$$(4.18) \quad \text{Adjacent}(m, n, \text{setsize}) \leftarrow \text{Successor}(m, n, \text{setsize}) \vee \text{Predecessor}(m, n, \text{setsize}).$$

Prove that every member is adjacent to every other member, where  $\text{setsize} \in \{1, 2, 3\}$ :

$$(4.19) \quad \text{Adjacent}(1, 1, 1) \leftarrow \text{Successor}(1, 1, 1) \leftarrow (m = \text{setsize} \wedge n = 1).$$

$$(4.20) \quad \text{Adjacent}(1, 2, 2) \leftarrow \text{Successor}(1, 2, 2) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(4.21) \quad \text{Adjacent}(2, 1, 2) \leftarrow \text{Successor}(2, 1, 2) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(4.22) \quad \text{Adjacent}(1, 2, 3) \leftarrow \text{Successor}(1, 2, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(4.23) \quad \text{Adjacent}(2, 1, 3) \leftarrow \text{Predecessor}(2, 1, 3) \leftarrow (n = m - 1 \geq 1).$$

$$(4.24) \quad \text{Adjacent}(3, 1, 3) \leftarrow \text{Successor}(3, 1, 3) \leftarrow (n = \text{setsize} \wedge m = 1).$$

$$(4.25) \quad \text{Adjacent}(1, 3, 3) \leftarrow \text{Predecessor}(1, 3, 3) \leftarrow (m = 1 \wedge n = \text{setsize}).$$

$$(4.26) \quad \text{Adjacent}(2, 3, 3) \leftarrow \text{Successor}(2, 3, 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

$$(4.27) \quad \text{Adjacent}(3, 2, 3) \leftarrow \text{Predecessor}(3, 2, 3) \leftarrow (n = m - 1 \geq 1).$$

Member 2 is the only successor of member 1 for all  $\text{setsize} > 3$ , which implies member 3 is not ( $\neg$ ) a successor of member 1 for all  $\text{setsize} > 3$ :

$$(4.28) \quad \neg \text{Successor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Successor}(1, 2, \text{setsize} > 3) \leftarrow (n = m + 1 \leq \text{setsize}).$$

Member  $n = \text{setsize} > 3$  is the only predecessor of member 1, which implies member 3 is not ( $\neg$ ) a predecessor of member 1 for all  $\text{setsize} > 3$ :

$$(4.29) \quad \neg \text{Predecessor}(1, 3, \text{setsize} > 3) \\ \leftarrow \text{Predecessor}(1, \text{setsize}, \text{setsize} > 3) \leftarrow (m = 1 \wedge n = \text{setsize} > 3).$$

For all  $\text{setsize} > 3$ , some elements are not ( $\neg$ ) sequentially adjacent to every other element (not symmetric):

$$(4.30) \quad \neg \text{Adjacent}(1, 3, \text{setsize} > 3) \\ \leftarrow \neg \text{Successor}(1, 3, \text{setsize} > 3) \wedge \neg \text{Predecessor}(1, 3, \text{setsize} > 3). \quad \square$$

## 5. Applications to physics

From the 3D proof (4.12), dividing a set of domain intervals into the same number of subintervals, a 3-dimensional distance subinterval length,  $r$ , maps to proportionately sized subinterval lengths of other types,  $t$ ,  $m$ , and  $q$ , where:

$$(5.1) \quad r = (r_c/t_c)t = (r_c/m_c)m = (r_c/q_c)q.$$

**5.1. Spacetime equations.** From the volume proof (3.2), in a local Euclidean space, two disjoint distance intervals,  $[0, r]$  and  $[0, r']$ , define a 2-volume (area). From the Minkowski distance proof (4.2), the distance interval lengths,  $r$  and  $r'$ , are functions of two areas,  $v$  and  $v'$  having the sizes,  $v = r^2$  and  $v' = r'^2$ . And  $\forall r \geq r' \exists r_\nu \in \mathbb{R} : r^2 = r'^2 + r_\nu^2$ . Combined with the 3D proof (4.12):

$$(5.2) \quad \exists \mu, \nu \in \mathbb{R} : r = \mu t \quad \wedge \quad r_\nu = \nu t \quad \wedge \quad \forall r \geq r' \exists r_\nu \in \mathbb{R} : r^2 = r'^2 + r_\nu^2 \\ \Rightarrow (\mu t)^2 = r'^2 + (\nu t)^2 \quad \Rightarrow \quad r' = \sqrt{(\mu t)^2 - (\nu t)^2} = \mu t \sqrt{1 - (\nu/\mu)^2}.$$

Local (proper) distance,  $r'$ , contracts relative to coordinate distance,  $r$ , as  $\nu \rightarrow \mu$ :

$$(5.3) \quad r' = \mu t \sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad \mu t = r \quad \Rightarrow \quad r' = r \sqrt{1 - (\nu/\mu)^2}.$$

From equation 5.2, coordinate length,  $t$ , dilates relative to local length,  $t'$ , as  $\nu \rightarrow \mu$ :

$$(5.4) \quad \mu t = r' / \sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad r' = \mu t' \quad \Rightarrow \quad t = t' / \sqrt{1 - (\nu/\mu)^2}.$$

Using  $r^2 = r'^2 + r_\nu^2$  from equation 5.2, where  $r_\nu$  is a 3-dimensional distance, one form of the flat Minkowski's spacetime event interval is:

$$(5.5) \quad dr^2 = dr'^2 + dr_\nu^2 \quad \wedge \quad dr_\nu^2 = dx_1^2 + dx_2^2 + dx_3^2 \quad \wedge \quad d(\mu t) = dr \\ \Rightarrow \quad dr'^2 = d(\mu t)^2 - dx_1^2 - dx_2^2 - dx_3^2.$$

**5.2. Newton's gravity force and the constant,  $G$ .** From equation 5.1:

$$(5.6) \quad \forall m_1, m_2, m, r \in \mathbb{R} : m_1 m_2 = m^2 \quad \wedge \quad m = (m_c/r_c)r \\ \Rightarrow \quad m_1 m_2 = ((m_c/r_c)r)^2 \quad \Rightarrow \quad (r_c/m_c)^2 m_1 m_2 / r^2 = 1.$$

$$(5.7) \quad r = (r_c/t_c)t = ct \quad \Rightarrow \quad mr = (m_c/r_c)(ct)^2 \quad \Rightarrow \quad ((r_c/m_c)/c^2)mr/t^2 = 1.$$

$$(5.8) \quad ((r_c/m_c)/c^2)mr/t^2 = 1 \quad \wedge \quad (r_c/m_c)^2 m_1 m_2 / r^2 = 1 \\ \Rightarrow \quad F := mr/t^2 = ((r_c/m_c)c^2)m_1 m_2 / r^2 = G m_1 m_2 / r^2,$$

where Newton's constant,  $G = (r_c/m_c)c^2$ , conforms to the SI units:  $m^3 \cdot kg^{-1} \cdot s^{-2}$ .

**5.3. Coulomb's charge force and constant.** From equation 5.1:

$$(5.9) \quad \forall q_1, q_2, q, r \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad q = (q_c/r_c)r \\ \Rightarrow \quad q_1 q_2 = ((q_c/r_c)r)^2 \quad \Rightarrow \quad (r_c/q_c)^2 q_1 q_2 / r^2 = 1.$$

$$(5.10) \quad r = (r_c/t_c)t = ct \quad \Rightarrow \quad mr = (m_c/r_c)(ct)^2 \quad \Rightarrow \quad ((r_c/m_c)/c^2)mr/t^2 = 1.$$

$$(5.11) \quad ((r_c/m_c)/c^2)mr/t^2 = 1 \quad \wedge \quad (r_c/q_c)^2 q_1 q_2 / r^2 = 1 \\ \Rightarrow \quad F := mr/t^2 = ((m_c/r_c)c^2)(r_c/q_c)^2 q_1 q_2 / r^2.$$



$$(5.12) \quad r_c/t_c = c \quad \wedge \quad F = ((m_c/r_c)c^2)(r_c/q_c)^2 q_1 q_2 / r^2 \\ \Rightarrow \quad F = (m_c(r_c/t_c^2))(r_c/q_c)^2 q_1 q_2 / r^2 = k_e q_1 q_2 / r^2,$$

where Coulomb's constant,  $k_e = (m_c(r_c/t_c^2))(r_c/q_c)^2$ , conforms to the SI units:  $N \cdot m^2 \cdot C^{-2}$ .

#### 5.4. 3 fundamental constants. $c_t$ , $c_m$ , and $c_q$ .

$$(5.13) \quad c_t = r_c/t_c \approx 2.99792458 \cdot 10^8 m \ s^{-1}.$$

$$(5.14) \quad G = (r_c/m_c)c_t^2 \quad \Rightarrow \quad c_m = r_c/m_c \approx 7.4261602691 \cdot 10^{-28} m \ kg^{-1}.$$

$$(5.15) \quad k_e = ((m_c/r_c)c_t^2)(r_c/q_c)^2 \Rightarrow \quad c_q = r_c/q_c \approx 8.6175172023 \cdot 10^{-18} m \ C^{-1}.$$

**5.5. Principle of conservation.** A change in distance corresponds to an inversely proportionate change in another type of measure. The ratios of  $c_t/c_m$  and  $c_t/c_q$  yields 3 conservation constants,  $k_t$ ,  $k_m$ , and  $k_q$  that are the basis of particle-wave behavior:

$$(5.16) \quad c_t/c_m = (m_c/r_c)(r_c/t_c) = (m_c r_c)/(t_c r_c) = k_m/k_t.$$

$$(5.17) \quad c_t/c_q = (q_c/r_c)(r_c/t_c) = (q_c r_c)/(t_c r_c) = k_q/k_t.$$

**5.6. Planck-Einstein equation:** Applying both the relative measure ratios 5.1 and the conservation ratios 5.5:

$$(5.18) \quad m(ct)^2 = mr^2 \quad \wedge \quad mr = m_c r_c = k_m \quad \Rightarrow \quad m(ct)^2 = k_m r.$$

$$(5.19) \quad m(ct)^2 = k_m r \quad \wedge \quad r_c/t_c = r/t = c \\ \Rightarrow \quad E := mc^2 = k_m r/t^2 = (k_m(r/t)) (1/t) = (k_m c)(1/t) = hf,$$

where the Planck constant  $h = k_m c$  and the frequency  $f = 1/t$ .

$$(5.20) \quad k_m = m_c r_c = h/c \approx 2.21022 \cdot 10^{-42} \ kg \ m.$$

$$(5.21) \quad k_t = t_c r_c = k_m/(c_t/c_m) \approx 5.47493 \cdot 10^{-78} \ s \ m.$$

$$(5.22) \quad k_q = q_c r_c = (c_t/c_q)k_t \approx 1.90466 \cdot 10^{-52} \ C \ m.$$

**5.7. Quantum-special relativity gravity force.** The total mass of a particle is  $m = \sqrt{m_0^2 + m_{ke}^2}$ , where  $m_0$  is the rest mass and  $m_{ke}$  is the kinetic energy-equivalent mass. Applying both the relative measure ratios 5.1 and the conservation ratios 5.5:

$$(5.23) \quad m_0 = (m_c/r_c)r \quad \wedge \quad m_{ke} = m_c r_c / r \quad \wedge \quad m = \sqrt{m_0^2 + m_{ke}^2} \\ \Rightarrow \quad m = \sqrt{((m_c/r_c)r)^2 + ((m_c r_c)/r)^2}.$$

Applying equation 5.23 to equation 5.6:

$$(5.24) \quad \exists m : m_1 m_2 = m^2 = ((m_c/r_c)r)^2 + ((m_c r_c)/r)^2 \\ \Rightarrow \quad m_1 m_2 / (((m_c/r_c)r)^2 + ((m_c r_c)/r)^2) = 1,$$

where  $r$  is the distance in the coordinate frame of reference. But, the experienced force in the proper (local) frame of reference is, from equation 5.2,  $r' = \sqrt{(ct)^2 - (vt)^2}$ :

$$(5.25) \quad r' = \sqrt{(ct)^2 - (vt)^2} \Rightarrow m_0 r' = (m_c/r_c)((ct)^2 - (vt)^2).$$

$$(5.26) \quad m_0 r' = (m_c/r_c)((ct)^2 - (vt)^2) \Rightarrow ((r_c/m_c)/(c^2 - v^2))m_0 r'/t^2 = 1.$$

$$(5.27) \quad ((r_c/m_c)/(c^2 - v^2))m_0 r'/t^2 = 1 \\ \wedge m_1 m_2 / (((m_c/r_c)r)^2 + ((m_c r_c)/r)^2) = 1 \\ \Rightarrow F := m_0 r'/t^2 = ((m_c/r_c)(c^2 - v^2))m_1 m_2 / (((m_c/r_c)r)^2 + ((m_c r_c)/r)^2).$$

**5.8. Quantum-special relativity charge force.** Applying  $m = (m_c/q_c)q$  to the quantum-relativistic gravity equation (5.7):

$$(5.28) \quad F = (m_c/q_c)^2 (m_c/r_c)(c^2 - v^2)q_1 q_2 / (((m_c/r_c)r)^2 + ((m_c r_c)/r)^2).$$

## 6. Insights and implications

- (1) Proving that volume and distance are derived from the same abstract, set-based definition of a countable n-volume provides a unifying and more rigorous set and limit-based foundation under integration, measure theory, the vector norm, and the metric space axioms without using the geometric primitives and relations required in Euclidean geometry [Joy98], axiomatic geometry [Lee10], and vector analysis [Wey52].
- (2) Every n-volume,  $v$ , has a corresponding  $d^n$ -volume. Therefore, a function that has the properties of a complete metric space, but no corresponding Minkowski distance,  $d : d^n = \sum_{i=1}^m d_i^n$ , is not sufficient to model the intuitive notion of an n-volume as a bijective function of distance (bijective implying distance also being an inverse function of an n-volume).
- (3) Euclid's proof that Euclidean distance is the smallest distance between two distinct points equate Euclidean distance to a straight line, where it is assumed that the straight line length is the smallest distance [Joy98]. And proofs that a straight line is the smallest distance equate the straight line to the Euclidean distance.

Using the calculus of variations for a shortest distance proof would result in circular logic due to the Euclidean assumptions in the definitions of the Riemann, Lebesgue, and line integrals.

All distance measures of an Euclidean 2-volume (area) are Minkowski distances (4.2), where  $1 \leq n \leq 2$ .  $n = 1$  is the Manhattan (largest) distance case,  $d = \sum_{i=1}^m s_i$ . And  $n = 2$  is the Euclidean (smallest) distance case,  $d = (\sum_{i=1}^m s_i^2)^{1/2}$ . Here, there is no need for geometric primitives, like straight line, triangle, etc.

- (4) Compare the distance sum inequality (4.4),

$$(6.1) \quad (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n},$$

to Minkowski's sum inequality:

$$(6.2) \quad (\sum_{i=1}^m (a_i^n + b_i^n)^n)^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}$$

[Min53]. Note the exponent difference in the left side of each equation:

$$(6.3) \quad (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \quad \text{vs.} \quad (\sum_{i=1}^m (a_i^n + b_i^n)^{\mathbf{n}})^{1/n}.$$

The two inequalities only intersect where  $n = 1$ . The distance sum inequality is a more fundamental inequality because its proof does not require the convexity and various inequality theorems required to prove the Minkowski sum inequality. And the distance sum inequality is derived from the definitions of volume and distance, which makes it more directly related to geometry.

- (5) The gravity (5.8), charge (5.12), and Planck (5.19) constants were all derived from more fundamental constants,  $(r_c/t_c) = c_t$ ,  $(r_c/m_c) = c_m$ ,  $(r_c/q_c) = c_q$ , and  $m_c r_c = k_m$ . And all depend on the speed of light constant,  $c_t$ : For example,  $G = c_m c_t^2$ ,  $k_e = (c_q^2/c_m) c_t^2$ , and  $h = k_m c_t$ .
- (6) Algebraic manipulation of Coulomb's constant,  $k_e = (r_c/q_c)^2 ((m_c/r_c) c^2) = (m_c (r_c/t_c^2)) (r_c/q_c)^2$ , yields the constant acceleration term,  $r_c/t_c^2$ , which suggests there might be a maximum acceleration constant.
- (7) The derivations of the spacetime equations, here (5.1), differ from other derivations.
  - (a) The derivations, here, are shorter and simpler.
  - (b) The derivations, here, do not rely on the Lorentz transformations or Einsteins' postulates [Ein15]. The derivations do not even require the notion of light.
  - (c) The derivations, here, rely only on geometry: the Euclidean volume proof (3.2), the Minkowski distances proof (4.1), and the 3D proof (4.12), which provides the insight that the properties of physical space creates a maximum speed, and the spacetime equations.
  - (d) The derivations are valid for spacetime, spacemass, and spacecharge.
- (8) Applying the ratios to derive Newton's gravity force (5.2) and Coulomb's charge force (5.3) equations provide:
  - (a) Derivations that do not assume the inverse square law or Gauss's flux divergence theorem. **Note:** the components of the Ricci and metric tensors in Einstein's field equations have the units,  $1/\text{distance}^2$  [Wey52], which is an assumption of the inverse square law.
  - (b) The first derivations to show that the inverse square law and the property of force as mass times acceleration are the result of the conversion ratios,  $r = (r_c/t_c)t = (r_c/m_c)m$ . And the derivation of the inverse square law does not rely on Gauss's flux divergence on the surface of a sphere.

The quantum-special relativity extension to Newton's gravity equation (5.26) makes empirically verifiable predictions.

- (a) In Newton's gravity force, Gauss's gravity law, and Einstein's field (general relativity) equations, the force,  $F \rightarrow \infty$  as the distance,  $r \rightarrow 0$ . But, in the quantum-special relativity extension to Newton's gravity equation,  $F \rightarrow 0$  as  $r \rightarrow 0$ . Where the distance between point-like particles is less than approximately  $10^{-5} m$ , the gravity force should be measurably smaller than at  $10^{-3} m$ , which implies larger black hole radii and maybe allows black hole evaporation. An approximation in Einstein's field (general relativity) equations is a second metric tensor where the components have the units, " $\text{distance}^2$ ."

- (b) Further, Newton’s gravity constant,  $G$ , Gauss’s constant,  $4\pi G$ , and Einstein’s gravity constant,  $k = 8\pi G/c^4$ , [Wey52], are only valid where the local velocity,  $v = 0$ . At relativistic speeds,  $G$  should be replaced with “ $((m_c/r_c)(c^2 - v^2))$ ”.

In the local (proper) frame of reference, for example, an observer on a rocket passing a star will measure a lower gravity force as the velocity increases. But, in the coordinate frame of reference, for example, an observer on earth looking at the rocket passing the distant star, the effect of the star’s gravity on the rocket would appear to increase as the velocity of the rocket increases.

- (9) There is no unit-factoring ratio converting a discrete state value to a continuously varying interval length. Therefore, the spin states of two quantum entangled particles and the polarization states of two quantum entangled photons are independent of varying distance and time interval lengths.
- (10) Linear algebra, vector analysis, differential geometry, etc. assume any number of possible dimensions. For example, the Gram-Schmidt process is a method to find an orthogonal vector for any  $n$ -dimensional vector [Coh21]. None of those disciplines have exposed the properties that can limit a geometry to 3 dimensions. But the proof that a strict linearly ordered and symmetric set is a cyclic set of at most 3 members (4.12) is the simplest explanation for observing only 3 dimensions of physical space.
- (a) If there are higher dimensions of ordered and symmetric geometric space, then there is a set of at most three members (4.12), each member being an ordered and symmetric set of 3 dimensions (three 3-dimensional balls).
- (b) Each of 3 ordered and symmetric dimensions of space can have at most 3 sequentially ordered and symmetric state values, for example, an ordered and symmetric set of 3 vector orientations,  $\{-1, 0, 1\}$ , per dimension of space and at most 3 spin states per dimension, etc.

## References

- [Coh21] M. Cohen, *Linear algebra: Theory, intuition, code*, Amazon Kindle, 2021. ↑12
- [Coq23] Coq, *Coq proof assistant*, 2023. <https://coq.inria.fr/documentation>. ↑2
- [Ein15] A. Einstein, *Relativity, the special and general theory*, Princeton University Press, 2015. ↑11
- [Gol76] R. R. Goldberg, *Methods of real analysis*, John Wiley and Sons, 1976. ↑1
- [Joy98] D. E. Joyce, *Euclid’s elements*, 1998. <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>. ↑10
- [Lee10] J. M. Lee, *Axiomatic geometry*, American Mathematical Society, 2010. ↑10
- [Min53] H. Minkowski, *Geometrie der zahlen*, Chelsea, 1953. reprint. ↑11
- [Rud76] W. Rudin, *Principles of mathematical analysis*, McGraw Hill Education, 1976. ↑1
- [Wey52] H. Weyl, *Space-time-matter*, Dover Publications Inc, 1952. ↑10, 11, 12