

Copyright © 2025 by George M. Van Treeck. Creative Commons Attribution License.

## **Some Set Properties Underlying Geometry and Physics**

George M. Van Treeck

(Dated: December 29, 2025)

# Abstract

Euclidean volume and some distance equations are instances of abstract sets of ordered combinations (n-tuples). The same sets of n-tuples include another combinatorial property limiting a set to at most 3 elements, e.g., a set of 3 distance dimensions and a set of 3 non-distance dimensions (time, mass, and charge). Where the space near each local coordinate point is Euclidean, each Cartesian axis distance unit corresponds to axis units of time, mass, and charge. Axis unit ratios are used for shorter and simpler derivations of gravity, charge, electromagnetic, relativity, quantum physics equations, and related constants. And the ratios are also used for simple quantum extensions to classical and relativity equations. All the proofs are verified in Rocq.

[1]

## CONTENTS

Introduction	3
Ruler measure and convergence	6
Volume	7
Euclidean volume	7
Distance	8
Vector inner product	8
Minkowski distance ( $p$ -norm)	9
Metric Space	10
Properties limiting a set to at most 3 elements	11
Applications to physics	13
Derivation of space-time-mass-charge	13
Derivation of $G$ , and the Newton, Gauss, and Poisson gravity laws	14
Derivation of Schwarzschild's gravitational time dilation and black hole metric	15
Simple method to find general relativity solutions	16
Derivation of $k_e$ , $\varepsilon_0$ , and the Coulomb and Gauss charge laws	17
Derivation of $B$ , $\mu_0$ , and Lorentz's law	18

Derivation of Faraday's law	19
3 direct proportion ratios: $c_t$ , $c_m$ , and $c_q$	19
3 inverse proportion ratios: $k_t$ , $k_m$ , and $k_q$	19
Derivation of $\hbar$ , $h$ , and the Planck relation	20
Derivation of 4 quantum (Planck) units: $r_p$ , $t_p$ , $m_p$ , $q_p$	20
Derivation of the fine structure constant, $\alpha$	20
Derivation of the Compton wavelength, $\lambda$	21
Derivation of Schrödinger's position-space wave equation	21
Derivation of Dirac's wave equation	22
Total of a type	23
Quantum extension to general relativity	23
Quantum extension to Newton's gravity force	24
Quantum extension to Coulomb's charge force	24
Insights and implications	25
References	30

**Keywords:** mathematical physics, combinatorics, set theory, distance measure, inner product, gravity, charge, electromagnetism, relativity, quantum physics.

## INTRODUCTION

Classical (Galilean coordinate) physics equations assume Euclidean space. And general relativity assumes a pseudo-Riemann geometry (an inner product space), where the space near each local coordinate point is Euclidean [2][3].

But integration, topology, measure theory, and vector analysis define volume, distance, and inner product, where the definitions (axioms) are motivated by geometry [4][5][6][3]. Deriving volume and distance equations without modeling the geometry notions of: point, straight line, side, angle, length, area, and volume, provides an analytic (set, sequence, limit, and combinatorial) perspective, which has applications to geometry and physics.

The proofs, in this article, have been verified using the Rocq proof verification system [7]. The formal proofs are in the Rocq files, "euclidrelations.v" and "threed.v," which are included as ancillary files.

Let  $|x_i|$  be the cardinal of (number of elements in) the countable set,  $x_i \in \{x_1, \dots, x_n\}$ . The Euclidean volume equation will be proved to be an instance of the number of ordered combinations (n-tuples),  $v_c$ , where:

$$\forall v_c, |x_i| \in \{0, \mathbb{N}\}, x_i \in \{x_1, \dots, x_n\}, v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i, s_i, v \in \mathbb{R}. \quad (1)$$

For all  $n > 1$ , there are some cases where multiple domain values,  $|x_1|, \dots, |x_n|$ , multiplied yield the same range value,  $v_c$ . Inferring a domain value,  $d_c$ , from  $v_c$ , requires an inverse (bijective) function,  $d_c = f_n^{-1}(v_c)$  and  $v_c = f_n(d_c)$ . There is only one bijective case for  $n = 1$ :  $v_c = |x_1| = d_c$ . There is only one bijective case for all  $n$  that includes the  $n = 1$  case:

$$\exists d_c, v_c, |x_i| \in \{0, \mathbb{N}\} : v_c = \prod_{i=1}^n |x_i| = \prod_{i=1}^n d_c = d_c^n. \quad (2)$$

A set of n-tuples being the union of subsets of n-tuples implies that the domain value,  $d_c$ , is also the inverse function of the sum of n-tuples, where it will be proved that:

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m (\prod_{j=1}^n |x_{i,j}|) \Rightarrow d^n = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad (3)$$

**Note:**  $s_{i,j} \in \mathbb{R}$  ( $s_{i,j}$   $\pm$ -signed) and the  $n = 2$  case is the vector inner product.

Where each  $v_{c_i}$  is also the bijective function,  $v_{c_i} = d_{c_i}^n$ :

$$d_c^n = \sum_{i=1}^m d_{c_i}^n \Rightarrow d^n = \sum_{i=1}^m d_i^n. \quad (4)$$

$|d|$  is the  $p$ -norm (Minkowski distance) [8], which will be proved to imply the metric space properties [5]. **Note:** the  $n = 2$  case is the Euclidean distance.

The definitions of the sets of n-tuples and the corresponding sets of volume and distance domain values assign a total order, strict linear sequences to the set elements, 1 through  $n$  via successor relations  $[(R_s(X, \rightarrow))]$  [6] and  $n$  through 1 via predecessor relations  $[(R_p(X, \leftarrow))]$ . For example,  $X = x_1 \cup \dots \cup x_n$  and  $d^n = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j})$  or  $d^n = \sum_{i=1}^m (\prod_{j=n}^1 s_{i,j})$ .

But the union, multiplication, and addition operations defining the total number of n-tuples and the corresponding volume and distance equations are commutative, where the commutative property also allows sequencing  $n$  number of set elements in each of the  $n!$  possible unique sequences. For example, for  $n = 3$  there are  $n! = 3! = 6$  possible sequences:  $X = x_1 \cup x_2 \cup x_3$ ,  $X = x_1 \cup x_3 \cup x_2$ ,  $X = x_2 \cup x_1 \cup x_3$ ,  $X = x_2 \cup x_3 \cup x_1$ ,  $X = x_3 \cup x_1 \cup x_2$ , and  $X = x_3 \cup x_2 \cup x_1$ .

Only a cyclic total order allows each element to be the first element of sequences via successor and predecessor relations. For example, for  $n = 3$  and starting at  $x_2$ :  $X = x_2 \cup x_3 \cup x_1$  (cyclic successors) and  $X = x_2 \cup x_1 \cup x_3$  (cyclic predecessors). And the only cyclic totally ordered set that allows sequencing in all  $n!$  possible sequences is where each set element is sequentially adjacent to every other element (either an *immediate* cyclic successor or an *immediate* cyclic predecessor to every other set element), herein, referred to as an immediate symmetric cyclic set (ISCS). An ISCS will be proved to have  $n \leq 3$  elements.

Application to physics uses the following 3 hypotheses:

1. **ISCS:**  $\{r_1, r_2, r_3\}$  is an ISCS of 3 distance dimensions and  $\{t$  (*time*),  $m$  (*mass*),  $q$  (*charge*) $\}$  is the ISCS of 3 “non-distance” dimensions, each dimension  $\subseteq \mathbb{R}$ . Physical space is 6-dimensional:  $r_1$ - $r_2$ - $r_3$ - $t$ - $m$ - $q$ .
2. **Cartesian:** Where the space near each local coordinate point is Euclidean, each Cartesian axis unit length,  $r_p$ , of distance,  $r$ , there are corresponding constant Cartesian axis unit lengths:  $t_p$  of time,  $t$ ;  $m_p$  of mass,  $m$ ; and  $q_p$  of charge,  $q$ , such that:  $r = (r_p/t_p)t = (r_p/m_p)m = (r_p/q_p)q$ .
3. **Maximum ratios** Let:  $r_p/t_p = c_t$ ,  $r_p/m_p = c_m$ , and  $r_p/q_p = c_q$ . The Cartesian axis unit ratios,  $c_t$ ,  $c_m$ , and  $c_q$  are the largest ratios. For example, the speeds of gravity and electromagnetic waves are limited to  $c_t$ .

The proofs and the 3 direct proportion ratios,  $c_t$ ,  $c_m$ , and  $c_q$ , are used to provide simpler and shorter derivations of: the special relativity equations[2][3], the gravitational constant,  $G = c_m c_t^2$ , the Newton, Gauss, and Poisson gravity equations [9][10], the Schwarzschild gravitational time dilation and black hole metric equations [11][12] (illustrating a simple method of finding solutions to Einstein’s general relativity equations), the Coulomb, Lorentz, Faraday electromagnetic laws[10] and their related constants  $k_e = c_q^2 c_t^2 / c_m$ ,  $\varepsilon_0 = 1/4\pi k_e$ ,  $\mu_0 = 4\pi c_q^2 / c_m$ . Derivations from the ratios show that  $G$ ,  $k_e$ ,  $\varepsilon_0$ ,  $\mu_0$ , and  $\hbar$  are **not** fundamental (atomic) constants.

Algebraic manipulation of the 3 direct proportion ratios yields 3 inverse proportion ratios,  $r = t_p r_p / t = m_p r_p / m = q_p r_p / q$ , letting  $k_t = t_p r_p$ ,  $k_m = m_p r_p$ , and  $k_q = q_p r_p$ . The direct and inverse proportion ratios are used to derive the Planck relation [10][13] and the reduced Planck constant,  $\hbar = k_m c_t$ . The values of  $k_t$ ,  $k_m$ , and  $k_q$  are derived from  $\hbar$ ,  $c_t$ ,  $c_m$ , and  $c_q$ . The values of  $r_p$ ,  $t_p$ ,  $m_p$ , and  $q_p$  are calculated from the ratios and are the Planck units.

The ratios and Planck relation are used to derive the Compton wavelength, the position-space Schrödinger, and the Dirac wave equations [10][15][16]. And the inverse proportion ratios are used to add quantum extensions to some general relativity and classical physics equations.



**Lemma .5.** *The number of  $n$ -tuples,  $v_c$ , is the sum of the number of  $n$ -tuples,  $v_{c_i}$ , in each subset of  $n$ -tuples, implies a volume is the sum of volumes,*

$$v_c = \sum_{i=1}^m v_{c_i} \quad \Rightarrow \quad v = \sum_{i=1}^m v_i, \quad v, v_i \in \mathbb{R}.$$

*The formal proof, “sum\_of\_volumes,” is in the Rocq file, euclidrelations.v.*

*Proof.* From the condition of this theorem:

$$v_c = \sum_{i=1}^m v_{c_i} \quad \Leftrightarrow \quad \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_i} \kappa). \quad (18)$$

Apply lemma .3 to equation 18:

$$\begin{aligned} \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} \left( \sum_{j=1}^m v_{c_i} \right) \kappa \quad \wedge \quad \lim_{\kappa \rightarrow 0} \kappa^n = \lim_{\kappa \rightarrow 0} \kappa \\ \Leftrightarrow \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_i} \kappa). \end{aligned} \quad (19)$$

Apply the ruler (.1) and ruler convergence theorem (.2) to equation 19:

$$\begin{aligned} \exists v, v_i : v = \lim_{\kappa \rightarrow 0} v_c \kappa \quad \wedge \quad \lim_{\kappa \rightarrow 0} v_c \kappa = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_i} \kappa) \\ \Rightarrow v = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_i} \kappa). \end{aligned} \quad (20)$$

Apply the ruler (.1) and ruler convergence theorem (.2) to equation 20:

$$v = \lim_{\kappa \rightarrow 0} \sum_{j=1}^m (v_{c_i} \kappa) \quad \wedge \quad \exists v_i, v_{c_i} : v_i = \lim_{\kappa \rightarrow 0} v_{c_i} \kappa \quad \Rightarrow \quad v = \sum_{j=1}^m v_i. \quad (21)$$

□

## DISTANCE

**Definition .6.** Bijective, countable domain value,  $d_c$ :

$$\forall v_c, d_c, |x_i| \in \{0, \mathbb{N}\}, \quad x_i \in \{x_1, \dots, x_n\}, \quad v_c = \prod_{i=1}^n |x_i| = \prod_{i=1}^n d_c = d_c^n. \quad (22)$$

## Vector inner product

**Theorem .7.** *Sum of volumes distance:*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} \quad \Rightarrow \quad d^n = \sum_{i=1}^m \left( \prod_{j=1}^n s_{ij} \right).$$

*The formal proof, “sum\_of\_volumes\_distance,” is in the Rocq file, euclidrelations.v.*



*Proof.* From the sum of volumes lemma .5 and the Euclidean volume theorem .4:

$$\begin{aligned} d_c^n = \sum_{i=1}^m v_{c_i} &\Rightarrow d^n = \sum_{i=1}^m v_i \quad \wedge \quad v_i = \prod_{j=1}^n s_{ij} \\ &\Rightarrow d^n = \sum_{i=1}^m v_i = \sum_{i=1}^m (\prod_{j=1}^n s_{ij}). \quad \square \quad (23) \end{aligned}$$

**Note:** In the volume proof (.4), where  $\kappa$  is negative, volume is negative. From the lemma (.3),  $\forall n \geq 1, 0 < |\kappa| < 1 : \lim_{\kappa \rightarrow 0} \kappa^n = \lim_{\kappa \rightarrow 0} \kappa, \exists \kappa_i \in \{\kappa_1, \dots, \kappa_n\}$ . Therefore,  $\kappa_i$  can have negative values, where  $\prod_{i=1}^n \kappa_i = \kappa^n$ . And where  $\kappa_i < 0$ , the corresponding  $s_{i,j} < 0$ . Therefore, the  $n = 2$  case of the sum of volumes distance is the vector inner product.

### Minkowski distance ( $p$ -norm)

**Theorem .8.** *Minkowski distance ( $p$ -norm):*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m d_{c_i}^n \Leftrightarrow d^n = \sum_{i=1}^m d_i^n.$$

*The formal proof, "Minkowski\_distance," is in the Rocq file, euclidrelations.v.*

*Proof.* From the sum of volumes distance theorem .7 and the Euclidean volume theorem .4:

$$\begin{aligned} d_c^n = v_c = \sum_{i=1}^m v_{c_i} &\Rightarrow d^n = v = \sum_{i=1}^m v_i \quad \wedge \quad v_i = \prod_{j=1}^n d_i = d_i^n \\ &\Rightarrow d^n = \sum_{i=1}^m d_i^n \quad \square \quad (24) \end{aligned}$$

**Theorem .9.** *Distance triangle inequality*

$$\forall n \in \mathbb{N}, v_a, v_b \geq 0 : (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}.$$

*The formal proof, distance\_inequality, is in the Rocq file, euclidrelations.v.*

*Proof.* Expand  $(v_a^{1/n} + v_b^{1/n})^n$  using the binomial expansion:

$$\begin{aligned} \forall v_a, v_b \geq 0 : v_a + v_b &\leq v_a + v_b + \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^{n-i} (v_b^{1/n})^i + \\ &\quad \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^i (v_b^{1/n})^{n-i} = (v_a^{1/n} + v_b^{1/n})^n. \quad (25) \end{aligned}$$

Take the  $n^{th}$  root of both sides of the inequality 25:

$$\forall v_a, v_b \geq 0, n \in \mathbb{N} : v_a + v_b \leq (v_a^{1/n} + v_b^{1/n})^n \Rightarrow (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}. \quad (26)$$

□

**Theorem .10.** *Distance sum inequality*

$$\forall m, n \in \mathbb{N}, a_i, b_i \geq 0 : (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}.$$

The formal proof, *distance\_sum\_inequality*, is in the Rocq file, *euclidrelations.v*.

*Proof.* Apply the distance triangle inequality (.9):

$$\begin{aligned} \forall m, n \in \mathbb{N}, v_a, v_b \geq 0 : v_a = \sum_{i=1}^m a_i^n \quad \wedge \quad v_b = \sum_{i=1}^m b_i^n \quad \wedge \quad (v_a + v_b)^{1/n} &\leq v_a^{1/n} + v_b^{1/n} \\ \Rightarrow ((\sum_{i=1}^m a_i^n) + (\sum_{i=1}^m b_i^n))^{1/n} &= (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq \\ &(\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad \square \quad (27) \end{aligned}$$

**Metric Space**

All Minkowski distances ( $p$ -norms) imply the metric space properties. The formal proofs: *triangle\_inequality*, *symmetry*, *non\_negativity*, and *identity\_of\_indiscernibles* are in the Rocq file, *euclidrelations.v*.

**Theorem .11.** *Triangle Inequality:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, w) \leq d(u, v) + d(v, w).$$

*Proof.*  $\forall p \geq 1, \quad k > 1, \quad u = s_1, \quad w = s_2, \quad v = w/k$ :

$$(u^p + w^p)^{1/p} \leq ((u^p + w^p) + 2v^p)^{1/p} = ((u^p + v^p) + (v^p + w^p))^{1/p}. \quad (28)$$

Apply the distance triangle inequality (.9) to the inequality 28:

$$\begin{aligned} (u^p + w^p)^{1/p} &\leq ((u^p + v^p) + (v^p + w^p))^{1/p} \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\ &\quad \wedge \quad v_a = u^p + v^p \quad \wedge \quad v_b = v^p + w^p \\ \Rightarrow (u^p + w^p)^{1/p} &\leq ((u^p + v^p) + (v^p + w^p))^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} \\ \Rightarrow d(u, w) &= (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \quad \square \quad (29) \end{aligned}$$

**Theorem .12.** *Symmetry:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, v) = d(v, u).$$

*Proof.* By the commutative law of addition:

$$\begin{aligned} \forall p : p \geq 1, \quad d(s_1, s_2) &= (\sum_{i=1}^2 s_i^p)^{1/p} = (s_1^p + s_2^p)^{1/p} \\ \Rightarrow d(u, v) &= (u^p + v^p)^{1/p} = (v^p + u^p)^{1/p} = d(v, u). \quad \square \quad (30) \end{aligned}$$

**Theorem .13.** *Non-negativity:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} \Rightarrow d(u, w) \geq 0.$$

*Proof.* By definition, the length of an interval is always  $\geq 0$ :

$$\forall [a_1, b_1], [a_2, b_2], \quad u = b_1 - a_1, \quad v = b_2 - a_2, \quad \Rightarrow \quad u \geq 0, \quad v \geq 0. \quad (31)$$

$$p \geq 1, \quad u, v \geq 0 \quad \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} \geq 0. \quad (32)$$

□

**Theorem .14.** *Identity of Indiscernibles:*  $d(u, u) = 0$ .

*Proof.* From the non-negativity property (.13):

$$\begin{aligned} d(u, w) \geq 0 \quad \wedge \quad d(u, v) \geq 0 \quad \wedge \quad d(v, w) \geq 0 \\ \Rightarrow \quad \exists d(u, w) = d(u, v) = d(v, w) = 0. \end{aligned} \quad (33)$$

$$d(u, w) = d(v, w) = 0 \quad \Rightarrow \quad u = v. \quad (34)$$

$$d(u, v) = 0 \quad \wedge \quad u = v \quad \Rightarrow \quad d(u, u) = 0. \quad (35)$$

□

### Properties limiting a set to at most 3 elements

The following definitions and proof use first order logic. A Horn clause-like expression is used, here, to make the proof easier to read. By convention, the proof goal is on the left side and supporting facts are on the right side of the implication sign ( $\leftarrow$ ). The formal proofs in the Rocq file `threed.v` are:

Lemmas: `adj111`, `adj122`, `adj212`, `adj123`,  
`adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and  
`not_all_mutually_adjacent_gt_3`.

**Definition .15.** Immediate Cyclic Successor of  $m$  is  $n$ :

$$\begin{aligned} \forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \\ \text{Successor}(m, n, \text{setsize}) \quad \leftarrow \quad (m = \text{setsize} \wedge n = 1) \quad \vee \quad (n = m + 1 \leq \text{setsize}). \end{aligned} \quad (36)$$

**Definition .16.** Immediate Cyclic Predecessor of  $m$  is  $n$ :

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : \\ Predecessor(m, n, setsize) \leftarrow (m = 1 \wedge n = setsize) \vee (n = m - 1 \geq 1). \quad (37)$$

**Definition .17.** Adjacent: element  $m$  is sequentially adjacent to element  $n$  if the immediate cyclic successor of  $m$  is  $n$  or the immediate cyclic predecessor of  $m$  is  $n$ . Notionally:

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : \\ Adjacent(m, n, setsize) \leftarrow Successor(m, n, setsize) \vee Predecessor(m, n, setsize). \quad (38)$$

**Definition .18.** Immediate Symmetric (every set element is sequentially adjacent to every other element):

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : Adjacent(m, n, setsize). \quad (39)$$

**Theorem .19.** An immediate symmetric cyclic set is limited to at most 3 elements.

*Proof.*

Every element is adjacent to every other element, where  $setsize \in \{1, 2, 3\}$ :

$$Adjacent(1, 1, 1) \leftarrow Successor(1, 1, 1) \leftarrow (m = setsize \wedge n = 1). \quad (40)$$

$$Adjacent(1, 2, 2) \leftarrow Successor(1, 2, 2) \leftarrow (n = m + 1 \leq setsize). \quad (41)$$

$$Adjacent(1, 2, 3) \leftarrow Successor(1, 2, 3) \leftarrow (n = m + 1 \leq setsize). \quad (42)$$

$$Adjacent(2, 1, 3) \leftarrow Predecessor(2, 1, 3) \leftarrow (n = m - 1 \geq 1). \quad (43)$$

$$Adjacent(3, 1, 3) \leftarrow Successor(3, 1, 3) \leftarrow (n = setsize \wedge m = 1). \quad (44)$$

$$Adjacent(1, 3, 3) \leftarrow Predecessor(1, 3, 3) \leftarrow (m = 1 \wedge n = setsize). \quad (45)$$

$$Adjacent(2, 3, 3) \leftarrow Successor(2, 3, 3) \leftarrow (n = m + 1 \leq setsize). \quad (46)$$

$$Adjacent(3, 2, 3) \leftarrow Predecessor(3, 2, 3) \leftarrow (n = m - 1 \geq 1). \quad (47)$$

Element 2 is the only immediate successor of element 1 for all  $setsize \geq 3$ , which implies element 3 is not ( $\neg$ ) an immediate successor of element 1 for all  $setsize \geq 3$ :

$$\neg Successor(1, 3, setsize \geq 3) \leftarrow Successor(1, 2, setsize \geq 3) \leftarrow (n = m + 1 \leq setsize). \quad (48)$$

Element  $n = \text{setsize} > 3$  is the only immediate predecessor of element 1, which implies element 3 is not ( $\neg$ ) an immediate predecessor of element 1 for all  $\text{setsize} > 3$ :

$$\neg \text{Predecessor}(1, 3, \text{setsize} \geq 3) \leftarrow \text{Predecessor}(1, \text{setsize}, \text{setsize} > 3) \leftarrow$$

$$(m = 1 \wedge n = \text{setsize} > 3). \quad (49)$$

For all  $\text{setsize} > 3$ , some elements are not ( $\neg$ ) sequentially adjacent to every other element (not immediate symmetric):

$$\neg \text{Adjacent}(1, 3, \text{setsize} > 3) \leftarrow$$

$$\neg \text{Successor}(1, 3, \text{setsize} > 3) \quad \wedge \quad \neg \text{Predecessor}(1, 3, \text{setsize} > 3). \quad \square \quad (50)$$

The Symmetric goal matches Adjacent goals 40 through 47 and fails for all “setsize” greater than three.

## APPLICATIONS TO PHYSICS

Where distance is an immediate symmetric cyclic set (ISCS) of dimensions, the 3D proof (.19) requires more dimensions to have non-distance types (elements of other sets). Let  $\tau = \{t \text{ (time)}, m \text{ (mass)}, q \text{ (charge)}\}$  be the ISCS of type “non-distance” dimensions, where for each Cartesian axis unit length,  $r_p$ , of distance,  $r$ , there are Cartesian axis unit lengths:  $t_p$  of time,  $t$ ;  $m_p$  of mass,  $m$ ; and  $q_p$  of charge,  $q$ , such that:

$$r = (r_p/t_p)t = (r_p/m_p)m = (r_p/q_p)q, \quad (51)$$

where  $c_t$ ,  $c_m$ , and  $c_q$  are the maximum ratios:

$$c = c_t = r_p/t_p, \quad c_m = r_p/m_p, \quad c_q = r_p/q_p. \quad (52)$$

### Derivation of space-time-mass-charge

Let  $r$  be an Euclidean distance. Then by the Minkowski distance theorem (.8),  $r^2 = \sum_{i=1}^m r_i^2$ . Let,  $r' = r_1$  and  $r_v^2 = (\sum_{i=2}^m r_i^2)$ . From the 3D theorem (.19) and Cartesian hypothesis (2):

$$\forall \tau \in \{t, m, q\}, \quad r^2 = r'^2 + r_v^2, \quad \exists \mu, \nu : \quad r = \mu\tau \quad \wedge \quad r_v = \nu\tau \quad \Rightarrow \quad (\mu\tau)^2 = r'^2 + (\nu\tau)^2$$

$$\Rightarrow \quad r' = \sqrt{(\mu\tau)^2 - (\nu\tau)^2} = \mu\tau \sqrt{1 - (\nu/\mu)^2}. \quad (53)$$

Local frame distance,  $r'$ , contracts relative to a distant observer frame distance,  $r$ , as  $\nu \rightarrow \mu$ :

$$r' = \mu\tau\sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad \mu\tau = r \quad \Rightarrow \quad r' = r\sqrt{1 - (\nu/\mu)^2}. \quad (54)$$

A distant observer frame type,  $\tau$ , dilates relative to the local observer frame type,  $\tau'$ , as  $\nu \rightarrow \mu$ :

$$\mu\tau = r'/\sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad r' = \mu\tau \quad \Rightarrow \quad \tau = \tau'/\sqrt{1 - (\nu/\mu)^2}. \quad (55)$$

Where  $\tau$  is type, time, the space-like flat Minkowski spacetime event interval is:

$$\begin{aligned} dr^2 &= dr'^2 + dr_v^2 \quad \wedge \quad dr_v^2 = dr_1^2 + dr_2^2 + dr_3^2 \quad \wedge \quad d(\mu\tau) = dr \\ &\Rightarrow \quad dr'^2 = d(\mu\tau)^2 - dr_1^2 - dr_2^2 - dr_3^2. \end{aligned} \quad (56)$$

### Derivation of $G$ , and the Newton, Gauss, and Poisson gravity laws

From equations 52:

$$r = c_m m \quad \wedge \quad r = c_t t \quad \Rightarrow \quad r/(c_t t)^2 = c_m m/r^2 \quad \Rightarrow \quad r/t^2 = (c_m c_t^2)m/r^2 = Gm/r^2, \quad (57)$$

where  $G = c_m c_t^2$ , has the SI units:  $m^3 \cdot kg^{-1} \cdot s^{-2}$  [9].

Newton's law follows from multiplying both sides of equation 57 by  $m$ :

$$r/t^2 = Gm/r^2 \quad \Leftrightarrow \quad F := mr/t^2 = Gm^2/r^2. \quad (58)$$

$$F = Gm^2/r^2 \quad \wedge \quad \forall m \in \mathbb{R} : \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \quad \Rightarrow \quad F = Gm_1 m_2 / r^2. \quad (59)$$

In this article, a new rationale for Gauss's and Poisson's laws for gravity is presented: Equation 57 relates linear (straight line) acceleration,  $r/t^2$ , to mass and distance. Gauss's gravity field,  $\mathbf{g}$ , and Poisson's gravity field,  $-\nabla\Phi(\tilde{\mathbf{r}}, t)$ , relate angular (orbital) acceleration,  $2\pi r/t^2$ , to mass and distance. Multiplying both sides of equation 57 by  $2\pi$  and differentiating yields Gauss's and Poisson's gravity laws [10]:

$$r/t^2 = Gm/|\tilde{\mathbf{r}}|^2 \quad \wedge \quad \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi r/t^2 \quad \Rightarrow \quad \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi Gm/|\tilde{\mathbf{r}}|^2 \quad (60)$$

$$\mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi Gm/|\tilde{\mathbf{r}}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{g} = \nabla^2\Phi(\tilde{\mathbf{r}}, t) = -4\pi Gm/|\tilde{\mathbf{r}}|^3 = -4\pi G\rho, \quad (61)$$

where  $\rho = m/|\tilde{\mathbf{r}}|^3$  is the mass density:

## Derivation of Schwarzschild's gravitational time dilation and black hole metric

From equations 54 and 51:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - (1)(v^2/c^2)} \wedge c_m m/r = 1 \Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - c_m m v^2 / r c^2}. \quad (62)$$

Where  $v_{\text{escape}}$  is the escape velocity:

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - c_m m v^2 / r c^2} \quad \wedge \quad KE = mv^2/2 = mv_{\text{escape}}^2/2 \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m v_{\text{escape}}^2 / r c^2}. \end{aligned} \quad (63)$$

For a photon, the escape velocity,  $v_{\text{escape}} = c$ .

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2c_m m v_{\text{escape}}^2 / r c^2} \quad \wedge \quad v_{\text{escape}} = c \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2}. \end{aligned} \quad (64)$$

Combining equation 64 with the derivation of  $G$  (59):

$$c_m c^2 = G \quad \wedge \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2} \Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm/r c^2}. \quad (65)$$

Combining equation 65 with equation 55 yields Schwarzschild's gravitational time dilation [11] [12]:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm/r c^2} \quad \wedge \quad t' = t \sqrt{1 - (v^2/c^2)} \Rightarrow t' = t \sqrt{1 - 2Gm/r c^2}. \quad (66)$$

From equations 54 and 65, where Schwarzschild's defined the black hole event horizon radius as  $\alpha := 2Gm/c^2$ :

$$r' = r \sqrt{1 - (v/c)^2} = r \sqrt{1 - 2Gm/r c^2} \quad \wedge \quad \alpha := 2Gm/c^2 \quad \Rightarrow \quad r' = r \sqrt{1 - \alpha/r}. \quad (67)$$

Applying equation 67 to the time-like spacetime interval equation 56:

$$\begin{aligned} r' &= r \sqrt{1 - \alpha/r} \quad \wedge \quad ds^2 = dr'^2 - dt'^2 \\ &\Rightarrow ds^2 = (\sqrt{1 - \alpha/r} dr)^2 - (dr'/\sqrt{1 - \alpha/r})^2 = (1 - \alpha/r) dr^2 - (1 - \alpha/r)^{-1} dt'^2. \end{aligned} \quad (68)$$

$$\begin{aligned} ds^2 &= (1 - \alpha/r) dr^2 - (1 - \alpha/r)^{-1} dt'^2 \quad \wedge \quad dr = d(ct) \quad \wedge \quad c = 1 \quad \wedge \quad \lim_{ds \rightarrow 0} dr' = dr \\ &\Rightarrow \lim_{ds \rightarrow 0} ds^2 = (1 - \alpha/r) dt^2 - (1 - \alpha/r)^{-1} dr^2. \end{aligned} \quad (69)$$

Using spherical coordinates to translate from 2D to 4D yields the  $+- --$  form of Schwarzschild's black hole metric [11] [12]:

$$\begin{aligned} ds^2 &= (1 - \alpha/r)dt^2 - (1 - \alpha/r)^{-1}dr^2 \\ \Rightarrow ds^2 &= (1 - \alpha/r)dt^2 - (1 - \alpha/r)^{-1}dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2) \\ \Rightarrow g_{\mu,\nu} &= \text{diag}[(1 - \alpha/r), -(1 - \alpha/r)^{-1}, -r^2(d\theta^2), -r^2(\sin^2\theta d\phi^2)]. \end{aligned} \quad (70)$$

### Simple method to find general relativity solutions

Einstein's field equation is:

$$G_{\mu,\nu} = \mathbf{R} + \frac{1}{2}Rg_{\mu,\nu} = \kappa T_{\mu,\nu}, \quad (71)$$

where:  $G_{\mu,\nu}$  is Einstein's tensor,  $\mathbf{R}$  is the Ricci curvatature,  $R$  is the scalar curvature,  $g_{\mu,\nu}$  is the metric tensor,  $\kappa = 8\pi G/c^4$ , and  $T_{\mu,\nu}$  is the stress-energy tensor [2][3].

The goal of the field equation is to determine the geodesic path and acceleration of a particle caused by the distribution of mass and energy as specified in the stress-energy tensor,  $T_{\mu,\nu}$ . This requires solving the equation for the metric tensor,  $g_{\mu,\nu}$ . But the metric tensor has a complex nonlinear relation to the Ricci and scalar curvature, which makes it complicated to determine the metric. Often there are no exact solutions.

In this article, the metric,  $g_{\mu,\nu}$ , is determined independent of Einstein's field equation. In this case the metric tensor is the independent variable and the distribution of mass and energy in the stress-energy tensor,  $T_{\mu,\nu}$ , are the dependent variables.

The infinitesimal space near every coordinate point on a pseudo-Riemann surface is Euclidean. In this case, the Cartesian unit ratios (91) (94) apply, where: all physics equations derived from the ratios and special relativity equations (53) are valid around the infinitesimal space at each coordinate point.

This leads to the following steps to solve for the metric tensor,  $g_{\mu,\nu}$ , independent of Einstein's field equation.

Step 1) Use the ratios and relativity equations to define functions returning scalar values for each component of the metric,  $g_{\nu,\mu}$ , in Einstein's field equations [2][3]: All functions derived from the ratios and relativity are valid metrics. An example is the previous Schwarzschild black hole metric derivation using the ratios and special relativity equations (62).



The diagonal components the metric,  $g_{\mu,\nu}$ , are determined as a generalization of the flat Minkowski spacetime interval equation:  $ds^2 = g_{1,1}dx_1^2 + \dots + g_{4,4}dx_4^2$ .  $ds$  is an element of the geodesic line (the affine connection between geodesics).  $x_1, \dots, x_4$  are the dimension of time and the 3 dimensions of space.

Step 2) Express the Einstein field equation as 2D tensors: As shown in equation 70, the Schwarzschild metric was first derived as a 2D metric and then expanded to a 4D metric. Further, the 4D flat spacetime interval equation (56) is an instance of the 2D equation,  $dr'^2 = d(ct)^2 - dr_v^2$ .

(Optional) The 2D metric tensor allows using the much simpler 2D Ricci curvature and scalar curvature. And the 2D tensors reduce the number of independent equations to solve, which can next be used to set constraints on the solutions in the 4D tensors.

Step 3) One simple method to translate from 2D to 4D is to use spherical coordinates, where  $r$  and  $t$  remain unchanged and two added dimensions are the angles,  $\phi$ , and  $\theta$ . For example, the 2D Schwarzschild metric was translated to 4D using this method in equation 70. The spherical coordinates can then be translated to other types of coordinates.

### Derivation of $k_e$ , $\varepsilon_0$ , and the Coulomb and Gauss charge laws

From equation 52:

$$r = c_q q \quad \Rightarrow \quad r^2 = c_q^2 q^2 \quad \Rightarrow \quad c_q^2 q^2 / r^2 = 1. \quad (72)$$

$$\begin{aligned} r = c_m m \quad \wedge \quad r = c_t t \quad \Rightarrow \quad mr &= ((1/c_m)r)(c_t t) = ((1/c_m)(c_t t))(c_t t) = (c_t^2/c_m)t^2 \\ &\Rightarrow \quad (c_m/c_t^2)mr/t^2 = 1. \end{aligned} \quad (73)$$

$$c_q^2 q^2 / r^2 = 1 \quad \wedge \quad (c_m/c_t^2)mr/t^2 = 1 \quad \Rightarrow \quad F := mr/t^2 = (c_q^2 c_t^2 / c_m) q^2 / r^2 = k_e q^2 / r^2, \quad (74)$$

where  $k_e = c_q^2 c_t^2 / c_m$ , conforms to the SI units:  $kg \cdot m^3 \cdot s^{-2} \cdot C^{-2} = N \cdot m^2 \cdot C^{-2}$  [10].

$$\forall q \in \mathbb{R} \exists q_1, q_2 \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad F = k_e q^2 / r^2 \quad \Rightarrow \quad F = k_e q_1 q_2 / r^2. \quad (75)$$

In this article, a new rationale for Gauss's electric field is presented: Coulomb's charge force equation (74) relates linear acceleration,  $r/t^2$ , to charge and distance. Gauss's electric field,  $\mathbf{E}$ , relates angular (orbital or rotational) acceleration,  $2\pi r/t^2$  to charge and distance:

From equation 75:

$$F_C = mr/t^2 = k_e q^2/r^2 \Rightarrow \exists F_E \in \mathbb{R} : F_E = 2\pi(mr/t^2) = 2\pi(k_e q^2/r^2). \quad (76)$$

$$F_E = 2\pi(k_e q^2/r^2) = q(2\pi(k_e/r^2)) \Rightarrow \exists E : \quad \wedge \quad E = 2\pi k_e q/r^2 \quad \wedge \quad F_E = qE. \quad (77)$$

The electric field,  $E := 2\pi k_e q/r'^2$ , conforms to the SI units  $kg \cdot m \cdot s^{-2} \cdot C^{-1} = N \cdot C^{-1}$ .

$$\mathbf{E} = 2\pi k_e q/|\mathbf{r}|^2 \Rightarrow \nabla \cdot \mathbf{E} = -4\pi k_e q/|\mathbf{r}|^3. \quad (78)$$

$$\nabla \cdot \mathbf{E} = -4\pi k_e q/|\mathbf{r}|^3 \quad \wedge \quad \varepsilon_0 := 1/4\pi k_e \quad \wedge \quad \rho = q/|\mathbf{r}|^3 \Rightarrow \nabla \cdot \mathbf{E} = -\rho/\varepsilon_0, \quad (79)$$

which is Gauss's electric field law [10].

### Derivation of $\mathbf{B}$ , $\mu_0$ , and Lorentz's law

Applying the distance contraction equation, 54, to equation 77, where  $r$  is the distant observer frame of reference and  $r'$  is moving particle local frame of reference:

$$r = r'/\sqrt{1 - v^2/c^2} \quad \wedge \quad F = 2\pi k_e q^2/r^2 \Rightarrow F = 2\pi k_e q^2(1 - v^2/c^2)/r'^2. \quad (80)$$

From equation 77:

$$E = 2\pi k_e q/r'^2 \Rightarrow F = q(E - (((2\pi k_e/c^2)q/r'^2))v^2). \quad (81)$$

$$F = q(E - (((2\pi k_e/c^2)q/r'^2))v^2) \Rightarrow \exists B : B = (2\pi k_e/c^2)vq/r'^2 \quad \wedge \quad F = q(E - Bv). \quad (82)$$

$$F = q(E - Bv) \Rightarrow \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{v}). \quad (83)$$

$$\mathbf{B} \times \vec{v} = -(\vec{v} \times \mathbf{B}) \quad \wedge \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{v}) \Rightarrow \mathbf{F} = q(\mathbf{E} + \vec{v} \times \mathbf{B}), \quad (84)$$

which is Lorentz law in the rest (observer on the moving particle) frame of reference, where the magnetic field,  $B = (2\pi k_e/c^2)vq/r'^2$ , conforms to the base SI units:  $kg \cdot s^{-1} \cdot C^{-1} = kg \cdot s^{-2} \cdot A^{-1} = T$ .

$$B = (2\pi k_e/c^2)vq/r'^2 \quad \wedge \quad B := \mu_0 H \quad \wedge \quad \mu_0 := 4\pi k_e/c^2 \Rightarrow H = vq/2r'^2, \quad (85)$$

where  $\mu_0 = 4\pi k_e/c^2 = 4\pi c_q^2/c_m$  conforms to the SI units  $kg \cdot m \cdot C^{-2} = kg \cdot m \cdot s^{-2} A^{-2}$  and  $H = vq/2r'^2$  conforms to the SI units  $C \cdot s^{-1} \cdot m^{-1} = A \cdot m^{-1}$ .

### Derivation of Faraday's law

From the magnetic field equation 83, where the electric and magnetic fields are propagating at the speed,  $v = c$ :

$$B = (2\pi k_e/c^2)qv/r^2 \quad \wedge \quad v = c \quad \wedge \quad r = ct \quad \Rightarrow \quad B = (2\pi k_e/c^3)q/t^2. \quad (86)$$

$$B = (2\pi k_e/c^3)q/t^2 \quad \Rightarrow \quad \partial B/\partial t = -(4\pi k_e/c^3)q/t^3. \quad (87)$$

$$\partial B/\partial t = -(4\pi k_e/c^3)q/t^3 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial B/\partial t = -4\pi k_e q/r^3. \quad (88)$$

From equation 78:

$$\mathbf{E} = 2\pi k_e q/|\vec{r}|^2 \quad \Rightarrow \quad \nabla \times \mathbf{E} = 4\pi k_e q/|\vec{r}|^3. \quad (89)$$

Combining equations 89 and 88 yields Faraday's law [10]:

$$\nabla \times \mathbf{E} = 4\pi k_e q/|\vec{r}|^3 \quad \wedge \quad \partial \mathbf{B}/\partial t = -4\pi k_e q/|\vec{r}|^3 \quad \Rightarrow \quad \nabla \times \mathbf{E} = -\partial \mathbf{B}/\partial t. \quad (90)$$

### 3 direct proportion ratios: $c_t$ , $c_m$ , and $c_q$

$$c_t = c \approx 2.99792458 \cdot 10^8 m \, s^{-1}. \quad (91)$$

$$G = c_m^2 c_t^2 \quad \wedge \quad G \approx 6.67418478 \cdot 10^{-11} m^3/kg/s^2 \quad \Rightarrow \quad c_m = G/c_t^2 \approx 7.42603211 \cdot 10^{-28} m \, kg^{-1}. \quad (92)$$

$$k_e = c_q^2 c_t^2 / c_m \quad \wedge \quad k_e \approx 8.9875517923 \cdot 10^9 Nm^2/C^2 \quad \Rightarrow \quad c_q = \sqrt{k_e c_m / c_t^2} \approx 8.61744282 \cdot 10^{-18} m \, C^{-1}. \quad (93)$$

### 3 inverse proportion ratios: $k_t$ , $k_m$ , and $k_q$

$$\begin{aligned} r/t = r_p/t_p \quad \wedge \quad r/m = r_p/m_p \quad &\Rightarrow \quad (r/t)/(r/m) = (r_p/t_p)/(r_p/m_p) \\ &\Rightarrow \quad (mr)/(tr) = (m_p r_p)/(t_p r_p) \quad \Rightarrow \quad mr = m_p r_p = k_m, \quad tr = t_p r_p = k_t. \end{aligned} \quad (94)$$

$$\begin{aligned} r/t = r_p/t_p \quad \wedge \quad r/q = r_p/q_p \quad &\Rightarrow \quad (r/t)/(r/q) = (r_p/t_p)/(r_p/q_p) \\ &\Rightarrow \quad (qr)/(tr) = (q_p r_p)/(t_p r_p) \quad \Rightarrow \quad qr = q_p r_p = k_q, \quad tr = t_p r_p = k_t. \end{aligned} \quad (95)$$

### Derivation of $\hbar$ , $h$ , and the Planck relation

[10][13] Applying both the direct proportion ratio (91), and inverse proportion ratio (94):

$$r = ct \quad \wedge \quad m = k_m/r \quad \Rightarrow \quad m(ct)^2 = (k_m/r)r^2 = k_m r. \quad (96)$$

$$\begin{aligned} m(ct)^2 = k_m r \quad \wedge \quad r/t = c \\ \Rightarrow E := mc^2 = k_m r/t^2 = (k_m c)(1/t) = \hbar\omega = \hbar\omega(2\pi/2\pi) = hf, \end{aligned} \quad (97)$$

where the reduced Planck constant,  $\hbar = k_m c$ , angular frequency,  $\omega = 1/t$ , the full Planck constant,  $h = 2\pi\hbar$ , and the cycles per second frequency (Hertz),  $f = \omega/2\pi$ .

Using  $\hbar \approx 1.054571817 \cdot 10^{-34}$ :

$$k_m = \hbar/c_t \approx 3.51767294 \cdot 10^{-43} \text{ kg } m. \quad (98)$$

$$k_t = k_m c_m/c_t \approx 8.71347873 \cdot 10^{-79} \text{ s } m. \quad (99)$$

$$k_q = k_t c_t/c_q \approx 3.03133454 \cdot 10^{-53} \text{ C } m. \quad (100)$$

### Derivation of 4 quantum (Planck) units: $r_p$ , $t_p$ , $m_p$ , $q_p$

$$: \quad r_p = \sqrt{r_p^2} = \sqrt{c_t k_t} = \sqrt{c_m k_m} = \sqrt{c_q k_q} \approx 1.61624107 \cdot 10^{-35} \text{ m}. \quad (101)$$

$$t_p = r_p/c_t \approx 5.39119991 \cdot 10^{-44} \text{ s}. \quad (102)$$

$$m_p = r_p/c_m \approx 2.17645313 \cdot 10^{-8} \text{ kg}. \quad (103)$$

$$q_p = r_p/c_q \approx 1.87554604 \cdot 10^{-18} \text{ C}. \quad (104)$$

### Derivation of the fine structure constant, $\alpha$

The ratios of two subtypes of force implies ratios of the form:

$$\alpha_\tau = \frac{F_{\tau_1}}{F_{\tau_2}} = \frac{K\tau_1^2/r^2}{K\tau_2^2/r^2} = \frac{\tau_1^2}{\tau_2^2}. \quad (105)$$

For example, where  $q_e$  is the elementary (electron) charge ( $1.60217663 \cdot 10^{-19} \text{ C}$ ), and  $q_p$  is Planck charge unit, the fine structure electron coupling constant is:

$$\alpha_q = q_e^2/q_p^2 \approx 0.0072973526. \quad (106)$$

### Derivation of the Compton wavelength, $\lambda$

[10][13] From equations 94 and 97:

$$r = k_m/m \quad \wedge \quad h = 2\pi k_m c \quad \Rightarrow \quad \lambda = 2\pi r = 2\pi k_m/m = (2\pi k_m/m)(c/c) = h/mc. \quad (107)$$

### Derivation of Schrödinger's position-space wave equation

Start with the previously derived Planck relation 97 and multiply the kinetic energy component by  $mc/mc$ :

$$\begin{aligned} mc^2 = \hbar\omega = \hbar/t \quad \Rightarrow \quad \exists V(r, t) : \hbar/t &= \hbar/2t + V(r, t) \\ &\Rightarrow \hbar/t = \hbar mc/2mct + V(r, t). \end{aligned} \quad (108)$$

And from the distance-to-time (speed of light) ratio (91):

$$\hbar/t = \hbar mc/2mct + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar mc^2/2mcr + V(r, t). \quad (109)$$

$$\hbar/t = \hbar mc^2/2mcr + V(r, t) \quad \wedge \quad \hbar/t = mc^2 \quad \Rightarrow \quad \hbar/t = \hbar^2/2mcr + V(r, t). \quad (110)$$

$$\hbar/t = \hbar^2/2mcr + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar^2/2mr^2 + V(r, t). \quad (111)$$

Multiply both sides of equation 111 by a function,  $\Psi(r, t)$ .

$$\hbar/t = \hbar^2/2mr^2 + V(r, t) \quad \Rightarrow \quad (\hbar/t)\Psi(r, t) = (\hbar^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t). \quad (112)$$

$$(\hbar/t)\Psi(r, t) = (\hbar^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t) \quad \wedge$$

$$\begin{aligned} \forall \Psi(r, t) : \partial^2 \Psi(r, t)/\partial r^2 &= (-1/r^2)\Psi(r, t) \quad \wedge \quad \partial \Psi(r, t)/\partial t = (i/t)\Psi(r, t) \\ &\Rightarrow \quad i\hbar \partial \Psi(r, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t), \end{aligned} \quad (113)$$

which is the one-dimensional position-space Schrödinger's equation [15][13].

$$\begin{aligned} i\hbar \partial \Psi(r, t)/\partial t &= -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t) \quad \wedge \quad ||\vec{r}'|| = r \\ &\Rightarrow \quad \exists \vec{r} : i\hbar \partial \Psi(\vec{r}, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(\vec{r}, t)/\partial \vec{r}^2 + V(\vec{r}, t)\Psi(\vec{r}, t), \end{aligned} \quad (114)$$

which is the 3-dimensional position-space Schrödinger's equation [15] [13].

## Derivation of Dirac's wave equation

Using the derived Planck relation 97:

$$mc^2 = \hbar/t \quad \Rightarrow \quad \exists V(r, t) : mc^2/2 + V(r, t) = \hbar/t \quad \Rightarrow \quad 2\hbar/t - 2V(r, t) = mc^2. \quad (115)$$

$$\begin{aligned} \forall V(r, t) : V(r, t) = i\hbar/t \quad \wedge \quad r = ct \quad \wedge \quad 2\hbar/t - 2V(r, t) = mc^2 \\ \Rightarrow \quad 2\hbar/t - i2\hbar c/r = mc^2. \end{aligned} \quad (116)$$

Use the ratios,  $r = c_q q$ , and,  $r = ct$ . to multiply each term on the left side of equation 116 by 1:

$$qc_q/r = qc_q/ct = 1 \wedge 2\hbar/t - i2\hbar c/r = mc^2 \Rightarrow 2\hbar(qc_q/c)/t^2 - i2\hbar((qc_q/c)/r^2)c = mc^2. \quad (117)$$

Applying a quantum amplitude equation in complex form to equation 118:

$$\begin{aligned} A_0 = (c_q/c)((1/t) - i(1/r)) \quad \wedge \quad 2\hbar(qc_q/c)/t^2 - i2\hbar((qc_q/c)/r^2)c = mc^2 \\ \Rightarrow \quad 2\hbar\partial(-qA_0)/\partial t - i2\hbar(\partial(-qA_0)/\partial r)c = mc^2. \end{aligned} \quad (118)$$

Translating equation 118 to moving (rest frame) coordinates via the Lorentz factor,  $\gamma_0 = 1/\sqrt{1 - (v/c)^2}$ :

$$\begin{aligned} 2\hbar\partial(-qA_0)/\partial t - i\hbar h(\partial(-qA_0)/\partial r)c = mc^2 \\ \Rightarrow \quad \gamma_0 2\hbar\partial(-qA_0)/\partial t - \gamma_0 i2\hbar(\partial(-qA_0)/\partial r)c = mc^2. \end{aligned} \quad (119)$$

Multiplying both sides of equation 119 by  $\Psi(r, t)$ :

$$\begin{aligned} \gamma_0 2\hbar\partial(-qA_0)/\partial t - \gamma_0 i2\hbar(\partial(-qA_0)/\partial r)c = mc^2 \quad \Rightarrow \\ \gamma_0 2\hbar(\partial(-qA_0)/\partial t)\Psi(r, t) - \gamma_0 i2\hbar(\partial(-qA_0)/\partial r)c\Psi(r, t) = mc^2\Psi(r, t). \end{aligned} \quad (120)$$

Applying the vectors to equation 120:

$$\begin{aligned} \gamma_0 2\hbar(\partial(-qA_0)/\partial t)\Psi(r, t) - \gamma_0 i2\hbar(\partial(-qA_0)/\partial r)c\Psi(r, t) = mc^2\Psi(r, t) \quad \wedge \\ ||\vec{r}|| = r \quad \wedge \quad ||\vec{A}|| = A_0 \quad \wedge \quad ||\vec{\gamma}|| = \gamma_0 \\ \Leftrightarrow \quad \exists \vec{r}, \vec{A}, \vec{\gamma} : \gamma_0 2\hbar(\partial(-qA_0)/\partial t)\Psi(r, t) - \vec{\gamma} \cdot i2\hbar(\partial(-q\vec{A})/\partial r)c\Psi(\vec{r}, t) = mc^2\Psi(\vec{r}, t). \end{aligned} \quad (121)$$

Adding a  $\frac{1}{2}$  spin to equation 118 yields Dirac's wave equation [16] [13]:

$$\begin{aligned} \gamma_0 2\hbar(\partial(-qA_0)/\partial t)\Psi(r, t) - \vec{\gamma} \cdot i2\hbar(\partial(-q\vec{\mathbf{A}})/\partial r)c\Psi(\vec{\mathbf{r}}, t) &= mc^2\Psi(\vec{\mathbf{r}}, t) \\ \wedge \quad A_0 &= \frac{1}{2}(c_q/c)((1/t) - i(1/r)) \\ \Rightarrow \quad \gamma_0 \hbar(\partial(-qA_0)/\partial t)\Psi(r, t) - \vec{\gamma} \cdot i\hbar(\partial(-q\vec{\mathbf{A}})/\partial r)c\Psi(\vec{\mathbf{r}}, t) &= mc^2\Psi(\vec{\mathbf{r}}, t). \end{aligned} \quad (122)$$

### Total of a type

Applying both the direct (91) and inverse proportion ratios (95):

$$r = \sqrt{(c_\tau\tau)^2 + (k_\tau/\tau)^2}, \tau \in \{t, m, q\} \quad \Leftrightarrow \quad \tau = \sqrt{(r/c_\tau)^2 + (k_\tau/r)^2}. \quad (123)$$

### Quantum extension to general relativity

The simplest way to demonstrate how to add quantum physics to general relativity is by extending Schwarzschild's gravitational time dilation equation and black hole metric. Apply the total of a type equation 123 to the derivation of Schwarzschild's time dilation and metric (62):

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - (v^2/c^2)(r/r)} \quad \wedge \quad r = \sqrt{(c_m m)^2 + (k_m/m)^2} = Q_m \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2/rc^2}. \end{aligned} \quad (124)$$

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2/rc^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2 \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2/rc^2}. \end{aligned} \quad (125)$$

For a photon, the escape velocity,  $v_{escape} = c$ .

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2/rc^2}. \quad \wedge \quad v_{escape} = c \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m c^2/rc^2} = \sqrt{1 - 2Q_m/r}. \end{aligned} \quad (126)$$

Combining equation 126 with equation 55 yields Schwarzschild's gravitational time dilation with a quantum mass effect:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m/r} \quad \wedge \quad t' = t\sqrt{1 - (v^2/c^2)} \quad \Rightarrow \quad t' = t\sqrt{1 - 2Q_m/r}. \quad (127)$$

Schwarzschild defined the black hole event horizon radius,  $\alpha := 2Gm/c^2$ . The radius with the quantum extension is  $\alpha := 2Q_m$ . At this point the exact same equations 67 through 70 yield what looks like the same Schwarzschild black hole metric.

### Quantum extension to Newton's gravity force

The quantum mass effect is easier to understand in the context Newton's gravity equation than in general relativity, because the metric equations and solutions in the EFEs are much more complex. From equations 123 and 51:

$$\begin{aligned} m/\sqrt{(r/c_m)^2 + (k_m/r)^2} = 1 \quad \wedge \quad r^2/(ct)^2 = 1 &\Rightarrow r^2/(ct)^2 = m/\sqrt{(r/c_m)^2 + (k_m/r)^2} \\ &\Rightarrow r^2/t^2 = c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2}. \end{aligned} \quad (128)$$

$$\begin{aligned} r^2/t^2 = c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2} &\Rightarrow (m/r)(r^2/t^2) = (m/r)(c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2}) \\ &\Rightarrow F := mr/t^2 = c^2 m^2/\sqrt{(r^4/c_m^2) + k_m^2}. \end{aligned} \quad (129)$$

$$\begin{aligned} F = c^2 m^2/\sqrt{(r^4/c_m^2) + k_m^2} \quad \wedge \quad \forall m \in \mathbb{R}, \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \\ &\Rightarrow F = c^2 m_1 m_2/\sqrt{(r^4/c_m^2) + k_m^2}. \end{aligned} \quad (130)$$

### Quantum extension to Coulomb's charge force

$$\begin{aligned} q^2/((r/c_q)^2 + (k_q/r)^2) = 1 \quad \wedge \quad r^2/(ct)^2 = 1 &\Rightarrow r^2/(ct)^2 = q^2/((r/c_q)^2 + (k_q/r)^2) \\ &\Rightarrow r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2). \end{aligned} \quad (131)$$

$$(1/r)(r^2/t^2) = (1/r)(c^2 q^2/((r/c_q)^2 + (k_q/r)^2)) \Rightarrow r/t^2 = c^2 q^2/(r^3/c_q^2 + k_q^2/r). \quad (132)$$

$$\begin{aligned} \forall q \in \mathbb{R} : \exists q_1, q_2 \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad r/t^2 = c^2 q^2/(r^3/c_q^2 + k_q^2/r) &\Rightarrow \\ &\exists q_1, q_2 \in \mathbb{R} : r/t^2 = c^2 q_1 q_2/(r^3/c_q^2 + k_q^2/r). \end{aligned} \quad (133)$$

$$\begin{aligned} r/t^2 = c^2 q_1 q_2/(r^3/c_q^2 + k_q^2/r) \quad \wedge \quad m = r/c_m &\Rightarrow F := mr/t^2 = (c^2/c_m) q_1 q_2/(r^2/c_q^2 + k_q^2/r^2). \\ &\quad (134) \end{aligned}$$



## INSIGHTS AND IMPLICATIONS

1. The ruler measure (.1) and convergence theorem (.2) were shown to be useful tools for proving that a countable sets of n-tuples imply a corresponding real-valued equation.
2. Defining all Euclidean and non-Euclidean distance measures as the inverse function of the sum of subset volumes:

$$\forall n, d: \quad d = f_n^{-1}(v) = f_n^{-1}(\sum_{i=1}^m v_i) : \quad (135)$$

- (a) shows the intimate relation between distance and volume that definitions, like inner product space and metric space, ignore [3][4][5];
- (b) is a more simple and concise definition of a distance measure that includes the properties of inner product space and metric space [3][4][5].
3. The left side of the distance sum inequality (.10),

$$(\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}, \quad (136)$$

differs from the left side of Minkowski's sum inequality [8]:

$$(\sum_{i=1}^m (a_i^n + b_i^n)^{\mathbf{n}})^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad (137)$$

- (a) The two inequalities are only the same where  $n = 1$ .
- (b) The distance sum inequality (.10) is a more fundamental inequality because the proof does not require the convexity and Hölder's inequality assumptions required to prove the Minkowski sum inequality [8].
- (c)  $\forall n > 1, v > 0$ : The distance sum inequality term,  $v_i^n = a_i^n + b_i^n$ :  $d = v^{1/n} = (\sum_{i=1}^m v_i^n)^{1/n}$ , is the Minkowski distance, which makes it directly related to geometry. But the Minkowski sum inequality term,  $d = v^{1/n} = (\sum_{i=1}^m ((v_i^n)^{\mathbf{n}}))^{1/n} = (\sum_{i=1}^m v_i^{\mathbf{n}^2})^{1/n}$ , is *not* a Minkowski distance.
- (d) The distance sum inequality might be applicable to machine learning.
4. **Combinatorics.** The number of n-tuples,  $v_c = \prod_{i=1}^n |x_i|$ , was proven to imply: the Euclidean volume equation (.4), the inverse function of the sum of volumes equation (.7) (which includes the inner product) and the Minkowski distance equation (.8)

(which includes the Manhattan and Euclidean distance equations), without relying on the geometric primitives and relations in Euclidean geometry [17][18], axiomatic geometry [19][20][21] [22][23], trigonometry [24] [25] calculus [26][24] [27], and vector analysis [3].

5. **Combinatorics.** The commutative property of the operations defining volume and distance equations limits a set to  $n \leq 3$  elements (.19). Other dimensions must have different types (elements of different sets).
  - (a) For example, the vector inner product space can only be extended beyond 3 dimensions if and only if the other dimensions have non-distance types, for example, dimensions of time, mass, and charge.
  - (b) As shown in the special relativity section (53), there is 6-dimensional space-time-mass-charge.
  - (c) If each type of quantum state is an ISCS, then there are at most 3 states of the same type: 3 orientations per dimension of space, 3 quark color charges, {red, green, blue}, 3 quark anti-color charges, and so on.
  - (d) If the states are not ordered (a bag of states), then a state value is undetermined (or superimposed) until observed (like Schrödinger's poisoned cat being both alive and dead until the box is opened [15]).
  - (e) A discrete (point) value has measure 0 (zero-length interval). The ratio of a time or distance interval length to zero is undefined, which is the reason quantum entangled (discrete) state values exist independent of time and distance.
6. For each Cartesian axis unit,  $r_p$ , of a 3-dimensional distance interval having a length,  $r$ , there are Cartesian axis units of other types of intervals forming unit ratios (91):  $c_t = r_p/t_p$ ,  $c_m = r_p/m_p$ ,  $c_q = r_p/q_p \Leftrightarrow$  the inverse proportion ratios (94):  $k_t = r_p t_p$ ,  $k_m = r_p m_p$ ,  $k_t = r_p q_p$ , where  $r_p$ ,  $t_p$ ,  $m_p$ , and  $q_p$  are the Planck units (101).
7. Empirical laws *describe* relations. Deriving empirical laws from the ratios *explains* the relations. Further, all the derivations of the physics equations from the ratios were much shorter and simpler than other derivations, which shows that the ratios are an important tool for physicists and engineers.

8. As shown in the subsection deriving the Schwarzschild's gravitational time dilation and black hole metric (62) [11][12] using ratios illustrates a way of simplifying the finding of solutions to Einstein's field equations.

9. The speed of light is limited to the ratio,  $c_t$ . And  $c_t$  is a component of the ratio-derived constants:  $G = c_m c_t^2$ ,  $k_e = c_q^2 c_t^2 / c_m$ ,  $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2 / c_m)$ , and  $\hbar = k_m c_t$ .

The only ratio-derived constant, in this article, that does not contain  $c_t$  is vacuum permeability:  $\mu_0 = 4\pi k_e / c_t^2 = 4\pi c_q^2 / c_m$ .

10. In the derivation of the Planck relation (97), one could start with  $k_m c = h$  (the full Planck constant) instead of  $k_m c = \hbar$  (the reduced Planck constant). But,  $k_m c = h$  would require: 1) the derived quantum units (101) to not be the Planck units (that is, a quantum unit would = Planck unit  $\times \sqrt{2\pi}$ ), 2) make the derivation of the fine structure constant (106) more complicated, 3) make the derivation of Schrödinger (114) and Dirac (122) wave equations more complicated and finding solutions to those wave equations more complicated.

11. Currently, the full Planck constant,  $h$  is assigned a standard value and the reduced Planck constant,  $\hbar$  is defined as  $\hbar = h/2\pi$  [14]. But, the derivations, in this article, suggest that the reduced Planck constant should have a standard value and the full Planck constant,  $h$ , defined as  $h = 2\pi\hbar$ .

12. Using the quantum (Planck) units,  $r_p$  and  $t_p$ :  $r_p/t_p^2$ , suggests a maximum linear acceleration for masses. And  $2\pi r_p/t_p^2$  suggests a maximum orbital or rotational acceleration.

13. The simplification of  $\mu_0$  into the quantum units shows two interesting relationships:

$$\begin{aligned} \mu_0 &= 4\pi \frac{k_e}{c_t^2} = 4\pi \frac{c_q^2}{c_m} = 4\pi \frac{(r_p/q_p)^2}{r_p/m_p} = 4\pi \frac{m_p r_p}{q_p^2} = 4\pi \frac{k_m}{q_p^2} \\ &\approx 4\pi \frac{3.5176729162 \cdot 10^{-43}}{3.5176729162 \cdot 10^{-35}} = 4\pi \cdot 10^{-7} \text{ kg m C}^{-2} = 4\pi \cdot 10^{-7} \text{ H m}^{-1}. \end{aligned} \quad (138)$$

(a) The first time  $k_m = m_p r_p$  appears is in the derivation of the Planck relation and reduced Planck constant,  $\hbar = k_m c$  (96), the second time in the Compton wavelength,  $r = k_m / m$  (107). And now,  $k_m$  appears as a component of  $k_e$ , and, therefore,  $\varepsilon_0$  and  $\mu_0$ , which are defined in terms of  $k_e$ .

- (b) It is an open question why  $\frac{k_m}{q_p^2}$  seems to equal  $1.0 \cdot 10^{-7}$  exactly.
14. The fine structure constant,  $\alpha$  was derived from the ratio of two subtypes of charge force that reduces to ratio of the square of the subtypes  $\alpha = q_e^2/q_p^2 \approx 0.0072973526$  (106), which is the empirical CODATA value [14].

- (a) The CODATA electron coupling version of the fine structure constant,  $\alpha$  is defined as:  $\alpha = q_e^2/4\pi\epsilon_0\hbar c = q_e^2/2\epsilon_0\hbar c$  [14]. The following steps show that the CODATA definition reduces to the ratio-derived equation:

$$\begin{aligned}\epsilon_0 &:= 1/4\pi k_e = 1/(4\pi(c_q^2 c_t^2/c_m)) \quad \wedge \quad \hbar = k_m c_t \quad \wedge \quad h = 2\pi\hbar \\ \Rightarrow \quad \epsilon_0 \hbar c &= 2\pi k_m c_t^2 / (4\pi(c_q^2/c_m)c_t^2) = k_m / (2(c_q^2/c_m)) \\ &= m_p r_p / (2((r_p/q_p)^2 / (r_p/m_p))) = q_p^2/2. \quad (139)\end{aligned}$$

$$\alpha = q_e^2/2\epsilon_0\hbar c \quad \wedge \quad \epsilon_0 \hbar c = q_p^2/2 \quad \Rightarrow \quad \alpha = q_e^2/2(q_p^2/2) = q_e^2/q_p^2. \quad (140)$$

- (b) The fine structure electron coupling constant is the ratio of electron static charge force to the propagating quantum charge (photon/electromagnetic) wave force, caused by a moving charged particle.
- (c) Other fine structure constants can also be expressed more simply as the ratios of two subtypes of forces, for example, an electron gravity coupling constant can be expressed as the ratio of the rest electron mass to a Planck mass unit:  
 $\alpha_{G_m} = m_e^2/m_p^2$ .

15. Empirical and hypothesized laws of physics use an *opaque* constant,  $K$ , that is defined to make an equation, where the units balance,  $g = Kf(r, t, \dots)$ . The opacity has led to the *incorrect* assumptions of those constants,  $K$ , being fundamental (atomic) constants.

In this article, some opaque constants are derived directly from (composed of) the ratios: gravity,  $G = c_m c_t^2$  (59), charge,  $k_e = c_q^2 c_t^2 / c_m$  (74), and Planck  $h = k_m c_t$  (97).  $\epsilon_0 = 1/4\pi k_e = 1/4\pi c_m / ((c_q^2/c_m)c_t^2)$  (79) and  $\mu_0 = 4\pi k_e / c_t^2 = 4\pi c_q^2 / c_m$  (85).

And the quantum extensions to: Schwarzschild's gravitational time dilation (126) Newton's gravity force (130), and Coulomb's charge force show, that where the quantum effects become measurable, the constants  $G$ ,  $k_e$ ,  $\epsilon_0$ , and  $\mu_0$  no longer exist (are no longer valid).

Therefore,  $G$ ,  $k_e$ ,  $\varepsilon_0$ ,  $\mu_0$ , and  $h$  are **not** fundamental constants.

16. Constants that use notions of temperature, pressure, mole, etc, for example, the Avogadro and Boltzmann constants [14], are probably not possible to define solely in terms of the Planck units and unit ratios because statistical mechanics is also a component.
17. The derivations of:  $\nabla \cdot \mathbf{g} = -4\pi G\rho$  from  $\mathbf{g} = 2\pi Gm/|\tilde{\mathbf{r}}|^2$  (60),  $\nabla \cdot \mathbf{E} = -\rho/\varepsilon_0$  from  $\mathbf{E} = 2\pi k_e q/|\tilde{\mathbf{r}}|^2$  (79), and  $\partial \mathbf{B}/\partial t = -\mu_0 \rho$  from  $\mathbf{B} = 2\pi k_e q/|\tilde{\mathbf{r}}|^2$  (88), show that the use of mass and charge density,  $\rho$ , are unnecessary complications that obfuscates the pattern,  $\nabla \cdot f(x, r) = -2k_x x/|\tilde{\mathbf{r}}|^3$ , being derived from the inverse square pattern,  $f(x, r) = k_x x/|\tilde{\mathbf{r}}|^2$ . And the energy density in the stress-energy tensor,  $T_{\mu,\nu}$ , in Einstein's field equations [3] also obfuscates the inverse square assumption.
18. Einstein's relativity equations: assume the Lorentz transformations, assume the laws of physics are same at each coordinate point, assume the notion of light, and assume that the speed of light is the same at each coordinate point [2][3].

The derivations, in this article, were made without those assumptions (does not even require the notion of light). Assuming Cartesian coordinates at each coordinate point creates unit ratios, where all equations (laws) derived from the unit ratios must be the same at each coordinate point. Deriving numeric values for the ratios assumes that the ratio,  $c_t = r_p/t_p$ , is the maximum speed.

19. The derivation of the magnetic field from the ratios and special relativity (82) shows that a magnetic field,  $\mathbf{B}$ , is the spacetime bend of the electric field,  $\mathbf{E}$  caused by relativistic charged particle angular (orbital or spin) velocities.
20. The quantum extensions to: Schwarzschild's gravitational time dilation (126), black hole metric (70), Newton's gravity force (130), and Coulomb's charge force (133) make quantifiable predictions:

- (a) For gravity,  $\lim_{r \rightarrow 0} F = c^2 m_1 m_2 / k_m$ , and for charge,  $\lim_{r \rightarrow 0} F = 0$ . Finite maximum gravity and charge forces: 1) allows radioactivity without the need for a weak force, 2) finite sloped energy well walls reducing the need for quantum tunneling, and 3) eliminates the problem of forces going to infinity as  $r \rightarrow 0$ .

- (b) The quantum-extended relativity and classic equations reduce to the non-extended relativity and classic equations and constants, where the distance between masses and charges is sufficiently large or the masses and charges sufficiently large that the quantum effects are not measurable. **Note** that  $G$ ,  $k_e$ ,  $\varepsilon_0$ ,  $\mu_0$ , and  $\kappa$  (Einstein's constant, which contains  $G$ ) do not exist (are not valid), where the quantum effects becomes measurable.
- (c) The covariant tensor components, in Einstein's field equations, that had the units  $1/\text{distance}^2$ , will now have the more complex units,  $1/\sqrt{(\text{distance}^4/c_\tau^2) + k_\tau^2}$ ,  $\tau \in \{t, m\}$ .
- (d)  $1/\sqrt{(\text{distance}^4/c_\tau^2) + k_\tau^2}$  implies that as distance  $\rightarrow 0$ , spacetime curvature peaks at the Planck distance,  $r_p$  (101). The ratio of Planck units,  $m_p/r_p^3$ , might indicate a maximum mass density. A finite force (spacetime curvature) and finite mass density would imply that black holes have sizes  $> 0$  (are not singularities). Black hole evaporation might be possible. If there was a "big bang," then it might not have originated from a singularity.

- 
- [1] Copyright © 2012-2025 George M. Van Treeck. Creative Commons Attribution License.
- [2] A. Einstein, *Relativity, The Special and General Theory* (Princeton University Press, 2015).
- [3] H. Weyl, *Space-Time-Matter* (Dover Publications Inc, 1952).
- [4] R. R. Goldberg, *Methods of Real Analysis* (John Wiley and Sons, 1976).
- [5] W. Rudin, *Principles of Mathematical Analysis* (McGraw Hill Education, 1976).
- [6] W. Conradie and V. Goranko, *Logic and Discrete Mathematics* (Wiley, 2015).
- [7] Rocq, Rocq proof assistant (2025), <https://rocq-prover.org/docs>.
- [8] H. Minkowski, *Geometrie der Zahlen* (Chelsea, 1953) reprint.
- [9] I. Newton, *The Mathematical Principles of Natural Philosophy* (Patristic Publishing, 2019) reprint.
- [10] R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics* (California Institute of Technology, 2010) <https://www.feynmanlectures.caltech.edu/>.
- [11] K. Schwarzschild, Über das gravitationsfeld eines massenpunktes nach der einsteinschen theo-

- rie, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften **7**, 198 (1916), <https://archive.org/stream/sitzungsberichte1916deutsch/page/188/mode/2up>.
- [12] S. Antoci and A. Loinger, An english translation of k. schwarzschild's 1916 article: On the gravitational field of a mass point according to einstein's theory, ARXIV.org (1999), <https://arxiv.org/abs/physics/9905030>.
- [13] M. Jain, *Quantum Mechanics: A Textbook for Undergraduates* (PHI Learning Private Limited, New Delhi, India, 2011).
- [14] CODATA, Physical constant values (2022), <https://physics.nist.gov/cuu/Constants/index.html>.
- [15] E. Schrödinger, *Collected Papers On Wave Mechanics* (Minkowski Institute Press, 2020).
- [16] P. Dirac, *The Principles of Quantum Mechanics* (Snowball Publishing, 1957).
- [17] D. E. Joyce, Euclid's elements (1998), <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>.
- [18] E. S. Loomis, *The Pythagorean Proposition* (NCTM, 1968).
- [19] G. D. Birkhoff, A set of postulates for plane geometry (based on scale and protractors), *Annals of Mathematics* **33** (1932).
- [20] D. Hilbert, *The Foundations of Geometry (2cd Ed)* (Chicago: Open Court, 1980) <http://www.gutenberg.org/ebooks/17384>.
- [21] J. M. Lee, *Axiomatic Geometry* (American Mathematical Society, 2010).
- [22] A. Tarski and S. Givant, Tarski's system of geometry, *The Bulletin of Symbolic Logic* **5**, 175 (1999).
- [23] O. Veblen, A system of axioms for geometry, *Trans. Amer. Math. Soc* **5**, 343 (1904), <https://doi.org/10.1090/S0002-9947-1904-1500678-X>.
- [24] A. Bogomolny, *Pythagorean Theorem* (Interactive Mathematics Miscellany and Puzzles, 2010) <http://www.cut-the-knot.org/pythagoras/CalculusProofCorrectedVersion.shtml>.
- [25] J. Zimba, On the possibility of trigonometric proofs of the pythagorean theorem, *Forum Geometricorum* **9**, 275 (2009).
- [26] B. C. Berndt, Ramanujan-100 years old (fashioned) or 100 years new (fangled)?, *The Mathematical Intelligencer* **10** (1988).
- [27] M. Staring, The pythagorean proposition a proof by means of calculus, *Mathematics Magazine* **69**, 45 (1996).