Copyright © 2026 by George M. Van Treeck. All rights reserved.

Mathematical physics made easy(er)

George M. Van Treeck

(Dated: March 30, 2026)

Abstract

Euclidean volume and some distance equations are proved to be instances of ordered combinations (n-tuples) of elements from a list of abstract, disjoint sets. Volume has a permutation property, where that permutation property is proved to limit a set of unordered elements to a cyclic list of at most 3 elements, for example, 3 unordered distance dimensions (height, width, and depth) and 3 unordered non-distance dimensions (time, mass, and charge). Where the dimensions of Euclidean distance, time, mass, and charge are each divided into a countably infinite number of same-sized intervals and i in N, the i-th unit interval length of Euclidean distance corresponds 1-1 to the i-th unit interval lengths of time, mass, and charge. Derivations of well-known gravity, charge, electromagnetic, relativity, quantum physics equations and constants from unit ratios are shorter and simpler. The unit ratios also allow simple quantum extensions to classical and relativity equations. The math proofs are verified in Rocq.

[1]

CONTENTS

Introduction 3

Insufficient reduction 3

First reduction step 4

Second reduction step 6

Volume 8

Euclidean volume 8

Distance 8

Vector inner product 8

Minkowski distance (p-norm) 9

Metric Space 10

Property limiting a set to at most 3 elements 11

Derivation of physics equations and constants 14

The unit ratios 14

G, Newton, Gauss, Poisson gravity laws 15
Special relativity 15
Simple general relativity solutions 16
Schwarzschild's time dilation and metric 17
k_e, ε_0, E, Coulomb and Gauss laws 18
B, μ_0, and Lorentz's law 19
Faraday's law 19
Maxwell's wave equations 20
The fine structure constant, α 21
$\hbar$, h, and Einstein-Planck relation 21
Unit ratio and reduced Planck unit values 22
Compton wavelength, λ 23
Schrödenger's wave equation 23
Dirac's wave equation 24
Total of a type 25
Quantum extension to general relativity 25
Quantum extension to Newton's gravity force 26
Quantum extension to Coulomb's charge force 26

Insights and implications 26

References 32

Keywords: mathematical physics, gravity, charge, electromagnetism, relativity, quantum physics.

INTRODUCTION

Insufficient reduction

The philosophy of reductionism posits that complex phenomena can be reduced to a few simple parts: a few simple objects, relations, operations, etc. But there has been no reduction of gravity, charge, electromagnetism, relativity, and quantum physics equations into a few simple objects and relations.

Instead, the current practice in physics is to either define a complex equation or derive a complex equation from other complex equations, for example: 1) Newton's law of gravity derived from Kepler's laws[2], or from Gauss's law[3], or from Einstein's field equations[4, 5]; 2) Maxwell's electromagnetic wave equations derived from several empirical laws[3]; 3) Einstein defined a complex tensor field equation and a geodesic equation[4, 5]; and 4) Schrödenger defined a complex wave equation[6].

And the definitions and derivations use "reasonable" assumptions. For example, relativity and quantum theory assume: 1) 3 dimensions of space and 2) the laws of physics are the same at every local Euclidean-like coordinate point[5–8]. But integration, topology, measure theory, Hilbert spaces, linear algebra, (pseudo-)Riemann manifolds, etc. have no restriction on the number of dimensions [5, 9–12]. And the properties of real and complex numbers place no restrictions on functions being the same at every Euclidean-like coordinate[9, 10].

Much of physics relies on volume and distance equations. But mathematical analysis defined Euclidean volume and distance equations and moved on to focusing on abstraction and generalization of geometric volume and distance. Examples of abstraction and generalization are: Hilbert space, inner product space, metric space, topological space, and (pseudo-)Riemann surfaces[9, 10, 12]. Mathematical analysts have failed to prove that volume and distance reduce to a few simple set relations.

The first step is to prove that well-known geometry equations are reduced to a few simple relations between countable set elements. The second step is to reduce well-known physics equations to the properties of the geometry set relations and equations.

Each math proof, in this article, has a corresponding formal proof, which has been verified using using the Rocq proof verification system [13], a verification tool used by mathematicians world-wide. The formal proofs are in the Rocq files, "euclidrelations.v" and "threed.v," which are included as ancillary files.

First reduction step

The following is an outline of the first step of proving that well-known geometry equations are reduced to (derived from) a few simple relations between countable set elements:

Let $|x_i|$ be the cardinal of (number of elements in) the abstract, countable set, $x_i \in \{x_1, \cdots, x_n\}$. And let v_c be the number of ordered combinations (n-tuples) of the elements

of $x_1, \cdots, x_n$: $v_c = \prod_{i=1}^{n} |x_i|$. The Euclidean volume equation will be proved to be an instance of of $v_c = \prod_{i=1}^{n} |x_i|$, where each n-tuple corresponds to a value, $\kappa \in \mathbb{R}$:

$$\forall\, v_c, |x_i| \in \{0, \mathbb{N}\} : \;\; v_c = \textstyle\prod_{i=1}^{n} |x_i| \quad \Rightarrow \quad v_c\kappa = (\prod_{i=1}^{n} |x_i|)\kappa = \prod_{i=1}^{n}(|x_i|\kappa^{1/n})$$
$$\Rightarrow \quad \forall\, v = v_c\kappa, \;\; s_i = |x_i|\kappa^{1/n} : \;\; v = \textstyle\prod_{i=1}^{n} s_i. \quad (1)$$

For all $n > 1$, there are many cases where different domain values, $|x_1|, \cdots, |x_n|$, multiplied yield the same range value, v_c. Inferring a domain value, d_c, from v_c, requires an inverse (bijective) function, $d_c = f_n^{-1}(v_c)$ and $v_c = f_n(d_c)$. The simplest bijective case for all n that includes the $n = 1$ case is where $|x_i| = d_c$:

$$v_c = \textstyle\prod_{i=1}^{n} |x_i| = \prod_{i=1}^{n} d_c = d_c^n \quad \Rightarrow \quad v = d^n. \quad (2)$$

A set of n-tuples being the union of disjoint subsets of n-tuples implies that the domain value, d_c, is the inverse function of the number of n-tuples, v_c, where v_c is the sum of the number of subset n-tuples:

$$d_c^n = v_c = \textstyle\sum_{i=1}^{m} v_{c_i} = \sum_{i=1}^{m}(\prod_{j=1}^{n} |x_{i,j}|) \quad \Rightarrow \quad d^n = v = \sum_{i=1}^{m}(\prod_{j=1}^{n} s_{i,j}). \quad (3)$$

Where each $s_{i,j}$ is $\pm$-signed, the $n = 2$ case is the dot product. And extending $s_{i,j}$ to include complex numbers and functions, the $n = 2$ case is the inner product.

And each summed volume, v_{c_i}, also has an inverse (bijective) function, $d_{c_i}^n = v_{c_i}$:

$$d_c^n = v_c = \textstyle\sum_{i=1}^{m} v_{c_i} = \sum_{i=1}^{m} d_{c_i}^n \quad \Rightarrow \quad d^n = \sum_{i=1}^{m} d_i^n. \quad (4)$$

$|d|$ is the p-norm (Minkowski distance)[14], which will be proved to imply the metric space properties[10]. The $n = 2$ case is the Euclidean distance.

Many sets contain unordered elements, where there is no intrinsic property making a set element the first, second, $\cdots$, or last (n-th). For example, the set elements, {height, width, depth} $\equiv$ {depth, width, height} are unordered and the set elements, {time, mass, charge} $\equiv$ {mass, charge, time} are unordered.

Sequencing a set of unordered elements reliably in the same order requires that the elements be placed into a "list," which assigns relative sequential positions, for example, labeling the unordered elements as $x_1, x_2, \cdots x_n$. The list can then be sequenced via successor and predecessor functions, $successor(x_i) = x_{i+1}$ and $predecessor(x_i) = x_{i-1}$. For example,

the volume successor sequence, $v = \prod_{i=1}^{n} s_i$, sequences the corresponding set elements, $x_1 =$ width, $x_2 =$ height, and $x_3 =$ depth, in the same order on each sequencing.

But the union, intersection, multiplication, and addition operations defining the total number of n-tuples and the corresponding volume and distance equations are commutative. The commutative property allows sequencing the *same* totally ordered list of n number of elements in all of the $n!$ possible unique sequences (permutations).

Only a cyclic list, where $successor(x_n) = x_1$ and $predecessor(x_1) = x_n$, allows each element to be the first element of some of those $n!$ permutations. For example, where $n = 3$, the volume cyclic successor sequences are: $v = s_1 \times s_2 \times s_3$. $v = s_2 \times s_3 \times s_1$, $v = s_3 \times s_1 \times s_2$, and the cyclic predecessor sequences are: $v = s_1 \times s_3 \times s_2$. $v = s_2 \times s_1 \times s_3$, $v = s_3 \times s_2 \times s_1$.

The only cyclic lists that allows sequencing all $n!$ permutations, using the cyclic successor and predecessor functions, are where each list element is sequentially adjacent (either an *immediate* cyclic successor or an *immediate* cyclic predecessor) to every other element, referred to, here, as an immediate symmetric cyclic list (ISCL). For example, the permutation, (x_2, x_3, x_1), requires x_1 and x_3 be sequentially adjacent. An ISCL will be proved to contain $n \leq 3$ elements.

Each volume value $\in \mathbb{R}$ and distance an inverse (bijective) functions of volume implies a corresponding distance value$\in \mathbb{R}$ and a distance dimension is $\subseteq \mathbb{R}$. If there is an ISCL of 3 distance dimensions, each $\subseteq \mathbb{R}$, then exists a complement ISCL of at most 3 non-distance dimensions, each $\subseteq \mathbb{R}$.

A 3-dimensional Euclidean distance dimension, $\subseteq \mathbb{R}$, can be related to an ISCL of 3 non-distance dimensions, each $\subseteq \mathbb{R}$, by dividing the Euclidean distance dimension and the 3 non-distance dimensions into countably infinite sets. Where $i \in \mathbb{N}$, the ith interval length of Euclidean distance, corresponds 1-1 to the ith interval lengths of each the 3 non-distance dimensions.

Second reduction step

The following is an outline of the second step of reducing well-known physics equations to the ISCL and property of the Euclidean distance dimension $\subseteq \mathbb{R}$, and non-distance dimensions, each $\subseteq \mathbb{R}$:

1. **ISCL:** $\{r_1, r_2, r_3\}$ is an ISCL of 3 "distance" dimensions and $\{t\ (time), m\ (mass),$

q $(charge)\}$ is the ISCL of 3 "non-distance" dimensions, each dimension $\subseteq \mathbb{R}$. There is a total of 6 unordered dimensions with the $\subseteq \mathbb{R}$ property: $r_1,\ r_2,\ r_3,\ t,\ m,\ q$.

2. **Corresponding lengths:** Where the dimensions of Euclidean distance, time, mass, and charge are each divided into countably infinite sets of same-sized intervals $\subset \mathbb{R}$ and $i \in \mathbb{N}$, the ith unit interval length, r_p of Euclidean distance r, corresponds 1-1 to the ith unit interval lengths: t_p of time t, m_p mass m, and q_p charge q, where "unit" means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

The 1-1 correspondence of unit interval lengths implies the unit ratios: $r = (r_p/t_p)t$, $r = (r_p/m_p)m$, and $r = (r_p/q_p)q$. Letting: $c_t = r_p/t_p$, $c_m = r_p/m_p$, and $c_q = r_p/q_p$, the gravity constant, G, will be shown to be $G = c_m c_t^2$, where c_t is the speed of light.

Algebraic manipulation of the 3 direct proportion ratios yields 3 inverse proportion ratios, $rt = t_p r_p$, $rm = m_p r_p$. and $rq = q_p r_p$. Letting: $k_t = t_p r_p$, $k_m = m_p r_p$, and $k_q = q_p r_p$, the reduced Planck constant, $\hbar$, will be shown to be $\hbar = k_m c_t$.

Note that the quantum units: r_p, t_p, m_p, and q_p are derived from the property:

$$r_p = \sqrt{r_p^2} = \sqrt{c_t k_t} = \sqrt{c_m k_m} = \sqrt{c_q k_q}. \tag{5}$$

And it will be shown that r_p, t_p, m_p, and q_p are the reduced Planck units.

Some published derivations of physics equations are long, with complex math, and many "reasonable" assumptions because the underlying principles are not well understood. As evidence of that assertion, the derivations of well-known gravity, charge, electromagnetic, relativity, quantum physics equations and constants from the unit ratios are shorter, simpler, and use fewer assumptions. All derivations, in this article, are either directly from unit ratios or from equations previously derived from the unit ratios.

The inverse proportion unit ratios are used to add quantum extensions Schwarzschild's gravitational time dilation and black hole metric[15, 16], Newton's gravity force[2], and Coulomb's charge force[3] equations. The quantum extended equations predict that where quantum effects are measurable, the constants: gravity, G, Einstein's $\kappa = 8\pi G/c^4$, charge, k_e, permittivity, ε_0, and permeability, μ_0, do not exist. And the quantum-extended equations also predict that as distance between two point masses and between two point charges go to zero, the forces do not go to infinity.

VOLUME

Euclidean volume

Theorem .1. *Euclidean volume,*

$$\forall\ v_c, d_c, |x_i| \in \{0,\ \mathbb{N}\},\ \ x_i \in \{x_1, \cdots, x_n\},\ v_c = \prod_{i=1}^{n} |x_i| \ \Rightarrow\ v = \prod_{i=1}^{n} s_i, \quad s_i, v \in \mathbb{R}. \tag{6}$$

The formal proof, "Euclidean_volume," is in the Rocq file, euclidrelations.v.

Proof.

$$\forall v_c, |x_i| \in \{0, \mathbb{N}\},\ \ \kappa \in \mathbb{R}:\ \ v_c = \prod_{i=1}^{n} |x_i| \quad \Leftrightarrow \quad v_c\kappa = (\prod_{i=1}^{n} |x_i|)\kappa = \prod_{i=1}^{n}(|x_i|\kappa^{1/n}) \tag{7}$$

$$\exists\ v, s_i \in \mathbb{R}:\ \ v = v_c\kappa\ \ \wedge\ \ s_i = |x_i|\kappa^{1/n}\ \ \wedge\ \ v_c\kappa = \prod_{i=1}^{n}(|x_i|\kappa^{1/n})\ \ \Rightarrow\ \ v = \prod_{i=1}^{n} s_i. \quad \square \tag{8}$$

Lemma .2. *The number of n-tuples, v_c, is the sum of the number of n-tuples, v_{c_i}, in each subset of n-tuples, implies a volume is the sum of volumes,*

$$v_c = \sum_{i=1}^{m} v_{c_i} \quad \Rightarrow \quad v = \sum_{i=1}^{m} v_i, \quad v, v_i \in \mathbb{R}.$$

The formal proof, "sum_of _volumes," is in the Rocq file, euclidrelations.v.

Proof. From the condition of this theorem and the Euclidean volume theorem (.1):

$$\forall\ \kappa \in \mathbb{R}:\ \ v_c = \sum_{i=1}^{m} v_{c_i} \quad \Rightarrow \quad v_c\kappa = (\sum_{i=1}^{m} v_{c_i})\kappa = \sum_{i=1}^{m}(v_{c_i}\kappa). \tag{9}$$

$$v_c\kappa = \sum_{i=1}^{m}(v_{c_i}\kappa) \quad \wedge \quad v = v_c\kappa \quad \wedge \quad v_i = v_{c_i}\kappa \quad \Rightarrow \quad v = \sum_{i=1}^{m} v_i. \quad \square \tag{10}$$

DISTANCE

Definition .3. Bijective, countable domain value, d_c:

$$\exists\ v_c, d_c, |x_i| \in \{0,\ \mathbb{N}\},\ \ x_i \in \{x_1, \cdots, x_n\}, \quad v_c = \prod_{i=1}^{n} |x_i| = \prod_{i=1}^{n} d_c = d_c^n. \tag{11}$$

Vector inner product

Theorem .4. *Sum of volumes distance:*

$$d_c^n = v_c = \sum_{i=1}^{m} v_{c_i} \quad \Rightarrow \quad d^n = v = \sum_{i=1}^{m}(\prod_{j=1}^{n} s_{i_j}).$$

The formal proof, "sum_of _volumes _distance," is in the Rocq file, euclidrelations.v.

Proof. Multiply both sides of the assumption by κ and apply the Euclidean volume proof (.1) to v_i:

$$v_c = \sum_{i=1}^{m} v_{c_i} \quad \Rightarrow \quad \forall\, \kappa \in \mathbb{R}: \;\; v_c\kappa = (\sum_{i=1}^{m} v_{c_i})\kappa = \sum_{i=1}^{m}(v_{c_i}\kappa). \tag{12}$$

$$v_c\kappa == \sum_{i=1}^{m}(v_{c_i}\kappa) \quad \wedge \quad \exists\, v, v_i \in \mathbb{R}: \;\; v = v_c\kappa \quad \wedge \quad v_i = v_{c_i}\kappa$$

$$\Rightarrow \quad v = \sum_{i=1}^{m}(\prod_{j=1}^{n} v_i. \tag{13}$$

$$d_c^n = v_c \quad \Rightarrow \quad d_c^n\kappa = (d_c\kappa^{1/n})^n = v_c\kappa. \tag{14}$$

$$\exists\, d, v \in \mathbb{R} : d = d_c\kappa^{1/n} \quad \wedge \quad v = v_c\kappa \quad \wedge \quad (d_c\kappa^{1/n})^n = v_c\kappa \quad \Rightarrow \quad d^n = v. \tag{15}$$

$$d^n = v \quad \wedge \quad v = \sum_{i=1}^{m}(\prod_{j=1}^{n} v_i \quad \Rightarrow \quad d^n = v = \sum_{i=1}^{m}(\prod_{j=1}^{n} v_i. \tag{16}$$

$$d^n = v = \sum_{i=1}^{m}(\prod_{j=1}^{n} v_i \quad \wedge \quad v_i = \prod_{j=1}^{n} s_{i,j} \quad \Rightarrow \quad d^n = v = \sum_{i=1}^{m}(\prod_{j=1}^{n} s_{i,j}). \quad \square \tag{17}$$

Note: $\exists\, \kappa_i \in \{\kappa_1, \cdots, \kappa_n\}$ such that $\kappa = \prod_{j=i}^{n} \kappa_i$. And where $\kappa_i < 0$, the corresponding $s_{i,j} < 0$. Therefore, the $n = 2$ case of the sum of volumes distance is the dot product. The dot product is generalized to the inner product by also allowing each $s_{i,j}$ term to be a complex number or function returning a real or complex number.

Minkowski distance (p-norm)

Theorem .5. *Minkowski distance (p-norm):*

$$d_c^n = v_c = \sum_{i=1}^{m} v_{c_i} = \sum_{i=1}^{m} d_{c_i}^n \quad \Leftrightarrow \quad d^n = \sum_{i=1}^{m} d_i^n.$$

The formal proof, "Minkowski_distance," is in the Rocq file, euclidrelations.v.

Proof. From the sum of volumes distance theorem .4:

$$d_c^n = v_c = \sum_{i=1}^{m} v_{c_i} \quad \Rightarrow \quad d^n = v = \sum_{i=1}^{m} v_i \tag{18}$$

$$d^n = v = \sum_{i=1}^{m} v_i \quad \wedge \quad d_i^n = v_i \Rightarrow \quad d^n = \sum_{i=1}^{m} d_i^n \quad \square \tag{19}$$

Theorem .6. *Distance triangle inequality*

$$\forall\, n \in \mathbb{N}, \;\; v_a, v_b \geq 0: \;\; (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}.$$

The formal proof, distance_inequality, is in the Rocq file, euclidrelations.v.

Proof. Expand $(v_a^{1/n} + v_b^{1/n})^n$ using the binomial expansion:

$$\forall\, v_a, v_b \geq 0: \quad v_a + v_b \leq v_a + v_b + \sum_{i=1}^{n} \binom{n}{k} (v_a^{1/n})^{n-k} (v_b^{1/n})^k + \sum_{i=1}^{n} \binom{n}{k} (v_a^{1/n})^k (v_b^{1/n})^{n-k} = (v_a^{1/n} + v_b^{1/n})^n. \quad (20)$$

Take the nth root of both sides of the inequality 20:

$$\forall\, v_a, v_b \geq 0,\ n \in \mathbb{N}: \quad v_a + v_b \leq (v_a^{1/n} + v_b^{1/n})^n \quad \Rightarrow \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}. \quad (21)$$

□

Theorem .7. *Distance sum inequality*

$$\forall\, m, n \in \mathbb{N},\ a_i, b_i \geq 0 : (\textstyle\sum_{i=1}^{m} (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^{m} a_i^n)^{1/n} + (\sum_{i=1}^{m} b_i^n)^{1/n}.$$

The formal proof, distance_sum_inequality, is in the Rocq file, euclidrelations.v.

Proof. Apply the distance triangle inequality (.6):

$$\forall\, m, n \in \mathbb{N},\ v_a, v_b \geq 0: \ v_a = \textstyle\sum_{i=1}^{m} a_i^n \quad \wedge \quad v_b = \sum_{i=1}^{m} b_i^n \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}$$
$$\Rightarrow \quad ((\textstyle\sum_{i=1}^{m} a_i^n) + (\sum_{i=1}^{m} b_i^n))^{1/n} = (\sum_{i=1}^{m} (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^{m} a_i^n)^{1/n} + (\sum_{i=1}^{m} b_i^n)^{1/n}. \quad \square \quad (22)$$

Metric Space

All Minkowski distances (p-norms) imply the metric space properties. The formal proofs: triangle_inequality, symmetry, non_negativity, and identity_of_indiscernibles are in the Rocq file, euclidrelations.v.

Theorem .8. *Triangle Inequality:*

$$d(s_1, s_2) = (\sum_{i=1}^{2} s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k$$
$$\Rightarrow \quad d(u, w) = (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w).$$

Proof. $\forall\, p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k$:

$$(u^p + w^p)^{1/p} \leq ((u^p + w^p) + 2v^p)^{1/p} = ((u^p + v^p) + (v^p + w^p))^{1/p}. \quad (23)$$

Apply the distance triangle inequality (.6) to the inequality 23:

$$(u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}$$
$$\wedge \quad v_a = u^p + v^p \quad \wedge \quad v_b = v^p + w^p$$
$$\Rightarrow \quad (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p}$$
$$\Rightarrow \quad d(u, w) = (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \quad \square \tag{24}$$

Theorem .9. *Symmetry:*
$d(s_1, s_2) = (\sum_{i=1}^{2} s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad v = s_2 \quad \Rightarrow \quad d(u, v) = d(v, u).$

Proof. By the commutative law of addition:

$$\forall\, p \,:\, p \geq 1, \quad d(s_1, s_2) = (\textstyle\sum_{i=1}^{2} s_i^p)^{1/p} = (s_1^p + s_2^p)^{1/p}$$
$$\Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} = (v^p + u^p)^{1/p} = d(v, u). \quad \square \tag{25}$$

Theorem .10. *Non-negativity:*

$$d(s_1, s_2) = (\textstyle\sum_{i=1}^{2} s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k \quad \Rightarrow \quad d(u, w) \geq 0.$$

Proof. By definition, the length of an interval is always ≥ 0:

$$\forall\, [a_1, b_1],\ [a_2, b_2], \quad u = b_1 - a_1,\ v = b_2 - a_2, \quad \Rightarrow \quad u \geq 0,\ v \geq 0. \tag{26}$$
$$p \geq 1,\ \ u, v \geq 0 \quad \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} \geq 0. \quad \square \tag{27}$$

Theorem .11. *Identity of Indiscernibles:* $d(u, u) = 0$.

Proof. From the non-negativity property (.10):

$$d(u, w) \geq 0 \quad \wedge \quad d(u, v) \geq 0 \quad \wedge \quad d(v, w) \geq 0 \tag{28}$$
$$\Rightarrow \quad \exists\, d(u, w) = d(u, v) = d(v, w) = 0. \tag{29}$$
$$d(u, w) = d(v, w) = 0 \quad \Rightarrow \quad u = v. \tag{30}$$
$$d(u, v) = 0 \quad \wedge \quad u = v \quad \Rightarrow \quad d(u, u) = 0. \quad \square \tag{31}$$

Property limiting a set to at most 3 elements

The following definitions and proof use first order logic. A Horn clause-like expression is used, here, to make the proof easier to read. By convention, the proof goal, in a Horn clause,

is on the left side of the implication sign ($\leftarrow$) and supporting facts are on the right side. The formal proofs in the Rocq file threed.v are: adj111, adj122, adj212, adj123, adj133, adj233, adj213, adj313, adj323, and not_all_mutually_adjacent_gt_3.

Definition .12. Immediate Cyclic Successor of m is n:

$$\forall\, x_m, x_n \in \{x_1, \cdots, x_{setsize}\} :$$
$$Successor(m, n, setsize) \quad \leftarrow \quad (m = setsize \wedge n = 1) \quad \vee \quad (n = m + 1 \leq setsize). \tag{32}$$

Definition .13. Immediate Cyclic Predecessor of m is n:

$$\forall\, x_m, x_n \in \{x_1, \cdots, x_{setsize}\} :$$
$$Predecessor(m, n, setsize) \quad \leftarrow \quad (m = 1 \wedge n = setsize) \quad \vee \quad (n = m - 1 \geq 1). \tag{33}$$

Definition .14. Adjacent: element m is sequentially adjacent to element n if the immediate cyclic successor of m is n or the immediate cyclic predecessor of m is n. Notionally:

$$\forall\, x_m, x_n \in \{x_1, \cdots, x_{setsize}\} :$$
$$Adjacent(m, n, setsize) \leftarrow Successor(m, n, setsize) \;\vee\; Predecessor(m, n, setsize). \tag{34}$$

Definition .15. Immediate Symmetric Cylic List (ISCL), where every set element is sequentially adjacent to every other element):

$$\forall\, x_m, x_n \in \{x_1, \cdots, x_{setsize}\} : \quad Adjacent(m, n, setsize). \tag{35}$$

Theorem .16. *An ISCL is limited to at most 3 elements.*

$$\forall\, x_m, x_n \in \{x_1, \cdots, x_{setsize}\} : \quad Adjacent(m, n, setsize) \quad \Rightarrow \quad setsize \leq 3. \tag{36}$$

Proof.

Every element is adjacent to every other element, where $setsize \in \{1,\ 2,\ 3\}$:

$$Adjacent(1,1,1) \leftarrow Successor(1,1,1) \leftarrow (m = setsize \wedge n = 1). \tag{37}$$

$$Adjacent(1,2,2) \leftarrow Successor(1,2,2) \leftarrow (n = m + 1 \leq setsize). \tag{38}$$

$$Adjacent(1,2,3) \leftarrow Successor(1,2,3) \leftarrow (n = m + 1 \leq setsize). \tag{39}$$

$$Adjacent(2,1,3) \leftarrow Predecessor(2,1,3) \leftarrow (n = m - 1 \geq 1). \tag{40}$$

$$Adjacent(3,1,3) \leftarrow Successor(3,1,3) \leftarrow (n = setsize \wedge m = 1). \tag{41}$$

$$Adjacent(1,3,3) \leftarrow Predecessor(1,3,3) \leftarrow (m = 1 \wedge n = setsize). \tag{42}$$

$$Adjacent(2,3,3) \leftarrow Successor(2,3,3) \leftarrow (n = m + 1 \leq setsize). \tag{43}$$

$$Adjacent(3,2,3) \leftarrow Predecessor(3,2,3) \leftarrow (n = m - 1 \geq 1). \tag{44}$$

Element 2 is the only immediate successor of element 1 for all $setsize \geq 3$, which implies element 3 is not ($\neg$) an immediate successor of element 1 for all $setsize \geq 3$:

$$\neg Successor(1,3,setsize \geq 3) \leftarrow \\ Successor(1,2,setsize \geq 3) \leftarrow (n = m + 1 \leq setsize). \tag{45}$$

Element $n = setsize > 3$ is the only immediate predecessor of element 1, which implies element 3 is not ($\neg$) an immediate predecessor of element 1 for all $setsize > 3$:

$$\neg Predecessor(1,3,setsize \geq 3) \leftarrow Predecessor(1,setsize,setsize > 3) \leftarrow \\ (m = 1 \wedge n = setsize > 3). \tag{46}$$

For all $setsize > 3$, some elements are not ($\neg$) sequentially adjacent to every other element (not immediate symmetric):

$$\neg Adjacent(1,3,setsize > 3) \leftarrow \\ \neg Successor(1,3,setsize > 3) \quad \wedge \quad \neg Predecessor(1,3,setsize > 3). \quad \square \tag{47}$$

The Symmetric goal matches the Adjacent goals 37 and fails for all "setsize" greater than three.

DERIVATION OF PHYSICS EQUATIONS AND CONSTANTS

The unit ratios

Application of the ISCL (.16) and properties of Euclidean geometry (.1) to physics:

1. **ISCL:** $\{r_1, r_2, r_3\}$ is an ISCL of 3 "distance" dimensions and $\{t\ (time), m\ (mass), q\ (charge)\}$ is the ISCL of 3 "non-distance" dimensions, each dimension $\subseteq \mathbb{R}$. There is a total of 6 unordered dimensions with the $\subseteq \mathbb{R}$ property: $r_1,\ r_2,\ r_3,\ t,\ m,\ q$.

2. **Corresponding lengths:** Where the dimensions of Euclidean distance, time, mass, and charge are each divided into countably infinite sets of same-sized intervals $\subset \mathbb{R}$ and $i \in \mathbb{N}$, the ith unit interval length, r_p of Euclidean distance r, corresponds 1-1 to the ith unit interval lengths: t_p of time t, m_p mass m, and q_p charge q, where "unit" means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

The 1-1 correspondence of unit interval lengths implies unit ratios of r_p to t_p, m_p, and q_p. A corresponding number of units can be expressed as, $r/r_p = \tau/\tau_p$, where $\tau \in \{t, m, q\}$. Multiplying both sides by r_p yields 3 direct proportion ratios:

$$\forall\, r_p, \tau_p, r, \tau \subseteq \mathbb{R},\ r_p, \tau_p\ \textbf{constants},\ \tau \in \{t, m, q\} :\ r/r_p = \tau/\tau_p \quad \Rightarrow \quad r = (r_p/\tau_p)/\tau$$
$$\Rightarrow \quad r = (r_p/t_p)t \quad \wedge \quad r = (r_p/m_p)m \quad \wedge \quad r = (r_p/q_p)q. \tag{48}$$

And for conciseness, let:

$$r_p/t_p = c_t \quad \wedge \quad r_p/m_p = c_m \quad \wedge \quad r_p/q_p = c_q. \tag{49}$$

Algebraic manipulation of the 3 direct proportion ratios yields 3 inverse proportion ratios:

$$r/t = r_p/t_p \quad \wedge \quad r/m = r_p/m_p \quad \Rightarrow \quad (r/t)/(r/m) = (r_p/t_p)/(r_p/m_p)$$
$$\Rightarrow \quad (mr)/(tr) = (m_p r_p)/(t_p r_p) \quad \Rightarrow \quad \exists\, r, t, m \in \mathbb{R} :\ mr = m_p r_p, \quad tr = t_p r_p. \tag{50}$$

$$r/t = r_p/t_p \quad \wedge \quad r/q = r_p/q_p \quad \Rightarrow \quad (r/t)/(r/q) = (r_p/t_p)/(r_p/q_p)$$
$$\Rightarrow \quad (qr)/(tr) = (q_p r_p)/(t_p r_p) \quad \Rightarrow \quad \exists\, r, t, q \in \mathbb{R} :\ qr = q_p r_p, \quad tr = t_p r_p. \tag{51}$$

And for conciseness, let:

$$t_p r_p = k_t \quad \wedge \quad m_p r_p = k_m \quad \wedge \quad q_p r_p = k_q. \tag{52}$$

G, Newton, Gauss, Poisson gravity laws

From equations 49, linear acceleration, r/t^2, can be related to mass, m, and distance, r:

$$r = c_m m \;\wedge\; r = c_t t \;\Rightarrow\; r/(c_t t)^2 = c_m m/r^2 \;\Rightarrow\; r/t^2 = (c_m c_t^2)m/r^2 = Gm/r^2, \tag{53}$$

where $G = c_m c_t^2$, conforms to the SI units: $m^3 \cdot kg^{-1} \cdot s^{-2}$ [2].

Newton's law follows from multiplying both sides of equation 53 by m:

$$r/t^2 = Gm/r^2 \;\Leftrightarrow\; F \stackrel{\text{def}}{=} mr/t^2 = Gm^2/r^2. \tag{54}$$

$$F = Gm^2/r^2 \;\wedge\; \forall\, m \in \mathbb{R} :\; \exists\, m_1, m_2 \in \mathbb{R} :\; m_1 m_2 = m^2 \quad\Rightarrow\quad F = Gm_1 m_2/r^2. \tag{55}$$

A new interpretation of Gauss's and Poisson's gravity laws is presented here. Equation 53 relates linear acceleration, r/t^2, to mass and distance. Gauss's gravity field, $\mathbf{g}$, and Poisson's gravity field, $-\nabla\Phi(\tilde{\mathbf{r}}, t)$ is the angular (orbital) acceleration, $2\pi r/t^2$. Applying equation 53:

$$r/t^2 = Gm/|\tilde{\mathbf{r}}|^2 \;\wedge\; \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi r/t^2 \quad\Rightarrow\quad \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi Gm/|\tilde{\mathbf{r}}|^2 \tag{56}$$

$$\mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = -2\pi Gm/|\tilde{\mathbf{r}}|^2 \quad\Rightarrow\quad \nabla\cdot\mathbf{g} = \nabla^2\Phi(\tilde{\mathbf{r}}, t) = -4\pi Gm/|\tilde{\mathbf{r}}|^3 = -4\pi G\rho, \tag{57}$$

where $\rho = m/|\tilde{\mathbf{r}}|^3$ is the mass density:

Special relativity

$$\forall\, \tau \in \{t, m, q\},\; r^2 = r'^2 + r_v^2,\; \exists\, \mu, \nu :\; r = \mu\tau \quad\wedge\quad r_v = \nu\tau \quad\Rightarrow\quad (\mu\tau)^2 = r'^2 + (\nu\tau)^2$$
$$\Rightarrow\quad r' = \sqrt{(\mu\tau)^2 - (\nu\tau)^2} = \mu\tau\sqrt{1 - (\nu/\mu)^2}. \tag{58}$$

Local frame distance, r', contracts relative to a distant observer frame distance, r, as $\nu \to \mu$:

$$r' = \mu\tau\sqrt{1 - (\nu/\mu)^2} \quad\wedge\quad \mu\tau = r \quad\Rightarrow\quad r' = r\sqrt{1 - (\nu/\mu)^2}. \tag{59}$$

A distant observer frame type, τ, dilates relative to the local observer frame type, τ', as $\nu \to \mu$:

$$\mu\tau = r'/\sqrt{1 - (\nu/\mu)^2} \quad\wedge\quad r' = \mu\tau \quad\Rightarrow\quad \tau = \tau'/\sqrt{1 - (\nu/\mu)^2}. \tag{60}$$

Where τ is type, time, the space-like flat Minkowski spacetime event interval is:

$$\mathrm{d}r^2 = \mathrm{d}r'^2 + \mathrm{d}r_v^2 \quad\wedge\quad \mathrm{d}r_v^2 = \mathrm{d}r_1^2 + \mathrm{d}r_2^2 + \mathrm{d}r_3^2 \quad\wedge\quad \mathrm{d}(\mu\tau) = \mathrm{d}r$$
$$\Rightarrow\; \mathrm{d}r'^2 = \mathrm{d}(\mu\tau)^2 - \mathrm{d}r_1^2 - \mathrm{d}r_2^2 - \mathrm{d}r_3^2. \tag{61}$$

Simple general relativity solutions

The Einstein field equation (EFE) is:

$$G_{\mu,\nu} + \Lambda g_{\mu,\nu} = \mathbf{R} + \frac{1}{2} R g_{\mu,\nu} + \Lambda g_{\mu,\nu} = \kappa T_{\mu,\nu}, \tag{62}$$

where: $G_{\mu,\nu}$ is Einstein's tensor, $\mathbf{R}$ is the Ricci curvatature, R is the scalar curvature, $g_{\mu,\nu}$ is the metric tensor, Λ is a cosmological constant, $\kappa = 8\pi G/c^4$ is Einstein's constant, and $T_{\mu,\nu}$ is the stress-energy-momentum tensor [5, 7].

The EFE is nonlinear, which has made determining the measure (metric) of the spacetime curvature, $g_{\mu,\nu}$, from the distribution of the stress-energy-momentum, $T_{\mu,\nu}$ very difficult. However, determining $g_{\mu,\nu}$ is very simple by noting that: 1) The infinitesimal space near every local coordinate point in the EFE is Euclidean-like[5, 7]; 2) The unit ratios are with respect to Euclidean distance and, therefore, all physics equations (laws) derived from the unit ratios are the same at every Euclidean-like local coordinate point; and 3) The spacetime curvature is specified by one function for each diagonal component of the metric, $g_{\mu,\nu}$ [7].

This leads to the following simple steps to solve for the metric tensor, $g_{\mu,\nu}$, independent of the EFE. The next subsection contains an example of using these steps for a much shorter and simpler derivation of Schwarzschild's black hole metric [15, 16].

Step 1) Use the unit ratios and other equations derived from the ratios, for example, the special relativity equations (59) and the equation defining the constant, G (53), to derive functions to be assigned to the diagonal components of $g_{\nu,\mu}$.

Step 2) The 4D flat spacetime interval equation is an instance of the 2D equation (61), $\mathrm{d}r'^2 = \mathrm{d}(ct)^2 - \mathrm{d}r_v^2$. Derive the metric first as a 2D metric and expand later to a 4D metric.

(Optional step 2d) The 2D metric tensor allows using the much simpler 2D Ricci curvature and scalar curvature. And the 2D tensors reduce the number of independent equations to solve. The 2D solutions can next be used to set constraints on the solutions in the 4D tensors.

Step 3) One simple method to translate from 2D to 4D is to use spherical coordinates, where the first 2 diagonal components of $g_{\nu,\mu}$ remain unchanged and the other 2 diagonal components are functions of the angles, longitude, θ, and latitude, ϕ.

Schwarzschild's time dilation and metric

From equations 59 and 48:

$$\sqrt{1-(v^2/c^2)} = \sqrt{1-(1)(v^2/c^2)} \quad \wedge \quad c_m m/r = 1$$
$$\Rightarrow \quad \sqrt{1-(v^2/c^2)} = \sqrt{1-c_m m v^2/rc^2}. \quad (63)$$

Where v_{escape} is the escape velocity:

$$\sqrt{1-(v^2/c^2)} = \sqrt{1-c_m m v^2/rc^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2$$
$$\Rightarrow \quad \sqrt{1-(v^2/c^2)} = \sqrt{1-2c_m m v_{escape}^2/rc^2}. \quad (64)$$

For a photon, the escape velocity, $v_{escape} = c$.

$$\sqrt{1-(v^2/c^2)} = \sqrt{1-2c_m m v_{escape}^2/rc^2}. \quad \wedge \quad v_{escape} = c$$
$$\Rightarrow \quad \sqrt{1-(v^2/c^2)} = \sqrt{1-2c_m mc^2/rc^2}. \quad (65)$$

Combining equation 65 with the derivation of G (53):

$$c_m c^2 = G \wedge \sqrt{1-(v^2/c^2)} = \sqrt{1-2c_m mc^2/rc^2} \Rightarrow \sqrt{1-(v^2/c^2)} = \sqrt{1-2Gm/rc^2}. \quad (66)$$

Combining equation 66 with equation 60 yields Schwarzschild's gravitational time dilation:

$$\sqrt{1-(v^2/c^2)} = \sqrt{1-2Gm/rc^2} \wedge t' = t\sqrt{1-(v^2/c^2)} \Rightarrow t' = t\sqrt{1-2Gm/rc^2}. \quad (67)$$

From equations 59 and 66, where Schwarzschild defined the black hole event horizon radius as $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$ [15, 16]:

$$r' = r\sqrt{1-(v/c)^2} = r\sqrt{1-2Gm/rc^2} \quad \wedge \quad \alpha \stackrel{\text{def}}{=} 2Gm/c^2 \quad \Rightarrow \quad r' = r\sqrt{1-\alpha/r}. \quad (68)$$

Applying equation 68 to the time-like spacetime interval equation 61, where $\mathrm{d}s^2 < 0$:

$$r' = r\sqrt{1-\alpha/r} \quad \wedge \quad \mathrm{d}s^2 = \mathrm{d}r'^2 - \mathrm{d}r^2$$
$$\Rightarrow \quad \mathrm{d}s^2 = (\sqrt{1-\alpha/r}\mathrm{d}r)^2 - (\mathrm{d}r'/\sqrt{1-\alpha/r})^2 = (1-\alpha/r)\mathrm{d}r^2 - (1-\alpha/r)^{-1}\mathrm{d}r'^2. \quad (69)$$
$$\mathrm{d}s^2 = (1-\alpha/r)\mathrm{d}r^2 - (1-\alpha/r)^{-1}\mathrm{d}r'^2 \quad \wedge \quad \mathrm{d}r = \mathrm{d}(ct) \quad \wedge \quad c = 1 \quad \wedge$$
$$\lim_{\mathrm{d}s\to 0} \mathrm{d}r' = \lim_{\mathrm{d}s\to 0} \mathrm{d}r \quad \Rightarrow \quad \lim_{\mathrm{d}s\to 0} \mathrm{d}s^2 = (1-\alpha/r)\mathrm{d}t^2 - (1-\alpha/r)^{-1}\mathrm{d}r^2. \quad (70)$$

Using spherical coordinates to translate from 2D to 4D yields the $+ - --$ form of Schwarzschild's black hole metric [15, 16]:

$$\begin{aligned} \mathrm{d}s^2 &= (1-\alpha/r)\mathrm{d}t^2 - (1-\alpha/r)^{-1}\mathrm{d}r^2 \\ &\Rightarrow \quad \mathrm{d}s^2 = (1-\alpha/r)\mathrm{d}t^2 - (1-\alpha/r)^{-1}\mathrm{d}r^2 - r^2(\mathrm{d}\theta^2 + sin^2\theta\mathrm{d}\phi^2) \\ &\Rightarrow \quad g_{\mu,\nu} = diag[(1-\alpha/r),\ -(1-\alpha/r)^{-1},\ -r^2(\mathrm{d}\theta^2),\ -r^2(sin^2\theta\mathrm{d}\phi^2)]. \end{aligned} \tag{71}$$

k_e, ε_0, E, Coulomb and Gauss laws

From equations 49, linear acceleration, r/t^2, can be related to charge, q, and distance, r:

$$r = c_q q \ \wedge\ r = c_t t \ \Rightarrow\ r/(c_t t)^2 = c_q q/r^2 \ \Rightarrow\ r/t^2 = (c_q c_t^2)q/r^2. \tag{72}$$

$$r = c_m m = c_q q \quad \Rightarrow \quad m = (c_q/c_m)q. \tag{73}$$

Combining equations 72 and 73 yields Coulombs's law:

$$r/t^2 = (c_q c_t^2)q/r^2 \ \wedge\ m = (c_q/c_m)q \ \Rightarrow\ F \stackrel{\text{def}}{=} mr/t^2 = (c_q^2 c_t^2/c_m)q^2/r^2 = k_e q^2/r^2. \tag{74}$$

where $k_e = c_q^2 c_t^2/c_m$, conforms to the SI units: $kg \cdot m^3 \cdot s^{-2} \cdot C^{-2} = N \cdot m^2 \cdot C^{-2}$ [3].

$$\forall\, q \in \mathbb{R}\ \exists\, q_1, q_2 \in \mathbb{R}:\ q_1 q_2 = q^2 \quad \wedge \quad F = k_e q^2/r^2 \quad \Rightarrow \quad F = k_e q_1 q_2/r^2. \tag{75}$$

A new interpretation of Gauss's electric field law is presented here. Equation 72 relates the linear acceleration, r/t^2, to charge and distance. Gauss's electric field, $\mathbf{E}$, relates angular (orbital) acceleration, $2\pi r/t^2$ to charge and distance. Applying equation 74:

$$F_C = mr/t^2 = k_e q^2/r^2 \quad \Rightarrow \quad \exists\, F_E \in \mathbb{R}:\ F_E = 2\pi(mr/t^2) = 2\pi k_e q^2/r^2. \tag{76}$$

$$F_E = 2\pi k_e q^2/r^2 = q(2\pi(k_e q/r^2)) \quad \wedge \quad \exists\, E \in R:\ E = 2\pi k_e q/r^2 \quad \Rightarrow \quad F_E = qE. \tag{77}$$

The electric field, $E = 2\pi k_e q/r^2$, conforms to the SI units $kg \cdot m \cdot s^{-2} \cdot C^{-1} = N \cdot C^{-1}$.

$$\mathbf{E} = 2\pi k_e q/|\vec{\mathbf{r}}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -4\pi k_e q/|\vec{\mathbf{r}}|^3. \tag{78}$$

$$\nabla \cdot \mathbf{E} = -4\pi k_e q/|\vec{\mathbf{r}}|^3 \quad \wedge \quad \varepsilon_0 \stackrel{\text{def}}{=} 1/4\pi k_e \quad \wedge \quad \rho \stackrel{\text{def}}{=} q/|\vec{\mathbf{r}}|^3 \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -\rho/\varepsilon_0, \tag{79}$$

which is Gauss's electric field law [3] in differential form.

B, μ_0, and Lorentz's law

Applying the distance contraction equation, 59, to equation 77, where r' is the distant observer frame of reference and r is local (rest) frame of reference:

$$r' = r/\sqrt{1 - v^2/c^2} \quad \wedge \quad F' = 2\pi k_e q^2/r'^2 \quad \Rightarrow \quad F = 2\pi k_e q^2(1 - v^2/c^2)/r^2. \tag{80}$$

From equation 77:

$$E = 2\pi k_e q/r^2 \ \wedge \ F = 2\pi k_e q^2(1 - v^2/c^2)/r^2 \ \Rightarrow \ F = q(E - (((2\pi k_e/c^2)q/r^2))v^2). \tag{81}$$

$$F = q(E - (((2\pi k_e/c^2)q/r^2))v^2) \ \wedge \ \exists\, B: \ B = (2\pi k_e/c^2)vq/r^2 \ \Rightarrow \ F = q(E - Bv). \tag{82}$$

Translating equation 82 to vector form:

$$F = q(E - Bv) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{\mathbf{v}}). \tag{83}$$

$$\mathbf{B} \times \vec{\mathbf{v}} = -(\vec{\mathbf{v}} \times \mathbf{B})) \quad \wedge \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{\mathbf{v}}) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} + \vec{\mathbf{v}} \times \mathbf{B}), \tag{84}$$

which is Lorentz law in the distant observer frame of reference, where the magnetic field, $B = (2\pi k_e/c^2)vq/r^2$, conforms to the base SI units: $kg \cdot s^{-1} \cdot C^{-1} \stackrel{\text{def}}{=} kg \cdot s^{-2} \cdot A^{-1} \stackrel{\text{def}}{=} T$.

Based on the assumption (having no math proof-motivation) of a magnetic dipole, the magnetic field divergence outward, $\nabla \cdot \mathbf{B}$, at a point, P, normal to a surface enclosing a charge equals the field divergence inward, $\nabla \cdot \mathbf{B}$:

$$\mathbf{B} = (2\pi k_e/c^2)vq/|\vec{\mathbf{r}}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{B} = -(4\pi k_e/c^2)q/|\vec{\mathbf{r}}|^3 + (4\pi k_e/c^2)q/|\vec{\mathbf{r}}|^3 = 0. \tag{85}$$

$$\nabla \cdot \mathbf{B} = -(4\pi k_e/c^2)q/|\vec{\mathbf{r}}|^3 + (4\pi k_e/c^2)q/|\vec{\mathbf{r}}|^3 = 0 \quad \Rightarrow \quad \mu_0 = (4\pi k_e/c^2), \tag{86}$$

where $\mu_0 = 4\pi k_e/c^2$ conforms to the SI units $kg \cdot m \cdot C^{-2} = kg \cdot m \cdot s^{-2}A^{-2}$.

Note: In the last section of this article, it will be shown that the CODATA definition, $\mu_0 = 2h\alpha/cq_e^2$ [17], and $\mu_0 = 4\pi k_e/c^2$ both reduce to the same equation, $\mu_0 = 4\pi k_m/q_p^2$.

Faraday's law

From the magnetic field equation 83, where the electric and magnetic fields are propagating at the speed, $v = c$:

$$B = (2\pi k_e/c^2)qv/r^2 \quad \wedge \quad v = c \quad \wedge \quad r = ct \quad \Rightarrow \quad B = (2\pi k_e/c^3)q/t^2. \tag{87}$$

$$B = (2\pi k_e/c^3)q/t^2 \quad \Rightarrow \quad \partial B/\partial t = -(4\pi k_e/c^3)q/t^3. \tag{88}$$

$$\partial B/\partial t = -(4\pi k_e/c^3)q/t^3 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial B/\partial t = -4\pi k_e q/r^3. \tag{89}$$

Faraday's law is the case, where the electric field, $\mathbf{E}$, is a single radial vector, $\mathbf{E}_{\tilde{\mathbf{r}}_1}$ acting on some point, P. Therefore, at point, P, $\mathbf{E}_{\tilde{\mathbf{r}}_2} = 0$ and $\mathbf{E}_{\tilde{\mathbf{r}}_3} = 0$. From the Lorentz equation 84, the magnetic field, $\mathbf{B}$ at point P, is perpendicular to $\mathbf{E}_{\tilde{\mathbf{r}}_1}$ (aligned on $\tilde{\mathbf{r}}_2$). For a constant force, the Lorentz equation makes the electric field, $\mathbf{E}_{\tilde{\mathbf{r}}_1}$, a function of $\mathbf{B}_{\tilde{\mathbf{r}}_2}$. Combining with equation 78:

$$\mathbf{E}_{\tilde{\mathbf{r}}_1} = 2\pi k_e q/|\tilde{\mathbf{r}}_2|^2 = f(\mathbf{B}_{\tilde{\mathbf{r}}_2}) \quad \wedge \quad \mathbf{E}_{\tilde{\mathbf{r}}_2} = 0 \quad \wedge \quad \mathbf{E}_{\tilde{\mathbf{r}}_3} = 0 \quad \Rightarrow \quad \nabla \times \mathbf{E} = \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ \frac{\partial}{\partial \tilde{\mathbf{r}}_1^2} & \frac{\partial}{\partial \tilde{\mathbf{r}}_2^2} & \frac{\partial}{\partial \tilde{\mathbf{r}}_3^2} \\ \mathbf{E}_{\tilde{\mathbf{r}}_1} & 0 & 0 \end{vmatrix}$$

$$= (\frac{\partial \mathbf{E}_{r_3}}{\partial \tilde{\mathbf{r}}_2^2} - \frac{\partial \mathbf{E}_{r_2}}{\partial \tilde{\mathbf{r}}_3^2})\hat{\mathbf{i}} + (\frac{\partial \mathbf{E}_{r_1}}{\partial \tilde{\mathbf{r}}_3^2} - \frac{\partial \mathbf{E}_{r_3}}{\partial \tilde{\mathbf{r}}_1^2})\hat{\mathbf{j}} + (\frac{\partial \mathbf{E}_{r_2}}{\partial \tilde{\mathbf{r}}_1^2} - \frac{\partial \mathbf{E}_{r_1}}{\partial \tilde{\mathbf{r}}_2^2})\hat{\mathbf{k}}$$

$$= 0\hat{\mathbf{i}} + 0\hat{\mathbf{j}} + (0 - \frac{\partial 2\pi k_e q/|\tilde{\mathbf{r}}_2|^2}{\partial |\tilde{\mathbf{r}}_2|^2})\hat{\mathbf{k}} \quad = \quad 4\pi k_e q/|\vec{\mathbf{r}}_2|^3\hat{\mathbf{k}} \quad \equiv \quad \nabla \times \mathbf{E} = 4\pi k_e q/|\vec{\mathbf{r}}|^3. \tag{90}$$

Combining equations 90 and 89 yields Maxwell-Faraday's law [3]:

$$\nabla \times \mathbf{E} = 4\pi k_e q/|\vec{\mathbf{r}}|^3 \quad \wedge \quad \partial \mathbf{B}/\partial t = -4\pi k_e q/|\vec{\mathbf{r}}|^3 \quad \Rightarrow \quad \nabla \times \mathbf{E} = -\partial \mathbf{B}/\partial t. \tag{91}$$

Maxwell's wave equations

From the Gauss's electric field equation 79:

$$\mathbf{E} = 2\pi k_e q/|\vec{\mathbf{r}}|^2 \quad \wedge \quad r = |\vec{\mathbf{r}}| \quad \Rightarrow \quad \nabla^2 \mathbf{E} = 12\pi k_e q/|\vec{\mathbf{r}}|^4. \tag{92}$$

$$\mathbf{E} = 2\pi k_e q/|\vec{\mathbf{r}}|^2 \ \wedge \ r = ct \quad \Rightarrow \quad \mathbf{E} = 2\pi k_e q/(ct)^2 \quad \Rightarrow \quad \partial^2 \mathbf{E}/\partial t^2 = (12\pi k_e/c^2)q/t^4. \tag{93}$$

$$\partial^2 \mathbf{E}/\partial t^2 = (12\pi k_e/c^2)q/t^4 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial^2 \mathbf{E}/\partial t^2 = 12\pi k_e c^2 q/|\vec{\mathbf{r}}|^4. \tag{94}$$

$$\nabla^2 \mathbf{E} = 12\pi k_e q/|\vec{\mathbf{r}}|^4 \ \wedge \ \partial^2 \mathbf{E}/\partial t^2 = 12\pi k_e c^2 q/|\vec{\mathbf{r}}|^4 \quad \Rightarrow \quad \nabla^2 \mathbf{E} = (1/c^2)\partial^2 \mathbf{E}/\partial t^2. \tag{95}$$

$$\nabla^2 \mathbf{E} = (1/c^2)\partial^2 \mathbf{E}/\partial t^2 \quad \wedge \quad \varepsilon_0 \mu_0 = 1/c^2 \quad \Rightarrow \quad \nabla^2 \mathbf{E} = \varepsilon_0 \mu_0 \partial^2 \mathbf{E}/\partial t^2, \tag{96}$$

which is Maxwell's electric field wave equation. And from equation (89):

$$\partial B/\partial t = -4\pi k_e q/r^3 \quad \wedge \quad r = |\vec{\mathbf{r}}| \quad \Rightarrow \quad \nabla^2 \mathbf{B} = 12\pi k_e q/|\vec{\mathbf{r}}|^4. \tag{97}$$

$$\mathbf{B} = 2\pi k_e q/|\vec{\mathbf{r}}|^2 \ \wedge \ r = ct \ \Rightarrow \quad \mathbf{B} = 2\pi k_e q/(ct)^2 \ \Rightarrow \quad \partial^2 \mathbf{B}/\partial t^2 = (12\pi k_e/c^2)q/t^4. \tag{98}$$

$$\partial^2 \mathbf{B}/\partial t^2 = (12\pi k_e/c^2)q/t^4 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial^2 \mathbf{B}/\partial t^2 = 12\pi k_e c^2 q/|\vec{\mathbf{r}}|^4. \tag{99}$$

$$\nabla^2 \mathbf{B} = -12\pi k_e q/|\vec{\mathbf{r}}|^4 \ \wedge \ \partial^2 \mathbf{B}/\partial t^2 = -12\pi k_e c^2 q/|\vec{\mathbf{r}}|^4 \quad \Rightarrow \quad \nabla^2 \mathbf{B} = (1/c^2)\partial^2 \mathbf{E}/\partial t^2. \tag{100}$$

$$\nabla^2 \mathbf{B} = (1/c^2)\partial^2 \mathbf{E}/\partial t^2 \quad \wedge \quad \varepsilon_0 \mu_0 = 1/c^2 \quad \Rightarrow \quad \nabla^2 \mathbf{B} = \varepsilon_0 \mu_0 \partial^2 \mathbf{B}/\partial t^2, \tag{101}$$

which is Maxwell's magnetic field wave equation.

The fine structure constant, α

The ratio of two subtypes of force implies ratios of the form:

$$\alpha_\tau = \frac{F_{\tau_1}}{F_{\tau_2}} = \frac{K\tau_1^2/r^2}{K\tau_2^2/r^2} = \frac{\tau_1^2}{\tau_2^2}. \tag{102}$$

For example, where q_e is the elementary (electron) charge ($1.60217663 \cdot 10^{-19}\ C$), and q_p is reduced Planck charge unit, and the fine structure electron coupling constant:

$$\alpha = q_e^2/q_p^2. \tag{103}$$

The gravitational fine structure constant, α_G, is the ratio of two subtypes of Newton's gravity force, which simplifies to:

$$\alpha_G = m_e^2/m_p^2. \tag{104}$$

$\hbar$, h, and Einstein-Planck relation

[3, 18] Applying both the direct proportion ratio (115), and inverse proportion ratio (50):

$$m = k_m/r \quad \wedge \quad r = ct \quad \Rightarrow \quad m(ct)^2 = (k_m/r)r^2 = k_m r. \tag{105}$$

$$m(ct)^2 = k_m r \quad \Rightarrow \quad E \stackrel{\text{def}}{=} mc^2 = k_m r/t^2. \tag{106}$$

$$E = mc^2 = k_m r/t^2 \ \wedge \ r/t = c \ \Rightarrow \ E = mc^2 = k_m c/t. \tag{107}$$

The next 4 steps show that $k_m c = \hbar$, the *reduced* Planck constant. From the fine structure equation 104:

$$\alpha = q_e^2/q_p^2 \quad \wedge \quad q_e = 1.60217663 \cdot 10^{-19}\ C \quad \wedge \quad \alpha \approx 7.2973525705 \cdot 10^{-3}$$
$$\Rightarrow \quad q_p = \sqrt{q_e^2/\alpha} \approx 1.87554604 \cdot 10^{-18}\ C. \tag{108}$$

From equations 86 and 108:

$$\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m = 4\pi(r_p/q_p)^2/(r_p/m_p) = 4\pi m_p r_p/q_p^2 = 4\pi k_m/q_p^2. \tag{109}$$

$$\mu_0 = 4\pi k_m/q_p^2 \ \wedge \ q_p \approx 1.87554604 \cdot 10^{-18}\ C \ \wedge \ \mu_0 \approx 1.25663706127(20) \cdot 10^{-6}\ kg\ m\ C^{-2}$$
$$\Rightarrow \quad k_m = \mu_0 q_p^2/4\pi \approx 3.51767294 \cdot 10^{-43}\ kg\ m. \tag{110}$$

$$k_m \approx 3.51767294 \cdot 10^{-43}\ kg\ m \quad \wedge \quad c = 2.99792458 \cdot 10^8\ m\ s^{-1}$$
$$\Rightarrow \quad k_m c \approx 1.054571817 \cdot 10^{-34}\ kg\ m^2\ s^{-1} = \hbar. \tag{111}$$

Combining equations 107 and 111:

$$E = mc^2 = k_m c/t\ kg\ m^2\ s^{-2}\ \wedge\ k_m c = \hbar\ kg\ m^2\ s^{-1}\ \Rightarrow\ E = \hbar/t\ kg\ m^2\ s^{-2}. \tag{112}$$

$$E = \hbar/t\ kg\ m^2\ s^{-2} \quad \wedge \quad \omega = (2\pi/t)\ radian\ s^{-1} \quad \Rightarrow$$

$$E_\omega = \hbar(2\pi/t) = \hbar\omega\ kg\ m^2\ radian\ s^{-2}. \tag{113}$$

$$E_\omega = \hbar/t\ kg\ m^2\ s^{-2}\ \wedge\ f = (1/t)\ cycle\ s^{-1} = ((2\pi/t)\ radian\ s^{-1})/(2\pi\ radian\ cycle^{-1})$$

$$\Rightarrow \quad E_f = \hbar\omega(2\pi/2\pi) = 2\pi\hbar f = hf\ kg\ m^2\ cycle\ s^{-2} = hf\ kg\ m^2\ s^{-1}\ Hz. \tag{114}$$

Unit ratio and reduced Planck unit values

The direct proportion ratios are:

$$c_t = c \approx 2.99792458 \cdot 10^8\ m\ s^{-1}. \tag{115}$$

$$G = c_m c_t^2 \quad \wedge \quad G \approx 6.67430(15) \cdot 10^{-11}\ m^3\ kg^{-1}\ s^{-2} \quad \Rightarrow$$

$$c_m = G/c_t^2 \approx 7.426162 \cdot 10^{-28}\ m\ kg^{-1}. \tag{116}$$

$$k_e = c_q^2 c_t^2/c_m \quad \wedge \quad k_e \approx 8.9875517923(13) \cdot 10^9 Nm^2\ C^{-2} \quad \Rightarrow$$

$$c_q = \sqrt{k_e c_m/c_t^2} \approx 8.6175182 \cdot 10^{-18} m\ C^{-1}. \tag{117}$$

From equation 111, the inverse proportion ratio values are:

$$\hbar \approx 1.054571783 \cdot 10^{-34}\ kg\ m^2/s \quad \wedge \quad c_t = 2.99792458 \cdot 10^8\ m/s$$

$$\Rightarrow \quad k_m = \hbar/c_t \approx 3.51767283 \cdot 10^{-43}\ kg\ m. \tag{118}$$

$$r_p = \sqrt{r_p^2} = \sqrt{k_m c_m} \approx 1.616255 \cdot 10^{-35}\ m. \tag{119}$$

$$k_t = r_p/c_t \approx 8.71363 \cdot 10^{-79}\ s\ m. \tag{120}$$

$$k_q = r_p/c_q \approx 3.031361 \cdot 10^{-53}\ C\ m. \tag{121}$$

From the fine structure equation 108, the reduced Planck unit values are:

$$q_p = \sqrt{q_e^2/\alpha} \approx 1.87554604 \cdot 10^{-18}\ C. \tag{122}$$

$$r_p = q_p c_q \approx 1.616255 \cdot 10^{-35}\ m. \tag{123}$$

$$t_p = r_p/c_t \approx 5.391246 \cdot 10^{-44}\ s. \tag{124}$$

$$m_p = r_p/c_m \approx 2.176434 \cdot 10^{-8}\ kg. \tag{125}$$

Compton wavelength, λ

[3, 18] From equations 52 and 108:

$$\lambda = 2\pi r = 2\pi k_m/m = 2\pi k_m c/mc \quad \wedge \quad h = 2\pi\hbar = 2\pi k_m c \quad \Rightarrow \quad \lambda = h/mc. \tag{126}$$

Schrödenger's wave equation

Start with the previously derived "base" Einstein-Planck relation 112 and multiply the kinetic energy component by mc/mc:

$$mc^2 = \hbar/t \quad \Rightarrow \quad \exists\, V(r,t) : \ \hbar/t = \hbar/2t + V(r,t). \tag{127}$$

$$\hbar/t = h/2t + V(r,t) \quad \Rightarrow \quad \hbar/t = \hbar mc/2mct + V(r,t). \tag{128}$$

And from the distance-to-time (speed of light) ratio (115):

$$\hbar/t = \hbar mc/2mct + V(r,t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar mc^2/2mcr + V(r,t). \tag{129}$$

$$\hbar/t = \hbar mc^2/2mcr + V(r,t) \ \wedge \ \hbar/t = mc^2 \quad \Rightarrow \quad \hbar/t = \hbar^2/2mcrt + V(r,t). \tag{130}$$

$$\hbar/t = \hbar^2/2mcrt + V(r,t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar^2/2mr^2 + V(r,t). \tag{131}$$

Multiply both sides of equation 131 by a probability density function, $\Psi(r,t)$.

$$\hbar/t = \hbar^2/2mr^2 + V(r,t) \quad \Rightarrow \quad (\hbar/t)\Psi(r,t) = (\hbar^2/2mr^2)\Psi(r,t) + V(r,t)\Psi(r,t). \tag{132}$$

$$\begin{gathered}(\hbar/t)\Psi(r,t) = ((\hbar)^2/2mr^2)\Psi(r,t) + V(r,t)\Psi(r,t) \quad \wedge \\ \forall\ \Psi(r,t) : \ \partial^2\Psi(r,t)/\partial r^2 = (-1/r^2)\Psi(r,t) \quad \wedge \quad \partial\Psi(r,t)/\partial t = (i\ 1/t)\Psi(r,t) \\ \Rightarrow \quad i\hbar\partial\Psi(r,t)/\partial t = -(\hbar^2/2m)\partial^2\Psi(r,t)/\partial r^2 + V(r,t)\Psi(r,t),\end{gathered} \tag{133}$$

which is the one-dimensional position-space Schrödenger's equation [6, 18].

$$\begin{gathered}i\hbar\partial\Psi(r,t)/\partial t = -(\hbar^2/2m)\partial^2\Psi(r,t)/\partial r^2 + V(r,t)\Psi(r,t) \quad \wedge \quad ||\vec{\mathbf{r}}|| = r \\ \Rightarrow \quad \exists\, \vec{\mathbf{r}} : \ i\hbar\partial\Psi(\vec{\mathbf{r}},t)/\partial t = -(\hbar^2/2m)\partial^2\Psi(\vec{\mathbf{r}},t)/\partial\vec{\mathbf{r}}^2 + V(\vec{\mathbf{r}},t)\Psi(\vec{\mathbf{r}},t),\end{gathered} \tag{134}$$

which is the 3-dimensional position-space Schrödenger's equation [6, 18].

Dirac's wave equation

Start with the previously derived "base" Einstein-Planck relation 112:

$$mc^2 = \hbar/t \quad \Rightarrow \quad \exists\, V(r,t):\ mc^2/2 + V(r,t) = \hbar/t \quad \Rightarrow \quad 2\hbar/t - 2V(r,t) = mc^2. \tag{135}$$

$$\forall\, V(r,t):\ V(r,t) = \hbar/t \quad \wedge \quad r = ct \quad \wedge \quad 2\hbar/t - 2V(r,t) = mc^2$$
$$\Rightarrow \quad 2\hbar/t - i2\hbar c/r = mc^2. \tag{136}$$

Use the ratios, $r = c_q q$, and, $r = ct$. to multiply each term on the left side of equation by 1:

$$qc_q/r = qc_q/ct = 1 \wedge 2\hbar/t - i2\hbar c/r = mc^2 \ \Rightarrow\ 2\hbar(qc_q/c)/t^2 - i\hbar((qc_q/c)/r^2)c = mc^2. \tag{137}$$

Applying a quantum amplitude equation in complex form to equation 137:

$$A_0 = 2(c_q/c)((1/t) - i(1/r)) \quad \wedge \quad 2\hbar(qc_q/c)/t^2 - i\hbar((qc_q/c)/r^2)c = mc^2$$
$$\Rightarrow\ \hbar\partial(-qA_0)/\partial t - i\hbar(\partial(-qA_0)/\partial r)c = mc^2. \tag{138}$$

Translating equation 139 to moving (rest frame) coordinates via the Lorentz factor, $\gamma_0 = 1/\sqrt{1-(v/c)^2}$:

$$\hbar\partial(-qA_0)/\partial t - i\hbar(\partial(-qA_0)/\partial r)c = mc^2$$
$$\Rightarrow \quad \gamma_0\hbar\partial(-qA_0)/\partial t - \gamma_0 i\hbar(\partial(-qA_0)/\partial r)c = mc^2. \tag{139}$$

Translating equation 139 to vector form:

$$\gamma_0\hbar(\partial(-qA_0)/\partial t) - \gamma_0 i\hbar(\partial(-qA_0)/\partial r)c = mc^2 \quad \wedge$$
$$||\vec{\mathbf{r}}|| = r \quad \wedge \quad ||\vec{\mathbf{A}}|| = A_0 \quad \wedge \quad ||\vec{\gamma}|| = \gamma_0$$
$$\Leftrightarrow \quad \exists\, \vec{\mathbf{r}}, \vec{\mathbf{A}}, \vec{\gamma} : \gamma_0\hbar(\partial(-qA_0)/\partial t) - \vec{\gamma}\cdot i\hbar(\partial(-q\vec{\mathbf{A}})/\partial r)c = mc^2. \tag{140}$$

Multiplying both sides of equation 140 by the spinor, Ψ, yields:Dirac's equation [8]:

$$\gamma_0\hbar(\partial(-qA_0)/\partial t) - \vec{\gamma}\cdot i\hbar(\partial(-q\vec{\mathbf{A}})/\partial r)c = mc^2 \quad \Rightarrow$$
$$\gamma_0\hbar(\partial(-qA_0)/\partial t)\Psi - \vec{\gamma}\cdot i\hbar(\partial(-q\vec{\mathbf{A}})/\partial r)c\Psi = mc^2\Psi \tag{141}$$

Total of a type

Applying both the direct (115) and inverse proportion ratios (118) to the Euclidean distance:

$$r = \sqrt{r_1^2 + r_2^2}, \quad r_1 = c_\tau \tau, \quad r_2 = k_\tau/\tau, \quad \tau \in \{t, m, q\}$$
$$\Rightarrow \quad r = \sqrt{(c_\tau \tau)^2 + (k_\tau/\tau)^2} \quad \Leftrightarrow \quad \tau = \sqrt{(r/c_\tau)^2 + (k_\tau/r)^2}. \quad (142)$$

Quantum extension to general relativity

The simplest way to demonstrate how to add quantum physics to general relativity is by extending Schwarzschild's gravitational time dilation equation and black hole metric. Apply the total of a type equation 142 to the derivation of Schwarzschild's time dilation and metric (63):

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - (v^2/c^2)(r/r)} \quad \wedge \quad r = \sqrt{(c_m m)^2 + (k_m/m)^2} = Q_m$$
$$\Rightarrow \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - Q_m v^2/rc^2}. \quad (143)$$
$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - Q_m v^2/rc^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2$$
$$\Rightarrow \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m v_{escape}^2/rc^2}. \quad (144)$$

For a photon, the escape velocity, $v_{escape} = c$.

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m v_{escape}^2/rc^2}. \ \wedge \ v_{escape} = c$$
$$\Rightarrow \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m c^2/rc^2} = \sqrt{1 - 2Q_m/r}. \quad (145)$$

Combining equation 145 with equation 60 yields Schwarzschild's gravitational time dilation with a quantum mass effect:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m/r} \quad \wedge \quad t' = t\sqrt{1 - (v^2/c^2)} \quad \Rightarrow \quad t' = t\sqrt{1 - 2Q_m/r}. \quad (146)$$

Schwarzschild defined the black hole event horizon radius, $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$. The radius with the quantum extension is $\alpha \stackrel{\text{def}}{=} 2Q_m$. At this point the exact same equations 68 through 71 yield what looks like the same Schwarzschild black hole metric.

Quantum extension to Newton's gravity force

The quantum mass effect is easier to understand in the context Newton's gravity equation than in general relativity, because the metric equations and solutions in the EFEs are much more complex. From equations 142 and 48:

$$m/\sqrt{(r/c_m)^2 + (k_m/r)^2} = 1 \quad \wedge \quad r^2/(ct)^2 = 1 \quad \Rightarrow \quad r^2/(ct)^2 = m/\sqrt{(r/c_m)^2 + (k_m/r)^2}$$
$$\Rightarrow \quad r^2/t^2 = c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2}. \tag{147}$$
$$r^2/t^2 = c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2} \quad \Rightarrow \quad (m/r)(r^2/t^2) = (m/r)(c^2 m/\sqrt{(r/c_m)^2 + (k_m/r)^2})$$
$$\Rightarrow \quad F \stackrel{\text{def}}{=} mr/t^2 = c^2 m^2/\sqrt{(r^4/c_m^2) + k_m^2}. \tag{148}$$
$$F = c^2 m^2/\sqrt{(r^4/c_m^2) + k_m^2} \quad \wedge \quad \forall\, m \in \mathbb{R},\, \exists\, m_1, m_2 \in \mathbb{R} :\ m_1 m_2 = m^2$$
$$\Rightarrow \quad F = c^2 m_1 m_2/\sqrt{(r^4/c_m^2) + k_m^2}. \tag{149}$$

Quantum extension to Coulomb's charge force

$$q^2/((r/c_q)^2 + (k_q/r)^2) = 1 \quad \wedge \quad r^2/(ct)^2 = 1 \quad \Rightarrow \quad r^2/(ct)^2 = q^2/((r/c_q)^2 + (k_q/r)^2)$$
$$\Rightarrow \quad r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2). \tag{150}$$
$$(1/r)(r^2/t^2) = (1/r)(c^2 q^2/((r/c_q)^2 + (k_q/r)^2)) \quad \Rightarrow \quad r/t^2 = c^2 q^2/(r^3/c_q^2 + k_q^2/r). \tag{151}$$
$$\forall\, q \in \mathbb{R} :\ \exists\, q_1, q_2 \in \mathbb{R} :\ q_1 q_2 = q^2 \quad \wedge \quad r/t^2 = c^2 q^2/(r^3/c_q^2 + k_q^2/r) \quad \Rightarrow \ \textit{nonumber} \tag{152}$$
$$\exists\, q_1, q_2 \in \mathbb{R} :\ r/t^2 = c^2 q_1 q_2/(r^3/c_q^2 + k_q^2/r). \tag{153}$$
$$r/t^2 = c^2 q_1 q_2/(r^3/c_q^2 + k_q^2/r) \quad \wedge \quad m = r/c_m \quad \Rightarrow \quad F \stackrel{\text{def}}{=} mr/t^2 = (c^2/c_m) q_1 q_2/(r^2/c_q^2 + k_q^2/r^2). \tag{154}$$

INSIGHTS AND IMPLICATIONS

1. The notion of Euclidean distance as the inverse function of a volume, where that volume is the sum of subset Euclidean volumes(.4), suggests defining a distance measure as the inverse function of a volume that is the sum of subset Euclidean and non-Euclidean volumes,

$$\forall\, n \in \mathbb{N}, \quad f_n, g \in C^2, \quad d = f_n^{-1} v = f_n^{-1}(\textstyle\sum_{i=1}^m f_n(v_i)) = f_n^{-1}(\sum_{i=1}^m g(s_{i,1}, \cdots, s_{i,n})) : \tag{155}$$

(a) shows the intimate relation between distance and volume that definitions, like inner product space and metric space, ignore [9, 10];

(b) is a more simple and concise definition of a distance measure, having the metric space properties [9, 10].

2. The left side of the distance sum inequality (.7),

$$(\sum_{i=1}^{m}(a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^{m} a_i^n)^{1/n} + (\sum_{i=1}^{m} b_i^n)^{1/n}, \tag{156}$$

differs from the left side of Minkowski's sum inequality [14]:

$$(\sum_{i=1}^{m}(a_i^n + b_i^n)^{\mathbf{n}})^{1/n} \leq (\sum_{i=1}^{m} a_i^n)^{1/n} + (\sum_{i=1}^{m} b_i^n)^{1/n}. \tag{157}$$

(a) The two inequalities are only the same where $n = 1$.

(b) The distance sum inequality (.7) is a more fundamental inequality because the proof does not require the convexity and Hölder's inequality assumptions required to prove the Minkowski sum inequality [14].

(c) $\forall\ n > 1,\ v > 0$: The distance sum inequality term, $v_i^n = a_i^n + b_i^n$: $d = v^{1/n} = (\sum_{i=1}^{m} v_i^n)^{1/n}$, is the Minkowski distance, which makes it directly related to geometry. But the Minkowski sum inequality term, $d = v^{1/n} = (\sum_{i=1}^{m}((v_i^n)^{\mathbf{n}}))^{1/n} = (\sum_{i=1}^{m} v_i^{\mathbf{n^2}})^{1/n}$, is *not* a Minkowski distance.

(d) The distance sum inequality might be applicable to machine learning.

3. *Combinatorics.* The number of n-tuples, $v_c = \prod_{i=1}^{n} |x_i|$, was proven to imply the Euclidean volume equation (.1). And the notion of distance was derived as the inverse function of the sum of the number of n-tuples (.4). These equations were derived without relying on the geometric primitives and relations in Euclidean geometry [19, 20], axiomatic geometry [21–25], trigonometry [26, 27] calculus [26, 28, 29], and vector analysis [5].

4. *Combinatorics.* The commutative property of the operations defining volume and distance equations limits a domain set to $n \leq 3$ elements (.16). Other elements must have different types (are elements of different sets).

(a) If there are more than 6 dimensions, $\{r_1, r_2, r_3, t, m, q\}$, then the dimensions are not unordered dimensions $\subseteq \mathbb{R}$. For example, there can be more dimensions

$\subseteq \mathbb{R}$, where each dimension has an intrinsic property giving it a position in strict totally ordered set.

(b) If each type of quantum state is an ISCL, then there are at most 3 states of the same type: 3 orientations per dimension of space (+1,0-1), 3 quark color charges, {red, green, blue}, 3 quark anti-color charges, and so on.

5. A discrete (point) valued state has measure 0 (zero-length interval). The ratio of a time or distance interval length to zero is undefined, which is the reason quantum entangled (discrete) state values exist independent of (non-deterministic with respect to) distance (position), time, mass, and charge. Therefore, a state value in space and time is undetermined until observed, like Schrödenger's poisoned cat being both alive and dead until the box is opened [6].

6. For each unit interval length, r_p, of a 3-dimensional distance, $r \subseteq \mathbb{R}$, there are unit interval lengths of other types of dimensions, forming unit ratios (115): $c_t = r_p/t_p$, $c_m = r_p/m_p$, $c_q = r_p/q_p \Leftrightarrow$ the inverse proportion ratios (50): $k_t = r_p t_p$, $k_m = r_p m_p$, $k_t = r_p q_p$, where r_p, t_p, m_p, and q_p are the reduced Planck units (122).

7. An empirical law *describes* the relations between the variables in an equation. Deriving empirical laws from the ratios *explains* the reasons for relations between the variables in an equation. Further, all the derivations of the physics equations from the ratios are much shorter and simpler than other derivations, which shows that the ratios are an important tool for physicists and engineers.

8. As shown in the subsection deriving the Schwarzschild's gravitational time dilation and black hole metric (63) [15, 16] using ratios illustrates a simple way of finding solutions to Einstein's field equations.

9. c_t, c_m, and c_q are the *maximum* ratios: $r = \sqrt{r_1^2 + r_2^2 + r_3^2} \Rightarrow r_1,\ r_2,\ r_3 \leq r \Rightarrow \forall\, \tau \in \{t, m, q\} : r_1/\tau,\ r_2/\tau,\ r_3/\tau \leq r/\tau = r_p/\tau_p = c_\tau$. For example, the speeds of gravity and light waves are limited to the speed, c_t.

10. c_t is a component of the constants: $G = c_m c_t^2$, $k_e = c_q^2 c_t^2/c_m$, $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2/c_m)$, and $\hbar = k_m c_t$. The only constant, derived in this article, that does not contain c_t is vacuum permeability: $\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m$.

11. The fine structure electron coupling constant, α was derived from the ratio of two subtypes of charge force that reduces to the unit-less ratio $\alpha = q_e^2/q_p^2 \approx 0.0072973526$ (104), which is the empirical CODATA value [17].

    (a) The CODATA electron coupling version of the fine structure constant, α is defined as: $\alpha = q_e^2/4\pi\varepsilon_0\hbar c) = q_e^2/2\varepsilon_0 hc$ [17]. The following steps show that the CODATA definition reduces to the ratio-derived equation:

$$\varepsilon_0 = 1/4\pi k_e = 1/(4\pi(c_q^2 c_t^2/c_m) \quad \wedge \quad \hbar = k_m c_t \quad \wedge \quad h = 2\pi\hbar$$
$$\Rightarrow \quad \varepsilon_0 hc = 2\pi k_m c_t^2/(4\pi(c_q^2/c_m)c_t^2) = k_m/(2(c_q^2/c_m))$$
$$= m_p r_p/(2((r_p/q_p)^2/(r_p/m_p))) = q_p^2/2. \quad (158)$$

$$\alpha = q_e^2/2\varepsilon_0 hc \quad \wedge \quad \varepsilon_0 hc = q_p^2/2 \quad \Rightarrow \quad \alpha = q_e^2/2(q_p^2/2) = q_e^2/q_p^2. \quad (159)$$

    (b) The fine structure electron coupling constant is the ratio of electron static charge force to the propagating quantum charge (photon/electromagnetic) wave force, caused by a moving charged particle.

    (c) The gravitational fine structure electron coupling constant, α_G, can be expressed as the ratio of two subtypes of Newton's gravity force, where m_e is the rest electron mass and m_p is the reduced Planck mass unit: $\alpha_G = m_e^2/m_p^2$ (104).

    The currently used gravitational fine structure constant reduces to the equation derived in this article: $\alpha_G = Gm_e^2/\hbar c = (c_m c_t^2)m_e^2/(k_m c_t)c_t = c_m m_e^2/k_m = (r_p/m_p)m_e^2/m_p r_p = m_e^2/m_p^2$.

12. The assumption that the old definition, $\mu_0 = 4\pi k_e/c_t^2 = 1/\varepsilon c^2$, and the newer NIST/CODATA definition, $\mu_0 = 2h\alpha/cq_e^2$[17], are different is *incorrect*! The following steps will show that both equations reduce to the same base equation, $\mu_0 = 4\pi k_m/q_p^2$, which shows that the old and newer NIST/CODATA definitions differ in empirical value because of different measurement methods.

    Using the ratios to reduce the old definition of μ_0 derived in this article:

$$\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m = 4\pi(r_p/q_p)^2/(r_p/m_p) = 4\pi m_p r_p/q_p^2 = 4\pi k_m/q_p^2. \quad (160)$$

    Using the ratios to reduce the CODATA definition:

$$h = 2\pi k_m c_t, \;\; \alpha = q_e^q/q_p^2 \;\; \Rightarrow \;\; \mu_0 = 2h\alpha/cq_e^2 = 4\pi k_m c_t(q_e^2/q_p^2)/c_t q_e^2 = 4\pi k_m/q_p^2. \quad (161)$$

13. Empirical and hypothesized laws of physics use an *opaque* constant to make the units in an equation balance. The opacity has led to the *incorrect* assumptions of those constants being fundamental (atomic) constants.

    In this article, some opaque constants are derived directly from (composed of) the ratios: gravity, $G = c_m c_t^2$ (55), charge, $k_e = c_q^2 c_t^2 / c_m$ (74), and reduced Planck $\hbar = k_m c_t$ (108). $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2/c_m)$ (79) and $\mu_0 = 4\pi k_e / c_t^2 = 4\pi c_q^2 / c_m$ (85).

    And the quantum extensions to: Schwarzschild's gravitational time dilation (145), Newton's gravity force (149), and Coulomb's charge force (154), show that, where the quantum effects become measurable, the constants G, k_e, ε_0, and μ_0 no longer exist (are no longer valid).

    Therefore, G, k_e, ε_0, μ_0, and $\hbar$ are *not* fundamental constants.

14. Constants that use notions of temperature and pressure for example, the Boltzmann constant [17], might not possible to define solely in terms of unit ratios.

15. The derivations of: $\nabla \cdot \mathbf{g} = -4\pi G\rho$ from $\mathbf{g} = 2\pi G m / |\tilde{\mathbf{r}}|^2$ (56), and $\nabla \cdot \mathbf{E} = -\rho/\varepsilon_0$ from $\mathbf{E} = 2\pi k_e q / |\tilde{\mathbf{r}}|^2$ (79), show that the use of mass and charge density, ρ show that:

    (a) Mass and charge density as motivations are unnecessary complexities. Equation 53 used to derive Newton's law and equation 72 used to derive Coulomb's law relate linear acceleration, r/t^2 to mass and charge. It is small step to recognize Gauss's $\mathbf{g}$, Poisson's $-\nabla\Phi(\tilde{\mathbf{r}}, t)$, and Gauss's $\mathbf{E}$ relate the angular (orbital) acceleration, $2\pi r/t^2$, to mass and charge, where differentiation yields the gravity field and electric field laws.

    The derivations are much simpler and shorter using angular (orbital) acceleration, $2\pi r/t^2$. The orbit length, $2\pi r$, corresponds to the Gauss's line integral length of a curve circumscribing a sphere containing the mass or charge.

    (b) Mass and charge density obfuscates the pattern, $\nabla \cdot f(x, r) = -4\pi k_x x / |\tilde{\mathbf{r}}|^3$, being derived from the inverse square pattern, $f(x, r) = 2\pi k_x x / |\tilde{\mathbf{r}}|^2$. The constant, G (derived from the inverse square relation (53)), is used in Einstein's constant, $\kappa = 8\pi G/c^4$, and the use of energy density in the stress energy tensor, $T_{\mu,\nu}$, in Einstein's field equations [5] also obfuscate the inverse square pattern.

16. The derivation Lorentz's law here (83) differs from other relativistic derivations, in that equations for the magnetic field, $\mathbf{B}$, and magnetic permeability, μ_0, are derived in the process. The equation, $\mathbf{B} = (2\pi k_e/c^2)vq/|\vec{\mathbf{r}}|^2$, allows a simple derivation of the Faraday's law (91) and the magnetic component of Maxwell's electromagnetic equations (101).

17. Maxwell's derivation of the electric and magnetic field wave equations required: Gauss's electric field law in a vacuum: $\nabla \cdot \mathbf{E} = 0$, Gauss's magnetic field law in a vacuum: $\nabla \cdot \mathbf{B} = 0$, Faraday's Law: $\nabla \times \mathbf{E} = -\partial \mathbf{B}/\partial t$, and Ampere's Law: $\nabla \times \mathbf{B} = \mu_0 \varepsilon_0 \partial \mathbf{E}/\partial t$[3]. In contrast, the derivations (96) (101), in this article, did not require any of those laws and relied solely on equations derived from the ratios.

18. Einstein's relativity equations: 1) assume the Lorentz transformations, 2) assume the laws of physics are same at each coordinate point, 3) assume the notion of light, and 4) assume that the speed of light is the same at each coordinate point [5, 7].

    The derivations, in this article, were made without those assumptions (does not even require the notion of light). The unit ratios are ratios of Euclidean distance to time, mass, and charge. Therefore, the unit ratios and all equations (laws) and constants derived from those ratios must be the same at each Euclidean-like coordinate point.

19. Using the quantum units, r_p and t_p: r_p/t_p^2, suggests a maximum linear acceleration for masses. And $2\pi r_p/t_p^2$ suggests a maximum orbital or rotational acceleration.

20. The Einstein-Planck relations, $E_\omega = \hbar\omega\ kg\ m^2\ radian\ s^{-2}$ (113) and $E_f = hf\ kg\ m^2\ s^{-1} Hz$ (114), were both derived from the base Einstein-Planck relation, $E = \hbar/t\ kg\ m^2\ s^{-2}$ (112). And the Schrödenger's wave equation (134) [6] and Dirac's wave equation (140) [8] were both derived from the base Einstein-Planck relation.

21. The quantum extensions to: Schwarzschild's gravitational time dilation (145), black hole metric (71), Newton's gravity force (149), and Coulomb's charge force (152) make quantifiable predictions:

    (a) For gravity, $\lim_{r\to 0} F = c^2 m_1 m_2/k_m$, and for charge, $\lim_{r\to 0} F = 0$. Finite maximum gravity and charge forces: 1) eliminates the problem of forces going to

infinity as $r \to 0$, 2) might allow radioactivity without the need for a weak force, and 3) finite sloped energy well walls might explain quantum tunneling.

(b) The quantum-extended relativity and classic equations reduce to the non-extended relativity and classic equations and constants, where the distance between masses and charges is sufficiently large or the masses and charges sufficiently large that the quantum effects are not measurable. *Note* that G, k_e, ε_0, μ_0, and κ (Einstein's constant, which contains G) do not exist (are not valid), where the quantum effects becomes measurable.

(c) The covariant tensor components, in Einstein's field equations, that had the units $1/distance^2$, will now have the more complex units, $1/\sqrt{(distance^4/c_\tau^2) + k_\tau^2}$, $\tau \in \{t, m\}$.

(d) $1/\sqrt{(distance^4/c_\tau^2) + k_\tau^2}$ implies that as distance $\to 0$, spacetime curvature peaks at the Planck unit distance, r_p (122). The ratio of quantum units, m_p/r_p^3, might indicate a maximum mass density. A finite force (spacetime curvature) and finite mass density would imply that black holes have sizes > 0 (are not singularities). Black hole evaporation might be possible. If there was a "big bang," then it might not have originated from a singularity.

---

[1] Copyright © 2026 George M. Van Treeck. All rights reserved.

[2] I. Newton, *The Mathematical Principles of Natural Philosophy* (Patristic Publishing, 2019) reprint.

[3] R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics* (California Institute of Technology, 2010) https://www.feynmanlectures.caltech.edu/.

[4] A. Einstein, Die grundlage der allgemeinen relativitätstheorie (the foundation of the general theory of relativity), Annalen der Physik (1916).

[5] H. Weyl, *Space-Time-Matter* (Dover Publications Inc, 1952).

[6] E. Schrödenger, *Collected Papers On Wave Mechanics* (Minkowski Institute Press, 2020).

[7] A. Einstein, *Relativity, The Special and General Theory* (Princeton University Press, 2015).

[8] P. Dirac, *The Principles of Quantum Mechanics* (Snowball Publishing, 1957).

[9] R. R. Goldberg, *Methods of Real Analysis* (John Wiley and Sons, 1976).

[10] W. Rudin, *Principles of Mathematical Analysis* (McGraw Hill Education, 1976).

[11] W. Conradie and V. Goranko, *Logic and Discrete Mathematics* (Wiley, 2015).

[12] X. P. N. Guigui, N. Miolane, Introduction to riemannian geometry and geometric statistics: from basic theory to implementation with geomstats, Foundations and Trends in Machine Learning **16**, 182 (2023), https://inria.hal.science/hal-03766900v2, hal-00644459.

[13] Rocq, Rocq proof assistant (2025), https://rocq-prover.org/docs.

[14] H. Minkowski, *Geometrie der Zahlen* (Chelsea, 1953) reprint.

[15] K. Schwarzschild, Über das gravitationsfeld eines massenpunktes nach der einsteinschen theorie, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften **7**, 198 (1916), https://archive.org/stream/sitzungsberichte1916deutsch #page/188/mode/2up.

[16] S. Antoci and A. Loinger, An english translation of k. schwarzschild's 1916 article: On the gravitational field of a mass point according to einstein's theory, ARXIV.org (1999), https://arxiv.org/abs/physics/9905030.

[17] CODATA, Physical constant values (2022), https://physics.nist.gov/cuu/Constants/index.html.

[18] M. Jain, *Quantum Mechanics: A Textbook for Undergraduates* (PHI Learning Private Limited, New Delhi, India, 2011).

[19] D. E. Joyce, Euclid's elements (1998), http://aleph0.clarku.edu/ djoyce/java/elements/elements .html.

[20] E. S. Loomis, *The Pythagorean Proposition* (NCTM, 1968).

[21] G. D. Birkhoff, A set of postulates for plane geometry (based on scale and protractors), Annals of Mathematics **33** (1932).

[22] D. Hilbert, *The Foundations of Geometry (2cd Ed)* (Chicago: Open Court, 1980) http://www.gutenberg.org/ebooks/17384.

[23] J. M. Lee, *Axiomatic Geometry* (American Mathematical Society, 2010).

[24] A. Tarski and S. Givant, Tarski's system of geometry, The Bulletin of Symbolic Logic **5**, 175 (1999).

[25] O. Veblen, A system of axioms for geometry, Trans. Amer. Math. Soc **5**, 343 (1904), https://doi.org/10.1090/S0002-9947-1904-1500678-X.

[26] A. Bogomolny, *Pythagorean Theorem* (Interactive Mathematics Miscellany and Puzzles, 2010) http://www.cut-the-knot.org/pythagoras/ CalculusProofCorrectedVersion.shtml.

[27] J. Zimba, On the possibility of trigonometric proofs of the pythagorean theorem, Forum Geometricorum **9**, 275 (2009).

[28] B. C. Berndt, Ramanujan-100 years old (fashioned) or 100 years new (fangled)?, The Mathematical Intelligencer **10** (1988).

[29] M. Staring, The pythagorean proposition a proof by means of calculus, Mathematics Magazine **69**, 45 (1996).