

Copyright © 2026 by George M. Van Treeck. All rights reserved.

## **Mathematical physics made easy(er)**

George M. Van Treeck

(Dated: May 7, 2026)

# Abstract

Euclidean volume and some distance equations are proved to be instances of ordered combinations (n-tuples) of elements from a list of disjoint, countable sets. A permutation property of volume is proved to limit a set of unordered elements to a cyclic list of at most 3 elements: a set of 3 unordered distance dimensions (height, width, and depth) and a set of 3 unordered non-distance dimensions (time, mass, and charge), each dimension subset  $R$ . Where the dimensions of Euclidean distance, time, mass, and charge are each divided into a countably infinite number of constant-sized intervals and for all  $i$  in  $N$ , there are 3 direct and 3 inverse proportion ratios of the  $i$ -th unit interval length of Euclidean distance to the  $i$ -th unit interval lengths of time, mass, and charge. Derivations of well-known gravity, charge, electromagnetic, relativity, quantum physics equations and related constants from the ratios are much shorter and simpler. Inverse ratios are used to add quantum extensions to gravity, charge, and relativity equations. The math proofs are verified in Rocq.

[1]

## CONTENTS

|                                               |    |
|-----------------------------------------------|----|
| Introduction                                  | 3  |
| Insufficient reduction                        | 3  |
| First reduction step                          | 4  |
| Second reduction step                         | 5  |
| Third reduction step                          | 7  |
| Volume                                        | 8  |
| Euclidean volume                              | 8  |
| Distance                                      | 8  |
| Vector inner product                          | 8  |
| Minkowski distance ( $p$ -norm)               | 9  |
| Distance inequalities                         | 9  |
| Metric Space                                  | 10 |
| Property limiting a set to at most 3 elements | 11 |

|                                                        |    |
|--------------------------------------------------------|----|
| Derivation of physics equations and constants          | 14 |
| Direct and inverse proportion unit ratios              | 14 |
| $G$ , Newton, Gauss, Poisson gravity laws              | 15 |
| Special relativity                                     | 15 |
| Simple general relativity solutions                    | 16 |
| Schwarzschild's time dilation and metric               | 17 |
| $k_e$ , $\varepsilon_0$ , $E$ , Coulomb and Gauss laws | 18 |
| $B$ , $\mu_0$ , and Lorentz's law                      | 19 |
| Faraday's law                                          | 19 |
| Maxwell's wave equations                               | 20 |
| The fine structure constant, $\alpha$                  | 21 |
| $\hbar$ , $h$ , and Einstein-Planck relation           | 21 |
| Values of the unit ratios and Planck units             | 22 |
| Compton wavelength, $\lambda$                          | 23 |
| Schrödinger's wave equation                            | 23 |
| Dirac's wave equation                                  | 24 |
| Total of a type                                        | 24 |
| Quantum extension to Schwarzschild's metric            | 24 |
| Quantum extension to Newton's gravity force            | 25 |
| Quantum extension to Coulomb's charge force            | 26 |
| Insights and implications                              | 26 |
| References                                             | 32 |

**Keywords:** mathematical physics, gravity, charge, electromagnetism, relativity, quantum physics.

## INTRODUCTION

### Insufficient reduction

The trend in physics has been for increasingly complicated mathematics, assumptions, and equations, with increasing difficulty in finding exact solutions to the equations. Exam-

ples are Einstein’s field equations (general theory of relativity)[2, 3] and Schrödinger’s wave equation[4].

And physics equations often rely on assumptions that have not had a mathematical justification. For example, relativity and quantum theory both assume: 1) 3 dimensions of space and 2) the laws of physics are the same at every local Euclidean-like coordinate point[2–5]. But integration, measure theory, Hilbert spaces, linear algebra, (pseudo-)Riemann manifolds, and so on, have no restriction on the number of dimensions [3, 6–9]. And no properties of real and complex numbers have been identified that require functions to be the same at every Euclidean-like coordinate point[6, 7].

Further, much of physics relies on volume and distance equations. Real analysis defines Euclidean volume as the product of interval lengths[6, 7] and defines the Euclidean distance equation, dot product, inner product, and metric space[6, 7]. Justification for the definitions is vigorous finger-pointing to Euclidean and Cartesian geometry.

The philosophy of reductionism posits that complex phenomena can be reduced to a few simple parts: a few simple objects, relations, operations, etc. The first reduction step, in this article, is to prove that well-known geometry equations are reduced to (derived from) a few simple set and number properties. The second step is to identify some consequences of the properties of volume and distance. The third reduction step is to show that well-known physics equations, constants, and assumptions are reduced to (derived from) those consequences.

### **First reduction step**

This section is an overview the first step of proving that well-known geometry equations are reduced to (derived from) a few simple relations between countable set elements.

Each math proof, in this article, has a corresponding formal proof, which has been verified using the Rocq proof verification system [10], a verification tool used by mathematicians world-wide. The formal proofs are in the Rocq files, “euclidrelations.v” and “threed.v,” which are included as ancillary files.

Let  $|x_i|$  be the cardinal of (number of elements in) the abstract, countable set,  $x_i \in \{x_1, \dots, x_n\}$ . Let the range value,  $v_c$ , be the number of ordered combinations (n-tuples) of

the elements of  $x_1, \dots, x_n$ :

$$x_i \in \{x_1, \dots, x_n\}, \quad \bigcap_{i=1}^n x_i = \emptyset, \quad |x_i| \in \{0, \mathbb{N}\} \quad \wedge \quad v_c = \prod_{i=1}^n |x_i|. \quad (1)$$

Where each n-tuple corresponds to a real value,  $\kappa \in \mathbb{R}$ , it will be proved that:

$$v_c = \prod_{i=1}^n |x_i| \quad \Rightarrow \quad v = \prod_{i=1}^n s_i, \quad v, s_i \in \mathbb{R}. \quad (2)$$

For all  $n > 1$ , there are cases where different domain values,  $|x_1|, \dots, |x_n|$ , multiplied yield the same range value,  $v_c$ . Inferring a domain value,  $d_c$ , from  $v_c$ , requires an inverse (bijective) function,  $d_c = f_n^{-1}(v_c)$  and  $v_c = f_n(d_c)$ . The simplest bijective case for all  $n$  that includes the  $n = 1$  case is the cuboid case,  $|x_i| = d_c$ . And it will be proved that:

$$v_c = \prod_{i=1}^n |x_i| \quad \wedge \quad d_c = |x_i| \quad \Rightarrow \quad v = d^n. \quad (3)$$

$v_c$  is the sum of the number of subset n-tuples. It will be proved that:

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} \quad \Rightarrow \quad d^n = v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad (4)$$

Note that volume,  $v$ , is a scalar quantity and where each  $s_{i,j}$  is  $\pm$ -signed, the  $n = 2$  case is the dot product:

$$v = \sum_{i=1}^m (\prod_{j=1}^{n=2} s_{i,j}) \quad \wedge \quad \exists a_i, b_i \in \mathbb{R} : s_{i,1} = a_i, s_{i,2} = b_i \quad \Rightarrow \quad v = \sum_{i=1}^m a_i b_i \stackrel{\text{def}}{=} \mathbf{a} \cdot \mathbf{b}. \quad (5)$$

The inner product is where  $a_i$  and  $b_i$  are also complex numbers, functions, and matrices.

And, where each summed volume,  $v_{c_i}$ , is the inverse (bijective) function,  $d_{c_i}^n = v_{c_i}$ :

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m d_{c_i}^n \quad \Rightarrow \quad d^n = \sum_{i=1}^m d_i^n. \quad (6)$$

$|d|$  is the  $p$ -norm (Minkowski distance)[11], which will be proved to imply the metric space properties[7]. The  $n = 2$  case is the Euclidean distance.

## Second reduction step

This section provides the rationale leading to the identification of two consequences of the set and number properties generating the geometry equations.

The commutative property of the union of subsets of n-tuples and the multiplication operation defining the number n-tuples and corresponding Euclidean volume (2) allow  $n$

number of domain elements to be sequenced in all  $n!$  permutations, for example,  $v = \text{width} \times \text{height} \times \text{depth} = \text{depth} \times \text{width} \times \text{height} = \dots$ . The commutative property implies that each domain element has no intrinsic property that makes an element the first, second, ... , or last, which is a set of unordered elements.

Reliably sequencing a set of unordered elements in the same order requires that the elements be placed into a “list,” which assigns relative sequential positions, for example, labeling the unordered elements as  $x_1, x_2, \dots, x_n$ . The list can be sequenced via the successor and predecessor functions,  $\text{successor}(x_i) = x_{i+1}$  and  $\text{predecessor}(x_i) = x_{i-1}$ . For example, assigning  $x_1 = \text{width}$ ,  $x_2 = \text{height}$ , and  $x_3 = \text{depth}$ , the successor sequence,  $v = \prod_{i=1}^3 s_i$ , reliably sequences the unordered set elements in the order, width→height→depth.

Only a cyclic list, where  $\text{successor}(x_n) = x_1$  and  $\text{predecessor}(x_1) = x_n$ , allows each unordered element to be the first element of some of those  $n!$  permutations. For example, where  $n = 3$ , the volume cyclic successor sequences are:  $v = s_1 \times s_2 \times s_3$ ,  $v = s_2 \times s_3 \times s_1$ ,  $v = s_3 \times s_1 \times s_2$ , and the cyclic predecessor sequences are:  $v = s_1 \times s_3 \times s_2$ ,  $v = s_2 \times s_1 \times s_3$ ,  $v = s_3 \times s_2 \times s_1$ .

The only cyclic lists that allows sequencing the unordered elements in all  $n!$  permutations, is where each list element is sequentially adjacent (either an *immediate* cyclic successor or an *immediate* cyclic predecessor) to every other element, referred to, in this article, as an immediate symmetric cyclic list (ISCL). For example, the cyclic permutations,  $[x_2, x_3, x_1]$  and  $[x_2, x_1, x_3]$ , require  $x_1$  and  $x_3$  be sequentially adjacent. An ISCL will be proved to contain  $n \leq 3$  elements.

There are two consequence of the ISCL property of volume and the property of distance being bijective function of a cuboid volume:

1. A consequence of the commutative property of volume is an ISCL of at most 3 unordered distance dimensions,  $\{r_1, r_2, r_3\}$ , and an ISCL of at most 3 unordered non-distance dimensions  $\{x, y, z\}$ , each dimension  $\subseteq \mathbb{R}$ .
2. A consequence of Euclidean distance being derived as inverse (bijective) functions of a cuboid volume,  $v \in \mathbb{R}$ , is the 3-dimensional Euclidean distance dimension  $\subseteq \mathbb{R}$ .

### Third reduction step

This section is an overview of the second step of reducing well-known physics equations to the ISCL property of 3 distance dimensions and 3 non-distance dimensions, each dimension  $\subseteq \mathbb{R}$ , and the property of 3-dimensional Euclidean distance dimension  $\subseteq \mathbb{R}$ :

1. **ISCL:**  $\{r_1, r_2, r_3\}$  is an ISCL of 3 “distance” dimensions and  $\{t \text{ (time)}, m \text{ (mass)}, q \text{ (charge)}\}$  is the ISCL of 3 “non-distance” dimensions, each dimension  $\subseteq \mathbb{R}$ . There is a total of 6 unordered dimensions with the  $\subseteq \mathbb{R}$  property:  $r_1, r_2, r_3, t, m, q$ .
2. **Corresponding lengths:** Where the dimensions of Euclidean distance, time, mass, and charge are each divided into a countably infinite number of constant-sized intervals and  $i \in \mathbb{N}$ , the  $i^{th}$  unit interval length,  $r_p$  of Euclidean distance, corresponds to the  $i^{th}$  unit interval lengths:  $t_p$  of time,  $m_p$  of mass, and  $q_p$  of charge, where “unit” means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

The correspondence of unit interval lengths implies 3 direct proportion unit ratios:  $r/t = r_p/t_p$ ,  $rm = r_p/m_p$ , and  $rq = r_p/q_p$ , letting:  $c_t = r_p/t_p$ ,  $c_m = r_p/m_p$ , and  $c_q = r_p/q_p$ . And the correspondence of unit interval lengths implies 3 inverse proportion unit ratios:  $rt = t_p r_p$ ,  $rm = m_p r_p$ , and  $rq = q_p r_p$ . And let:  $k_t = t_p r_p$ ,  $k_m = m_p r_p$ , and  $k_q = q_p r_p$ .

The gravity[12, 13], special relativity[2, 3], Schwarzschild black hole metric[14, 15], charge[13], electromagnetism[13] equations and related constants are derived from the 3 direct proportion unit ratios. The Planck relation[13], Compton wavelength[13], Schrödinger wave[4], Dirac wave[5] equations and related constants are derived from combining the 3 direct and 3 inverse proportion unit ratios.

The quantum units:  $r_p$ ,  $t_p$ ,  $m_p$ , and  $q_p$  are derived from the property:

$$r_p = \sqrt{r_p^2} = \sqrt{c_t k_t} = \sqrt{c_m k_m} = \sqrt{c_q k_q}. \quad (7)$$

And it will be shown that  $r_p$ ,  $t_p$ ,  $m_p$ , and  $q_p$  are the reduced Planck units.

The inverse proportion ratios are used to add quantum extensions to Scharwzchild’s metric, Newton’s gravity, and Coulomb’s charge equations. The quantum-extended gravity, charge, and relativity equations predict that where quantum effects are measurable, the constants: gravity,  $G$ , Einstein’s  $\kappa = 8\pi G/c^4$ , charge,  $k_e$ , permittivity,  $\varepsilon_0$ , and permeability,  $\mu_0$ , do not exist. And the quantum-extended equations also predict that as distance between two point masses and between two point charges go to zero, the forces do not go to infinity.

## VOLUME

### Euclidean volume

**Theorem .1.** *Euclidean volume,*

$$\forall v_c, d_c, |x_i| \in \{0, \mathbb{N}\}, x_i \in \{x_1, \dots, x_n\}, v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i, \quad s_i, v \in \mathbb{R}. \quad (8)$$

The formal proof, “Euclidean\_volume,” is in the Rocq file, euclidrelations.v.

*Proof.*

$$\forall v_c, |x_i| \in \{0, \mathbb{N}\}, \kappa \in \mathbb{R} : v_c = \prod_{i=1}^n |x_i| \Leftrightarrow v_c \kappa = (\prod_{i=1}^n |x_i|) \kappa = \prod_{i=1}^n (|x_i| \kappa^{1/n}). \quad (9)$$

$$v_c \kappa = \prod_{i=1}^n (|x_i| \kappa^{1/n}) \wedge \exists v, s_i \in \mathbb{R} : v = v_c \kappa \wedge s_i = |x_i| \kappa^{1/n} \Rightarrow v = \prod_{i=1}^n s_i. \quad \square \quad (10)$$

**Lemma .2.** *The number of  $n$ -tuples,  $v_c$ , is the sum of the number of  $n$ -tuples,  $v_{c_i}$ , in each disjoint subset of  $n$ -tuples, implies a volume is the sum of volumes,*

$$v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v = \sum_{i=1}^m v_i, \quad v, v_i \in \mathbb{R}.$$

The formal proof, “sum\_of\_volumes,” is in the Rocq file, euclidrelations.v.

*Proof.* From the condition of this theorem and the Euclidean volume theorem (.1):

$$\forall \kappa \in \mathbb{R} : v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v_c \kappa = (\sum_{i=1}^m v_{c_i}) \kappa = \sum_{i=1}^m (v_{c_i} \kappa). \quad (11)$$

$$v_c \kappa = \sum_{i=1}^m (v_{c_i} \kappa) \wedge v = v_c \kappa \wedge v_i = v_{c_i} \kappa \Rightarrow v = \sum_{i=1}^m v_i. \quad \square \quad (12)$$

## DISTANCE

**Definition .3.** Bijective, countable domain value,  $d_c$ :

$$\exists v_c, d_c, |x_i| \in \{0, \mathbb{N}\}, x_i \in \{x_1, \dots, x_n\}, v_c = \prod_{i=1}^n |x_i| = \prod_{i=1}^n d_c = d_c^n. \quad (13)$$

### Vector inner product

**Theorem .4.** *Sum of volumes distance:*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} \Rightarrow d^n = v = \sum_{i=1}^m (\prod_{j=1}^n s_{ij}).$$

The formal proof, “sum\_of\_volumes\_distance,” is in the Rocq file, euclidrelations.v.



*Proof.* Multiply both sides of the assumption by  $\kappa$  and apply the Euclidean volume proof (.1) to  $v_i$ :

$$v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v = \sum_{i=1}^m v_i, \quad v, v_i \in \mathbb{R}. \quad (14)$$

$$v = \sum_{i=1}^m v_i \quad \wedge \quad v_i = \prod_j^n s_{i,j} \Rightarrow v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad (15)$$

$$d_c^n = v_c \Rightarrow d_c^n \kappa = (d_c \kappa^{1/n})^n = v_c \kappa. \quad (16)$$

$$\exists d, v \in \mathbb{R} : d = d_c \kappa^{1/n} \quad \wedge \quad v = v_c \kappa \quad \wedge \quad (d_c \kappa^{1/n})^n = v_c \kappa \Rightarrow d^n = v. \quad (17)$$

$$d^n = v \quad \wedge \quad v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}) \Rightarrow d^n = v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad \square \quad (18)$$

Volume,  $v$ , is a scalar quantity and where each  $s_{i,j}$  is  $\pm$ -signed, the  $n = 2$  case is the dot product:

$$v = \sum_{i=1}^m (\prod_{j=1}^{n=2} s_{i,j}) \quad \wedge \quad \exists a_{i,1}, a_{i,2} \in \mathbb{R} : s_{i,1} = a_i, s_{i,2} = b_i \Rightarrow v = \sum_{i=1}^m a_i b_i \stackrel{\text{def}}{=} \mathbf{a} \cdot \mathbf{b}. \quad (19)$$

The inner product is where  $a_i$  and  $b_i$  are also complex numbers, functions, and matrices.

### Minkowski distance ( $p$ -norm)

**Theorem .5.** *Minkowski distance ( $p$ -norm):*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m d_{c_i}^n \Leftrightarrow d^n = \sum_{i=1}^m d_i^n.$$

*The formal proof, "Minkowski\_distance," is in the Rocq file, euclidrelations.v.*

*Proof.* From the sum of volumes distance theorem .4:

$$d_{c_i}^n = v_{c_i} \Rightarrow d_i^n = v_i. \quad (20)$$

$$d^n = v = \sum_{i=1}^m v_i \quad \wedge \quad d_i^n = v_i \Rightarrow d^n = \sum_{i=1}^m d_i^n \quad \square \quad (21)$$

### Distance inequalities

**Theorem .6.** *Distance triangle inequality*

$$\forall n \in \mathbb{N}, v_a, v_b \geq 0 : (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}.$$

*The formal proof, distance\_inequality, is in the Rocq file, euclidrelations.v.*

*Proof.* Expand  $(v_a^{1/n} + v_b^{1/n})^n$  using the binomial expansion:

$$\begin{aligned} \forall v_a, v_b \geq 0 : \quad v_a + v_b &\leq v_a + v_b + \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^{n-i} (v_b^{1/n})^i + \\ &\quad \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^i (v_b^{1/n})^{n-i} = (v_a^{1/n} + v_b^{1/n})^n. \end{aligned} \quad (22)$$

Take the  $n^{th}$  root of both sides of the inequality 22:

$$\forall v_a, v_b \geq 0, \quad n \in \mathbb{N} : \quad v_a + v_b \leq (v_a^{1/n} + v_b^{1/n})^n \quad \Rightarrow \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}. \quad (23)$$

□

**Theorem .7.** *Distance sum inequality*

$$\forall m, n \in \mathbb{N}, \quad a_i, b_i \geq 0 : (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}.$$

The formal proof, *distance\_sum\_inequality*, is in the Rocq file, *euclidrelations.v*.

*Proof.* Apply the distance triangle inequality (.6):

$$\begin{aligned} \forall m, n \in \mathbb{N}, \quad v_a, v_b \geq 0 : \quad v_a &= \sum_{i=1}^m a_i^n \quad \wedge \quad v_b = \sum_{i=1}^m b_i^n \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\ \Rightarrow \quad ((\sum_{i=1}^m a_i^n) &+ (\sum_{i=1}^m b_i^n))^{1/n} = (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq \\ &(\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad \square \end{aligned} \quad (24)$$

## Metric Space

All Minkowski distances ( $p$ -norms) imply the metric space properties. The formal proofs: *triangle\_inequality*, *symmetry*, *non\_negativity*, and *identity\_of\_indiscernibles* are in the Rocq file, *euclidrelations.v*.

**Theorem .8.** *Triangle Inequality:*

$$\begin{aligned} d(s_1, s_2) &= (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k \\ \Rightarrow \quad d(u, w) &= (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \end{aligned}$$

*Proof.*  $\forall p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k:$

$$(u^p + w^p)^{1/p} \leq ((u^p + w^p) + 2v^p)^{1/p} = ((u^p + v^p) + (v^p + w^p))^{1/p}. \quad (25)$$

Apply the distance triangle inequality (.6) to the inequality 25:

$$\begin{aligned}
(u^p + w^p)^{1/p} &\leq ((u^p + v^p) + (v^p + w^p))^{1/p} \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\
&\quad \wedge \quad v_a = u^p + v^p \quad \wedge \quad v_b = v^p + w^p \\
&\Rightarrow (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} \\
&\Rightarrow d(u, w) = (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \quad \square \quad (26)
\end{aligned}$$

**Theorem .9.** *Symmetry:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad v = s_2 \quad \Rightarrow \quad d(u, v) = d(v, u).$$

*Proof.* By the commutative law of addition:

$$\begin{aligned}
\forall p : p \geq 1, \quad d(s_1, s_2) &= (\sum_{i=1}^2 s_i^p)^{1/p} = (s_1^p + s_2^p)^{1/p} \\
&\Rightarrow d(u, v) = (u^p + v^p)^{1/p} = (v^p + u^p)^{1/p} = d(v, u). \quad \square \quad (27)
\end{aligned}$$

**Theorem .10.** *Non-negativity:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \in N \quad \wedge \quad s_1, s_2 \geq 0 \quad \Rightarrow \quad d(u, v) \geq 0.$$

*Proof.*

$$\forall n \geq 1 \quad \wedge \quad s_1, s_2 \geq 0 \quad \Rightarrow \quad d(s_1, s_2) = (s_1^p + s_2^p)^{1/p} \geq 0. \quad (28)$$

$$\forall u = s_1, v = s_2 \quad \wedge \quad d(s_1, s_2) = (s_1^p + s_2^p)^{1/p} \geq 0 \quad \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} \geq 0. \quad \square \quad (29)$$

**Theorem .11.** *Identity of Indiscernibles:*  $d(u, u) = 0$ .

*Proof.* From the non-negativity property (.10):

$$d(u, w) \geq 0 \quad \wedge \quad d(u, v) \geq 0 \quad \wedge \quad d(v, w) \geq 0 \quad (30)$$

$$\Rightarrow \exists d(u, w) = d(u, v) = d(v, w) = 0. \quad (31)$$

$$d(u, w) = d(v, w) = 0 \quad \Rightarrow \quad u = v. \quad (32)$$

$$d(u, v) = 0 \quad \wedge \quad u = v \quad \Rightarrow \quad d(u, u) = 0. \quad \square \quad (33)$$

## PROPERTY LIMITING A SET TO AT MOST 3 ELEMENTS

The following definitions and proof use first order logic. A Horn clause-like expression is used, here, to make the proof easier to read. By convention, the proof goal, in a Horn clause,

is on the left side of the implication sign ( $\leftarrow$ ) and supporting facts are on the right side. The formal proofs in the Rocq file `threed.v` are: `adj111`, `adj122`, `adj212`, `adj123`, `adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and `not_all_mutually_adjacent_gt_3`.

**Definition .12.** Immediate Cyclic Successor of  $m$  is  $n$ :

$$\begin{aligned} & \forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \\ \text{Successor}(m, n, \text{setsize}) & \leftarrow (m = \text{setsize} \wedge n = 1) \quad \vee \quad (n = m + 1 \leq \text{setsize}). \end{aligned} \quad (34)$$

**Definition .13.** Immediate Cyclic Predecessor of  $m$  is  $n$ :

$$\begin{aligned} & \forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \\ \text{Predecessor}(m, n, \text{setsize}) & \leftarrow (m = 1 \wedge n = \text{setsize}) \quad \vee \quad (n = m - 1 \geq 1). \end{aligned} \quad (35)$$

**Definition .14.** Adjacent: Element  $m$  is sequentially adjacent to element  $n$  if the immediate cyclic successor of  $m$  is  $n$  or the immediate cyclic predecessor of  $m$  is  $n$ . Notionally:

$$\begin{aligned} & \forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \\ \text{Adjacent}(m, n, \text{setsize}) & \leftarrow \text{Successor}(m, n, \text{setsize}) \vee \text{Predecessor}(m, n, \text{setsize}). \end{aligned} \quad (36)$$

**Definition .15.** Immediate Symmetric Cyclic List (ISCL), where every set element is sequentially adjacent to every other element):

$$\forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \quad \text{Adjacent}(m, n, \text{setsize}). \quad (37)$$

**Theorem .16.** An ISCL is limited to at most 3 elements.

$$\forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} : \quad \text{Adjacent}(m, n, \text{setsize}) \Rightarrow \text{setsize} \leq 3. \quad (38)$$

*Proof.*

Every element is adjacent to every other element, where  $setsize \in \{1, 2, 3\}$ :

$$Adjacent(1, 1, 1) \leftarrow Successor(1, 1, 1) \leftarrow (m = setsize \wedge n = 1). \quad (39)$$

$$Adjacent(1, 2, 2) \leftarrow Successor(1, 2, 2) \leftarrow (n = m + 1 \leq setsize). \quad (40)$$

$$Adjacent(1, 2, 3) \leftarrow Successor(1, 2, 3) \leftarrow (n = m + 1 \leq setsize). \quad (41)$$

$$Adjacent(2, 1, 3) \leftarrow Predecessor(2, 1, 3) \leftarrow (n = m - 1 \geq 1). \quad (42)$$

$$Adjacent(3, 1, 3) \leftarrow Successor(3, 1, 3) \leftarrow (n = setsize \wedge m = 1). \quad (43)$$

$$Adjacent(1, 3, 3) \leftarrow Predecessor(1, 3, 3) \leftarrow (m = 1 \wedge n = setsize). \quad (44)$$

$$Adjacent(2, 3, 3) \leftarrow Successor(2, 3, 3) \leftarrow (n = m + 1 \leq setsize). \quad (45)$$

$$Adjacent(3, 2, 3) \leftarrow Predecessor(3, 2, 3) \leftarrow (n = m - 1 \geq 1). \quad (46)$$

Element 2 is the only immediate successor of element 1 for all  $setsize \geq 3$ , which implies element 3 is not ( $\neg$ ) an immediate successor of element 1 for all  $setsize \geq 3$ :

$$\neg Successor(1, 3, setsize \geq 3) \leftarrow Successor(1, 2, setsize \geq 3) \leftarrow (n = m + 1 \leq setsize). \quad (47)$$

Element  $n = setsize > 3$  is the only immediate predecessor of element 1, which implies element 3 is not ( $\neg$ ) an immediate predecessor of element 1 for all  $setsize > 3$ :

$$\neg Predecessor(1, 3, setsize \geq 3) \leftarrow Predecessor(1, setsize, setsize > 3) \leftarrow (m = 1 \wedge n = setsize > 3). \quad (48)$$

For all  $setsize > 3$ , some elements are not ( $\neg$ ) sequentially adjacent to every other element (not immediate symmetric):

$$\neg Adjacent(1, 3, setsize > 3) \leftarrow \neg Successor(1, 3, setsize > 3) \quad \wedge \quad \neg Predecessor(1, 3, setsize > 3). \quad \square \quad (49)$$

The Symmetric goal matches the Adjacent goals 39 and fails for all “setsize” greater than three.

## DERIVATION OF PHYSICS EQUATIONS AND CONSTANTS

### Direct and inverse proportion unit ratios

Application of the ISCL property of volume (.16) and the 3-dimensional Euclidean distance dimension  $\subseteq \mathbb{R}$  (.5):

1. **ISCL:**  $\{r_1, r_2, r_3\}$  is an ISCL of 3 “distance” dimensions and  $\{t$  (*time*),  $m$  (*mass*),  $q$  (*charge*) $\}$  is the ISCL of 3 “non-distance” dimensions, each dimension  $\subseteq \mathbb{R}$ . There is a total of 6 unordered dimensions with the  $\subseteq \mathbb{R}$  property:  $r_1, r_2, r_3, t, m, q$ .
2. **Corresponding lengths:** Where the dimensions of Euclidean distance, time, mass, and charge are each divided into a countably infinite number of constant-sized intervals and  $i \in \mathbb{N}$ , the  $i^{th}$  unit interval length,  $r_p$  of Euclidean distance, corresponds to the  $i^{th}$  unit interval lengths:  $t_p$  of time,  $m_p$  of mass, and  $q_p$  of charge, where “unit” means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

Their are as many unit interval lengths of  $r_p$  as the unit interval lengths  $t_p, m_p$ , and  $q_p$ , can be expressed as the direct proportion ratios,  $r/\tau = r_p/\tau_p$ , where  $\tau \in \{t, m, q\}$ .

$$\begin{aligned} \forall r_p, \tau_p, r, \tau \subseteq \mathbb{R}, r_p, \tau_p \text{ constants}, \tau \in \{t, m, q\} : r/\tau = r_p/\tau_p &\Rightarrow r = (r_p/\tau_p)/\tau \\ &\Rightarrow r = (r_p/t_p)t \quad \wedge \quad r = (r_p/m_p)m \quad \wedge \quad r = (r_p/q_p)q. \end{aligned} \quad (50)$$

And for conciseness, let:

$$r_p/t_p = c_t \quad \wedge \quad r_p/m_p = c_m \quad \wedge \quad r_p/q_p = c_q. \quad (51)$$

And there are 3 inverse proportion ratios:

$$\begin{aligned} r/t = r_p/t_p \quad \wedge \quad r/m = r_p/m_p &\Rightarrow (r/t)/(r/m) = (r_p/t_p)/(r_p/m_p) \\ &\Rightarrow (mr)/(tr) = (m_p r_p)/(t_p r_p) \Rightarrow \exists r, t, m \in \mathbb{R} : mr = m_p r_p, \quad tr = t_p r_p. \end{aligned} \quad (52)$$

$$\begin{aligned} r/t = r_p/t_p \quad \wedge \quad r/q = r_p/q_p &\Rightarrow (r/t)/(r/q) = (r_p/t_p)/(r_p/q_p) \\ &\Rightarrow (qr)/(tr) = (q_p r_p)/(t_p r_p) \Rightarrow \exists r, t, q \in \mathbb{R} : qr = q_p r_p, \quad tr = t_p r_p. \end{aligned} \quad (53)$$

And for conciseness, let:

$$t_p r_p = k_t \quad \wedge \quad m_p r_p = k_m \quad \wedge \quad q_p r_p = k_q. \quad (54)$$

## **G, Newton, Gauss, Poisson gravity laws**

From equations 51, linear acceleration,  $r/t^2$ , can be related to mass,  $m$ , and distance,  $r$ :

$$r = c_m m \quad \wedge \quad r = c_t t \quad \Rightarrow \quad r/(c_t t)^2 = c_m m/r^2 \quad \Rightarrow \quad r/t^2 = (c_m c_t^2) m/r^2 = Gm/r^2, \quad (55)$$

where  $G = c_m c_t^2 = r_p^3/m_p t_p^2$ , conforms to the SI units:  $m^3 \cdot kg^{-1} \cdot s^{-2}$  [12].

Newton's law follows from multiplying both sides of equation 55 by  $m$ :

$$r/t^2 = Gm/r^2 \quad \Leftrightarrow \quad F \stackrel{\text{def}}{=} mr/t^2 = Gm^2/r^2. \quad (56)$$

$$F = Gm^2/r^2 \quad \wedge \quad \forall m \in \mathbb{R} : \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \quad \Rightarrow \quad F = Gm_1 m_2/r^2. \quad (57)$$

A new interpretation of Gauss's and Poisson's gravity laws is presented here. Equation 55 relates linear acceleration,  $r/t^2$ , to mass and distance. Gauss's gravity field,  $\mathbf{g}$ , and Poisson's gravity field,  $-\nabla\Phi(\tilde{\mathbf{r}}, t)$ , is the angular (orbital) acceleration,  $2\pi r/t^2$ . Applying equation 55:

$$r/t^2 = -Gm/|\tilde{\mathbf{r}}|^2 \quad \wedge \quad \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = 2\pi r/t^2 \quad \Rightarrow \quad \mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = -2\pi Gm/|\tilde{\mathbf{r}}|^2 \quad (58)$$

$$\mathbf{g} = -\nabla\Phi(\tilde{\mathbf{r}}, t) = -2\pi Gm/|\tilde{\mathbf{r}}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{g} = \nabla^2\Phi(\tilde{\mathbf{r}}, t) = 4\pi Gm/|\tilde{\mathbf{r}}|^3 = 4\pi G\rho, \quad (59)$$

where  $\rho = m/|\tilde{\mathbf{r}}|^3$  is the mass density:

## **Special relativity**

$$\begin{aligned} \forall \tau \in \{t, m, q\}, \quad r^2 = r'^2 + r_v^2, \quad \exists \mu, \nu : r = \mu\tau \quad \wedge \quad r_v = \nu\tau \quad \Rightarrow \quad (\mu\tau)^2 = r'^2 + (\nu\tau)^2 \\ \Rightarrow \quad r' = \sqrt{(\mu\tau)^2 - (\nu\tau)^2} = \mu\tau\sqrt{1 - (\nu/\mu)^2}. \end{aligned} \quad (60)$$

Local frame distance,  $r'$ , contracts relative to a distant observer frame distance,  $r$ , as  $\nu \rightarrow \mu$ :

$$r' = \mu\tau\sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad \mu\tau = r \quad \Rightarrow \quad r' = r\sqrt{1 - (\nu/\mu)^2}. \quad (61)$$

A distant observer frame type,  $\tau$ , dilates relative to the local observer frame type,  $\tau'$ , as  $\nu \rightarrow \mu$ :

$$\mu\tau = r'/\sqrt{1 - (\nu/\mu)^2} \quad \wedge \quad r' = \mu\tau \quad \Rightarrow \quad \tau = \tau'/\sqrt{1 - (\nu/\mu)^2}. \quad (62)$$

Where  $\tau$  is type, time, the space-like flat Minkowski spacetime event interval is:

$$\begin{aligned} dr^2 = dr'^2 + dr_v^2 \quad \wedge \quad dr_v^2 = dr_1^2 + dr_2^2 + dr_3^2 \quad \wedge \quad d(\mu\tau) = dr \\ \Rightarrow \quad dr'^2 = d(\mu\tau)^2 - dr_1^2 - dr_2^2 - dr_3^2. \end{aligned} \quad (63)$$

## Simple general relativity solutions

The Einstein field equation (EFE) is:

$$G_{\mu,\nu} + \Lambda g_{\mu,\nu} = \kappa T_{\mu,\nu}, \quad G_{\mu,\nu} = \mathbf{R} + \frac{1}{2} R g_{\mu,\nu}. \quad (64)$$

where:  $G_{\mu,\nu}$  is Einstein's tensor,  $\mathbf{R}$  is the Ricci curvature,  $R$  is the scalar curvature,  $g_{\mu,\nu}$  is the metric tensor,  $\Lambda$  is a cosmological constant,  $\kappa = 8\pi G/c^4$  is Einstein's constant, and  $T_{\mu,\nu}$  is the stress-energy-momentum tensor [2, 3].

The EFE is nonlinear, which has made determining the measure (metric) of the space-time curvature,  $g_{\mu,\nu}$ , very difficult. In particular, it has been assumed that the measure of curvature,  $g_{\mu,\nu}$ , depends on the Ricci curvature,  $\mathbf{R}$ , and depends on the distribution of the stress, energy, and momentum,  $T_{\mu,\nu}$ , which bends spacetime [3, 14, 15].

But, in the next subsection, the Schwarzschild metric[14, 15] will be derived independent of the EFE by noting that: 1) The laws of physics are independent of the Ricci curvature and independent of the distribution of the stress, energy, and momentum. 2) The laws of physics are expressed in the components of the metric,  $g_{\mu,\nu}$ . This leads to the following simple steps to derive all metric tensors,  $g_{\mu,\nu}$ , independent of the EFE.

Step 1) Use the unit ratios and other equations derived from the ratios, for example, the ratio-derived special relativity equations(61) and the ratio-derived constant,  $G$  (55), to derive functions (laws) to be assigned to the components of  $g_{\nu,\mu}$ . For spherically symmetric cases, like the Schwarzschild metric, the ratio derived laws are assigned to the diagonal components of  $g_{\nu,\mu}$ . For non-symmetric cases, ratio-derived laws are also assigned to the other components of  $g_{\nu,\mu}$ .

Step 2) The 4D flat spacetime interval equation is an instance of the 2D equation (63). Therefore, derive the metric first as a 2D metric and expand later to a 4D metric.

(Optional step 2d) The 2D metric tensor allows using the much simpler 2D Ricci curvature (which, in 2D, is the Gaussian curvature) and scalar curvature. And the 2D tensors reduce the number of independent equations to solve. The 2D solutions can next be used to set constraints on the solutions in the 4D tensors.

Step 3) Translate from 2D to 4D. For spherically symmetric cases, the simplest method to translate from 2D to 4D is to use spherical coordinates, where the first 2 diagonal components of  $g_{\nu,\mu}$  remain unchanged and the other 2 diagonal components are functions of the longitude angle,  $\theta$ , and latitude angle,  $\phi$ .



### Schwarzschild's time dilation and metric

From equations 61 and 50:

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - (1)(v^2/c^2)} \quad \wedge \quad c_m m/r = 1 \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - c_m m v^2 / r c^2}.\end{aligned}\quad (65)$$

Where  $v_{escape}$  is the escape velocity:

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - c_m m v^2 / r c^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2 \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m v_{escape}^2 / r c^2}.\end{aligned}\quad (66)$$

For a photon, the escape velocity,  $v_{escape} = c$ .

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2c_m m v_{escape}^2 / r c^2} \quad \wedge \quad v_{escape} = c \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2}.\end{aligned}\quad (67)$$

Combining equation 67 with the derivation of  $G$  (55):

$$c_m c^2 = G \quad \wedge \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2} \Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm/r c^2}.\quad (68)$$

Combining equation 68 with equation 62 yields Schwarzschild's gravitational time dilation:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm/r c^2} \quad \wedge \quad t' = t\sqrt{1 - (v^2/c^2)} \Rightarrow t' = t\sqrt{1 - 2Gm/r c^2}.\quad (69)$$

From equations 61 and 68, where Schwarzschild defined the black hole event horizon radius as  $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$  [14, 15]:

$$r' = r\sqrt{1 - (v/c)^2} = r\sqrt{1 - 2Gm/r c^2} \quad \wedge \quad \alpha \stackrel{\text{def}}{=} 2Gm/c^2 \Rightarrow r' = r\sqrt{1 - \alpha/r}.\quad (70)$$

Applying equation 70 to the time-like spacetime interval equation 63, where  $ds^2 < 0$ :

$$\begin{aligned}r' &= r\sqrt{1 - \alpha/r} \quad \wedge \quad ds^2 = dr'^2 - dr^2 \\ \Rightarrow \quad ds^2 &= (\sqrt{1 - \alpha/r} dr)^2 - (dr'/\sqrt{1 - \alpha/r})^2 = (1 - \alpha/r)dr^2 - (1 - \alpha/r)^{-1}dr'^2.\end{aligned}\quad (71)$$

$$\begin{aligned}ds^2 &= (1 - \alpha/r)dr^2 - (1 - \alpha/r)^{-1}dr'^2 \quad \wedge \quad dr = d(ct) \quad \wedge \quad c = 1 \quad \wedge \\ \lim_{ds \rightarrow 0} dr' &= \lim_{ds \rightarrow 0} dr \Rightarrow \lim_{ds \rightarrow 0} ds^2 = (1 - \alpha/r)dt^2 - (1 - \alpha/r)^{-1}dr^2.\end{aligned}\quad (72)$$

Using spherical coordinates to translate from 2D to 4D yields the  $+- --$  form of Schwarzschild's black hole metric [14, 15]:

$$\begin{aligned} \lim_{ds \rightarrow 0} ds^2 &= (1 - \alpha/r) dt^2 - (1 - \alpha/r)^{-1} dr^2 \\ \Rightarrow \lim_{ds \rightarrow 0} ds^2 &= (1 - \alpha/r) dt^2 - (1 - \alpha/r)^{-1} dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2) \\ \Rightarrow g_{\mu,\nu} &= \text{diag}[(1 - \alpha/r), -(1 - \alpha/r)^{-1}, -r^2, -r^2 \sin^2\theta]. \end{aligned} \quad (73)$$

### $k_e, \varepsilon_0, \mathbf{E}$ , Coulomb and Gauss laws

From equations 51, linear acceleration,  $r/t^2$ , can be related to charge,  $q$ , and distance,  $r$ :

$$r = c_q q \quad \wedge \quad r = c_t t \quad \Rightarrow \quad r/(c_t t)^2 = c_q q/r^2 \quad \Rightarrow \quad r/t^2 = (c_q c_t^2) q/r^2. \quad (74)$$

$$r = c_m m \quad \wedge \quad r = c_q q \quad \Rightarrow \quad m = (c_q/c_m) q. \quad (75)$$

Combining equations 74 and 75 yields Coulombs's law:

$$r/t^2 = (c_q c_t^2) q/r^2 \quad \wedge \quad m = (c_q/c_m) q \quad \Rightarrow \quad F \stackrel{\text{def}}{=} mr/t^2 = (c_q^2 c_t^2/c_m) q^2/r^2 = k_e q^2/r^2. \quad (76)$$

$k_e = c_q^2 c_t^2/c_m = m_p r_p^3/t_p^2 q_p^2$ , conforms to the SI units:  $kg \cdot m^3 \cdot s^{-2} \cdot C^{-2} \stackrel{\text{def}}{=} N \cdot m^2 \cdot C^{-2}$  [13].

$$\forall q \in \mathbb{R} \exists q_1, q_2 \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad F = k_e q^2/r^2 \quad \Rightarrow \quad F = k_e q_1 q_2/r^2. \quad (77)$$

A new interpretation of Gauss's electric field law is presented here. Equation 74 relates the linear acceleration,  $r/t^2$ , to charge and distance. Gauss's electric field,  $\mathbf{E}$ , relates angular (orbital) acceleration,  $2\pi r/t^2$  to charge and distance. Applying equation 76:

$$F_C = mr/t^2 = k_e q^2/r^2 \quad \Rightarrow \quad \exists F_E \in \mathbb{R} : F_E = 2\pi(mr/t^2) = 2\pi k_e q^2/r^2. \quad (78)$$

$$F_E = 2\pi k_e q^2/r^2 = q(2\pi(k_e q/r^2)) \quad \wedge \quad \exists E \in \mathbb{R} : E = 2\pi k_e q/r^2 \quad \Rightarrow \quad F_E = qE. \quad (79)$$

The electric field,  $E = 2\pi k_e q/r^2$ , conforms to the SI units  $kg \cdot m \cdot s^{-2} \cdot C^{-1} \stackrel{\text{def}}{=} N \cdot C^{-1}$ .

$$\mathbf{E} = 2\pi k_e q/|\mathbf{r}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -4\pi k_e q/|\mathbf{r}|^3. \quad (80)$$

$$\nabla \cdot \mathbf{E} = -4\pi k_e q/|\mathbf{r}|^3 \quad \wedge \quad \varepsilon_0 \stackrel{\text{def}}{=} 1/4\pi k_e \quad \wedge \quad \rho \stackrel{\text{def}}{=} q/|\mathbf{r}|^3 \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -\rho/\varepsilon_0, \quad (81)$$

which is Gauss's electric field law [13] in differential form.

## **B, $\mu_0$ , and Lorentz's law**

Applying the distance contraction equation, 61, to equation 79, where  $r'$  is the distant observer frame of reference and  $r$  is local (rest) frame of reference:

$$r' = r/\sqrt{1 - v^2/c^2} \quad \wedge \quad F' = 2\pi k_e q^2/r'^2 \quad \Rightarrow \quad F = 2\pi k_e q^2(1 - v^2/c^2)/r^2. \quad (82)$$

From equation 79:

$$E = 2\pi k_e q/r^2 \quad \wedge \quad F = 2\pi k_e q^2(1 - v^2/c^2)/r^2 \quad \Rightarrow \quad F = q(E - ((2\pi k_e/c^2)q/r^2)v^2). \quad (83)$$

$$F = q(E - ((2\pi k_e/c^2)q/r^2)v^2) \quad \wedge \quad \exists B : B = (2\pi k_e/c^2)vq/r^2 \quad \Rightarrow \quad F = q(E - Bv). \quad (84)$$

Translating equation 84 to vector form:

$$F = q(E - Bv) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{v}). \quad (85)$$

$$\mathbf{B} \times \vec{v} = -(\vec{v} \times \mathbf{B}) \quad \wedge \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \vec{v}) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} + \vec{v} \times \mathbf{B}), \quad (86)$$

which is Lorentz law in the distant observer frame of reference, where the magnetic field,  $B = (2\pi k_e/c^2)vq/r^2$ , conforms to the base SI units:  $kg \cdot s^{-1} \cdot C^{-1} \stackrel{\text{def}}{=} kg \cdot s^{-2} \cdot A^{-1} \stackrel{\text{def}}{=} T$ .

Based on the assumption (having no math proof-motivation) of a magnetic dipole, the magnetic field divergence outward,  $\nabla \cdot \mathbf{B}$ , at a point,  $P$ , normal to a surface enclosing a charge equals the field divergence inward,  $\nabla \cdot \mathbf{B}$ :

$$\mathbf{B} = (2\pi k_e/c^2)vq/|\vec{r}|^2 \quad \Rightarrow \quad \nabla \cdot \mathbf{B} = -(4\pi k_e/c^2)q/|\vec{r}|^3 + (4\pi k_e/c^2)q/|\vec{r}|^3 = 0. \quad (87)$$

$$\nabla \cdot \mathbf{B} = -(4\pi k_e/c^2)q/|\vec{r}|^3 + (4\pi k_e/c^2)q/|\vec{r}|^3 = 0 \quad \Rightarrow \quad \mu_0 = (4\pi k_e/c^2), \quad (88)$$

where  $\mu_0 = 4\pi k_e/c^2$  conforms to the SI units  $kg \cdot m \cdot C^{-2} \stackrel{\text{def}}{=} kg \cdot m \cdot s^{-2} A^{-2}$ .

*Note:* In the last section of this article, it will be shown that the CODATA definition,  $\mu_0 = 2h\alpha/cq_e^2$  [16], and  $\mu_0 = 4\pi k_e/c^2$  both reduce to the same equation,  $\mu_0 = 4\pi k_m/q_p^2$ .

## **Faraday's law**

From the magnetic field equation 85, where the electric and magnetic fields are propagating at the speed,  $v = c$ :

$$B = (2\pi k_e/c^2)qv/r^2 \quad \wedge \quad v = c \quad \wedge \quad r = ct \quad \Rightarrow \quad B = (2\pi k_e/c^3)q/t^2. \quad (89)$$

$$B = (2\pi k_e/c^3)q/t^2 \quad \Rightarrow \quad \partial B/\partial t = -(4\pi k_e/c^3)q/t^3. \quad (90)$$

$$\partial B/\partial t = -(4\pi k_e/c^3)q/t^3 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial B/\partial t = -4\pi k_e q/r^3. \quad (91)$$

Faraday's law is the case, where the electric field,  $\mathbf{E}$ , is a single radial vector,  $\mathbf{E}_{\tilde{\mathbf{r}}_1}$  acting on some point,  $P$ . Therefore, at point,  $P$ ,  $\mathbf{E}_{\tilde{\mathbf{r}}_2} = 0$  and  $\mathbf{E}_{\tilde{\mathbf{r}}_3} = 0$ . From the Lorentz equation 86, the magnetic field,  $\mathbf{B}$  at point  $P$ , is perpendicular to  $\mathbf{E}_{\tilde{\mathbf{r}}_1}$  (aligned on  $\tilde{\mathbf{r}}_2$ ). For a constant force, the Lorentz equation makes the electric field,  $\mathbf{E}_{\tilde{\mathbf{r}}_1}$ , a function of  $\mathbf{B}_{\tilde{\mathbf{r}}_2}$ . Combining with equation 80:

$$\begin{aligned} \mathbf{E}_{\tilde{\mathbf{r}}_1} = 2\pi k_e q / |\tilde{\mathbf{r}}_2|^2 = f(\mathbf{B}_{\tilde{\mathbf{r}}_2}) \quad \wedge \quad \mathbf{E}_{\tilde{\mathbf{r}}_2} = 0 \quad \wedge \quad \mathbf{E}_{\tilde{\mathbf{r}}_3} = 0 \quad \Rightarrow \quad \nabla \times \mathbf{E} &= \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ \frac{\partial}{\partial \tilde{r}_1^2} & \frac{\partial}{\partial \tilde{r}_2^2} & \frac{\partial}{\partial \tilde{r}_3^2} \\ \mathbf{E}_{\tilde{\mathbf{r}}_1} & 0 & 0 \end{vmatrix} \\ &= \left( \frac{\partial \mathbf{E}_{r_3}}{\partial \tilde{r}_2^2} - \frac{\partial \mathbf{E}_{r_2}}{\partial \tilde{r}_3^2} \right) \hat{\mathbf{i}} + \left( \frac{\partial \mathbf{E}_{r_1}}{\partial \tilde{r}_3^2} - \frac{\partial \mathbf{E}_{r_3}}{\partial \tilde{r}_1^2} \right) \hat{\mathbf{j}} + \left( \frac{\partial \mathbf{E}_{r_2}}{\partial \tilde{r}_1^2} - \frac{\partial \mathbf{E}_{r_1}}{\partial \tilde{r}_2^2} \right) \hat{\mathbf{k}} \\ &= 0\hat{\mathbf{i}} + 0\hat{\mathbf{j}} + \left( 0 - \frac{\partial 2\pi k_e q / |\tilde{\mathbf{r}}_2|^2}{\partial |\tilde{\mathbf{r}}_2|^2} \right) \hat{\mathbf{k}} = 4\pi k_e q / |\tilde{\mathbf{r}}_2|^3 \hat{\mathbf{k}} \equiv \nabla \times \mathbf{E} = 4\pi k_e q / |\tilde{\mathbf{r}}|^3. \end{aligned} \quad (92)$$

Combining equations 92 and 91 yields Maxwell-Faraday's law [13]:

$$\nabla \times \mathbf{E} = 4\pi k_e q / |\tilde{\mathbf{r}}|^3 \quad \wedge \quad \partial \mathbf{B} / \partial t = -4\pi k_e q / |\tilde{\mathbf{r}}|^3 \quad \Rightarrow \quad \nabla \times \mathbf{E} = -\partial \mathbf{B} / \partial t. \quad (93)$$

### Maxwell's wave equations

From the Gauss's electric field equation 81:

$$\mathbf{E} = 2\pi k_e q / |\tilde{\mathbf{r}}|^2 \quad \wedge \quad r = |\tilde{\mathbf{r}}| \quad \Rightarrow \quad \nabla^2 \mathbf{E} = 12\pi k_e q / |\tilde{\mathbf{r}}|^4. \quad (94)$$

$$\mathbf{E} = 2\pi k_e q / |\tilde{\mathbf{r}}|^2 \quad \wedge \quad r = ct \quad \Rightarrow \quad \mathbf{E} = 2\pi k_e q / (ct)^2 \quad \Rightarrow \quad \partial^2 \mathbf{E} / \partial t^2 = (12\pi k_e / c^2) q / t^4. \quad (95)$$

$$\partial^2 \mathbf{E} / \partial t^2 = (12\pi k_e / c^2) q / t^4 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial^2 \mathbf{E} / \partial t^2 = 12\pi k_e c^2 q / |\tilde{\mathbf{r}}|^4. \quad (96)$$

$$\nabla^2 \mathbf{E} = 12\pi k_e q / |\tilde{\mathbf{r}}|^4 \quad \wedge \quad \partial^2 \mathbf{E} / \partial t^2 = 12\pi k_e c^2 q / |\tilde{\mathbf{r}}|^4 \quad \Rightarrow \quad \nabla^2 \mathbf{E} = (1/c^2) \partial^2 \mathbf{E} / \partial t^2. \quad (97)$$

$$\nabla^2 \mathbf{E} = (1/c^2) \partial^2 \mathbf{E} / \partial t^2 \quad \wedge \quad \varepsilon_0 \mu_0 = 1/c^2 \quad \Rightarrow \quad \nabla^2 \mathbf{E} = \varepsilon_0 \mu_0 \partial^2 \mathbf{E} / \partial t^2, \quad (98)$$

which is Maxwell's electric field wave equation. And from equation (91):

$$\partial \mathbf{B} / \partial t = -4\pi k_e q / r^3 \quad \wedge \quad r = |\tilde{\mathbf{r}}| \quad \Rightarrow \quad \nabla^2 \mathbf{B} = 12\pi k_e q / |\tilde{\mathbf{r}}|^4. \quad (99)$$

$$\mathbf{B} = 2\pi k_e q / |\tilde{\mathbf{r}}|^2 \quad \wedge \quad r = ct \quad \Rightarrow \quad \mathbf{B} = 2\pi k_e q / (ct)^2 \quad \Rightarrow \quad \partial^2 \mathbf{B} / \partial t^2 = (12\pi k_e / c^2) q / t^4. \quad (100)$$

$$\partial^2 \mathbf{B} / \partial t^2 = (12\pi k_e / c^2) q / t^4 \quad \wedge \quad r = ct \quad \Rightarrow \quad \partial^2 \mathbf{B} / \partial t^2 = 12\pi k_e c^2 q / |\tilde{\mathbf{r}}|^4. \quad (101)$$

$$\nabla^2 \mathbf{B} = -12\pi k_e q / |\tilde{\mathbf{r}}|^4 \quad \wedge \quad \partial^2 \mathbf{B} / \partial t^2 = -12\pi k_e c^2 q / |\tilde{\mathbf{r}}|^4 \quad \Rightarrow \quad \nabla^2 \mathbf{B} = (1/c^2) \partial^2 \mathbf{B} / \partial t^2. \quad (102)$$

$$\nabla^2 \mathbf{B} = (1/c^2) \partial^2 \mathbf{B} / \partial t^2 \quad \wedge \quad \varepsilon_0 \mu_0 = 1/c^2 \quad \Rightarrow \quad \nabla^2 \mathbf{B} = \varepsilon_0 \mu_0 \partial^2 \mathbf{B} / \partial t^2, \quad (103)$$

which is Maxwell's magnetic field wave equation.

### The fine structure constant, $\alpha$

The ratio of two subtypes of force implies ratios of the form:

$$\alpha_\tau = \frac{F_{\tau_1}}{F_{\tau_2}} = \frac{K\tau_1^2/r^2}{K\tau_2^2/r^2} = \frac{\tau_1^2}{\tau_2^2}. \quad (104)$$

For example, the fine structure electron coupling constant is the ratios of two types of Coulomb's charge force, which simplifies to:

$$\alpha = q_e^2/q_p^2, \quad (105)$$

where  $q_e$  is the elementary (electron) charge and  $q_p$  is the reduced Planck charge unit.

The gravitational fine structure constant,  $\alpha_G$ , is the ratio of two subtypes of Newton's gravity force, which simplifies to:

$$\alpha_G = m_e^2/m_p^2, \quad (106)$$

where  $m_e$  is the electron mass and  $m_p$  is the reduced Planck mass unit.

### $\hbar$ , $h$ , and Einstein-Planck relation

Applying both the direct proportion ratio (117), and inverse proportion ratio (52):

$$m = k_m/r \quad \wedge \quad r = ct \quad \Rightarrow \quad m(ct)^2 = (k_m/r)r^2 = k_mr. \quad (107)$$

$$m(ct)^2 = k_mr \quad \Rightarrow \quad E \stackrel{\text{def}}{=} mc^2 = k_mr/t^2. \quad (108)$$

$$E = mc^2 = k_mr/t^2 \quad \wedge \quad r/t = c \quad \Rightarrow \quad E = mc^2 = k_mc/t. \quad (109)$$

The next 4 steps show that  $k_mc = \hbar$ , the *reduced* Planck constant. From the fine structure equation 106:

$$\begin{aligned} \alpha = q_e^2/q_p^2 \quad \wedge \quad q_e = 1.60217663 \cdot 10^{-19} \text{ C} \quad \wedge \quad \alpha \approx 7.2973525705 \cdot 10^{-3} \\ \Rightarrow \quad q_p = \sqrt{q_e^2/\alpha} \approx 1.87554604 \cdot 10^{-18} \text{ C}. \end{aligned} \quad (110)$$

From equations 88 and 110:

$$\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m = 4\pi(r_p/q_p)^2/(r_p/m_p) = 4\pi m_p r_p/q_p^2 = 4\pi k_m/q_p^2. \quad (111)$$

$$\begin{aligned} \mu_0 = 4\pi k_m/q_p^2 \quad \wedge \quad q_p \approx 1.87554604 \cdot 10^{-18} \text{ C} \quad \wedge \quad \mu_0 \approx 1.25663706127(20) \cdot 10^{-6} \text{ kg m C}^{-2} \\ \Rightarrow \quad k_m = \mu_0 q_p^2/4\pi \approx 3.51767294 \cdot 10^{-43} \text{ kg m}. \end{aligned} \quad (112)$$

$$\begin{aligned} k_m \approx 3.51767294 \cdot 10^{-43} \text{ kg m} \quad \wedge \quad c = 2.99792458 \cdot 10^8 \text{ m s}^{-1} \\ \Rightarrow \quad \hbar = k_mc \approx 1.054571817 \cdot 10^{-34} \text{ kg m}^2 \text{ s}^{-1}. \end{aligned} \quad (113)$$

Combining equations 109 and 113 yields  $\hbar$ ,  $h$ . and the Einstein-Planck relation[13, 17]:

$$E = mc^2 = k_m c/t \text{ kg } m^2 s^{-2} \wedge k_m c = \hbar \text{ kg } m^2 s^{-1} \Rightarrow E = \hbar/t \text{ kg } m^2 s^{-2}. \quad (114)$$

$$E = \hbar/t \text{ kg } m^2 s^{-2} \wedge \omega = (2\pi/t) \text{ radian } s^{-1} \Rightarrow \\ E_\omega = \hbar(2\pi/t) = \hbar\omega \text{ kg } m^2 \text{ radian } s^{-2}. \quad (115)$$

$$E_\omega = \hbar/t \text{ kg } m^2 \text{ radian } s^{-2} \wedge \\ f = (1/t) \text{ cycle } s^{-1} = ((2\pi/t) \text{ radian } s^{-1})/(2\pi \text{ radian cycle}^{-1}) \\ \Rightarrow E_f = \hbar\omega(2\pi/2\pi) = 2\pi\hbar f = hf \text{ kg } m^2 \text{ cycle } s^{-2} \stackrel{\text{def}}{=} hf \text{ kg } m^2 \text{ Hz } s^{-1}. \quad (116)$$

### Values of the unit ratios and Planck units

The direct proportion ratios are:

$$c_t = c \approx 2.99792458 \cdot 10^8 \text{ m } s^{-1}. \quad (117)$$

$$G = c_m c_t^2 \wedge G \approx 6.67430(15) \cdot 10^{-11} \text{ m}^3 \text{ kg}^{-1} s^{-2} \Rightarrow \\ c_m = G/c_t^2 \approx 7.426162 \cdot 10^{-28} \text{ m } \text{kg}^{-1}. \quad (118)$$

$$k_e = c_q^2 c_t^2 / c_m \wedge k_e \approx 8.9875517923(13) \cdot 10^9 \text{ Nm}^2 \text{ C}^{-2} \Rightarrow \\ c_q = \sqrt{k_e c_m / c_t^2} \approx 8.6175182 \cdot 10^{-18} \text{ m } \text{C}^{-1}. \quad (119)$$

From equation 113, the inverse proportion ratio values are:

$$\hbar \approx 1.054571783 \cdot 10^{-34} \text{ kg } m^2/s \wedge c_t = 2.99792458 \cdot 10^8 \text{ m/s} \\ \Rightarrow k_m = \hbar/c_t \approx 3.51767283 \cdot 10^{-43} \text{ kg } m. \quad (120)$$

$$r_p = \sqrt{r_p^2} = \sqrt{k_m c_m} \approx 1.616255 \cdot 10^{-35} \text{ m}. \quad (121)$$

$$k_t = r_p/c_t \approx 8.71363 \cdot 10^{-79} \text{ s } m. \quad (122)$$

$$k_q = r_p/c_q \approx 3.031361 \cdot 10^{-53} \text{ C } m. \quad (123)$$

From the fine structure equation 110, the reduced Planck unit values are:

$$q_p = \sqrt{q_e^2/\alpha} \approx 1.87554604 \cdot 10^{-18} \text{ C}. \quad (124)$$

$$r_p = \sqrt{r_p^2} = \sqrt{c_t k_t} = \sqrt{c_m k_m} = \sqrt{c_q k_q} \approx 1.616255 \cdot 10^{-35} \text{ m}. \quad (125)$$

$$t_p = r_p/c_t \approx 5.391246 \cdot 10^{-44} \text{ s}. \quad (126)$$

$$m_p = r_p/c_m \approx 2.176434 \cdot 10^{-8} \text{ kg}. \quad (127)$$

### Compton wavelength, $\lambda$

[13, 17] From equations 54 and 110:

$$\lambda = 2\pi r = 2\pi k_m/m = 2\pi k_m c/mc \quad \wedge \quad h = 2\pi \hbar = 2\pi k_m c \quad \Rightarrow \quad \lambda = h/mc. \quad (128)$$

### Schrödinger's wave equation

Start with the previous unit ratio-derived “base” Einstein-Planck relation 114 and multiply the kinetic energy component by  $mc/mc$ :

$$mc^2 = \hbar/t \quad \Rightarrow \quad \exists V(r, t) : \hbar/t = \hbar/2t + V(r, t). \quad (129)$$

$$\hbar/t = \hbar/2t + V(r, t) \quad \Rightarrow \quad \hbar/t = \hbar mc/2mct + V(r, t). \quad (130)$$

And from the distance-to-time (speed of light) ratio (117):

$$\hbar/t = \hbar mc/2mct + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar mc^2/2mcr + V(r, t). \quad (131)$$

$$\hbar/t = \hbar mc^2/2mcr + V(r, t) \quad \wedge \quad \hbar/t = mc^2 \quad \Rightarrow \quad \hbar/t = \hbar^2/2mcrt + V(r, t). \quad (132)$$

$$\hbar/t = \hbar^2/2mcrt + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar^2/2mr^2 + V(r, t). \quad (133)$$

Multiply both sides of equation 133 by a probability density function,  $\Psi(r, t)$ .

$$\hbar/t = \hbar^2/2mr^2 + V(r, t) \quad \Rightarrow \quad (\hbar/t)\Psi(r, t) = (\hbar^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t). \quad (134)$$

$$(\hbar/t)\Psi(r, t) = ((\hbar)^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t) \quad \wedge$$

$$\forall \Psi(r, t) : \partial^2 \Psi(r, t)/\partial r^2 = (-1/r^2)\Psi(r, t) \quad \wedge \quad \partial \Psi(r, t)/\partial t = (i 1/t)\Psi(r, t)$$

$$\Rightarrow \quad i\hbar \partial \Psi(r, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t), \quad (135)$$

which is the one-dimensional position-space Schrödinger's equation [4, 17].

$$i\hbar \partial \Psi(r, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t) \quad \wedge \quad ||\vec{r}|| = r$$

$$\Rightarrow \quad \exists \vec{r} : i\hbar \partial \Psi(\vec{r}, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(\vec{r}, t)/\partial \vec{r}^2 + V(\vec{r}, t)\Psi(\vec{r}, t), \quad (136)$$

which is the 3-dimensional position-space Schrödinger's equation [4, 17].

## Dirac's wave equation

Start with the previous unit ratio-derived “base” Einstein-Planck relation 114:

$$\hbar/t = mc^2 \Leftrightarrow \hbar/2t + \hbar/2t. = mc^2 \Leftrightarrow \hbar/2t = -\hbar/2t. + mc^2. \quad (137)$$

$$\hbar/2t = -\hbar/2t. + mc^2 \wedge qc_q/r = qc_q/ct = 1 \Rightarrow \hbar qc_q/2ct^2 = -qc_q/2rt. + mc^2. \quad (138)$$

$$\hbar qc_q/2ct^2 = -qc_q/2rt. + mc^2 \wedge r = ct \Rightarrow \hbar qc_q/2ct^2 = -qcc_q/2r^2. + mc^2. \quad (139)$$

$$\hbar qc_q/2ct^2 = -qcc_q/2r^2. + mc^2 \Rightarrow \hbar(qc_q/2ct^2)\psi = (-qcc_q/2r^2)\psi + mc^2\psi. \quad (140)$$

$$\psi = e^{-iqc_q((1/2ct)-(1/2r))} \Rightarrow \partial\psi/\partial t = (-iqc_q/2ct^2)\psi \wedge \nabla\psi = (iqc_q/2r^2)\psi. \quad (141)$$

$$\begin{aligned} \partial\psi/\partial t &= (-iqc_q/2ct^2)\psi \wedge \nabla\psi = (iqc_q/2r^2)\psi \wedge \\ \hbar(qc_q/2ct^2)\psi &= (-qcc_q/2r^2)\psi + mc^2\psi \Rightarrow i\hbar\partial\psi/\partial t = -i\hbar c\nabla\psi + mc^2\psi. \end{aligned} \quad (142)$$

$$\begin{aligned} \forall y = a\mathbf{x} + b \exists \boldsymbol{\alpha}, \beta : y &= \boldsymbol{\alpha} \cdot a\mathbf{x} + b\beta \wedge i\hbar\partial\psi/\partial t = -i\hbar c\nabla\psi + mc^2\psi \\ \Rightarrow \exists \boldsymbol{\alpha}, \beta : i\hbar\partial\psi/\partial t &= i\hbar c\boldsymbol{\alpha} \cdot -\nabla\psi + mc^2\beta\psi \end{aligned} \quad (143)$$

which is the relativistic, free particle Dirac equation[5], where  $\beta = \gamma^0$ ,  $\boldsymbol{\alpha} = \gamma^0\boldsymbol{\gamma}^k$ ,  $k \in \{1, 2, 3\}$ , and each  $\gamma$  a matrix.

## Total of a type

Applying both the direct (117) and inverse proportion ratios (120) to the Euclidean distance:

$$\begin{aligned} r &= \sqrt{r_1^2 + r_2^2}, \quad r_1 = c_\tau\tau, \quad r_2 = k_\tau/\tau, \quad \tau \in \{t, m, q\} \\ \Rightarrow r &= \sqrt{(c_\tau\tau)^2 + (k_\tau/\tau)^2} \Leftrightarrow \tau = \sqrt{(r/c_\tau)^2 + (k_\tau/r)^2}. \end{aligned} \quad (144)$$

## Quantum extension to Schwarzschild's metric

The simplest way to demonstrate how to add quantum physics to general relativity is by extending Schwarzschild's gravitational time dilation equation and black hole metric. Apply the total of a type equation 144 to the derivation of Schwarzschild's time dilation and metric



(65):

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - (v^2/c^2)(r/r)} \quad \wedge \quad r = \sqrt{(c_m m)^2 + (k_m/m)^2} = Q_m \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2 / r c^2}.\end{aligned}\tag{145}$$

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2 / r c^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2 \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2 / r c^2}.\end{aligned}\tag{146}$$

For a photon, the escape velocity,  $v_{escape} = c$ .

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2 / r c^2} \quad \wedge \quad v_{escape} = c \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m c^2 / r c^2} = \sqrt{1 - 2Q_m / r}.\end{aligned}\tag{147}$$

Combining equation 147 with equation 62 yields Schwarzschild's gravitational time dilation with a quantum mass effect:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m / r} \quad \wedge \quad t' = t \sqrt{1 - (v^2/c^2)} \quad \Rightarrow \quad t' = t \sqrt{1 - 2Q_m / r}.\tag{148}$$

Schwarzschild defined the black hole event horizon radius,  $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$ . The radius with the quantum extension is  $\alpha \stackrel{\text{def}}{=} 2Q_m$ . At this point the exact same equations 70 through 73 yield what looks like the same Schwarzschild black hole metric.

### Quantum extension to Newton's gravity force

The quantum mass effect is easier to understand in the context Newton's gravity equation than in general relativity, because the metric equations and solutions in the EFEs are much more complex. From equations 144 and 50:

$$\begin{aligned}m / \sqrt{(r/c_m)^2 + (k_m/r)^2} &= 1 \quad \wedge \quad r^2 / (ct)^2 = 1 \quad \Rightarrow \quad r^2 / (ct)^2 = m / \sqrt{(r/c_m)^2 + (k_m/r)^2} \\ \Rightarrow \quad r^2 / t^2 &= c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2}.\end{aligned}\tag{149}$$

$$\begin{aligned}r^2 / t^2 = c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2} &\Rightarrow (m/r)(r^2 / t^2) = (m/r)(c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2}) \\ \Rightarrow \quad F \stackrel{\text{def}}{=} mr / t^2 &= c^2 m^2 / \sqrt{(r^4/c_m^2) + k_m^2}.\end{aligned}\tag{150}$$

$$\begin{aligned}F = c^2 m^2 / \sqrt{(r^4/c_m^2) + k_m^2} &\quad \wedge \quad \forall m \in \mathbb{R}, \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \\ \Rightarrow \quad F &= c^2 m_1 m_2 / \sqrt{(r^4/c_m^2) + k_m^2}.\end{aligned}\tag{151}$$

### Quantum extension to Coulomb's charge force

$$\begin{aligned} q^2/((r/c_q)^2 + (k_q/r)^2) = 1 \quad \wedge \quad r^2/(ct)^2 = 1 &\Rightarrow r^2/(ct)^2 = q^2/((r/c_q)^2 + (k_q/r)^2) \\ \Rightarrow r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2). \end{aligned} \quad (152)$$

$$\begin{aligned} r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2) \quad \wedge \quad r = c_m m \\ \Rightarrow F \stackrel{\text{def}}{=} mr/t^2 = (c^2/c_m) q^2/((r/c_q)^2 + (k_q/r)^2). \end{aligned} \quad (153)$$

$$\begin{aligned} \forall q \in \mathbb{R} : \exists q_1, q_2 \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad F = c^2/c_m q^2/((r/c_q)^2 + (k_q/r)^2) \\ \Rightarrow F = c^2/c_m q^2/((r/c_q)^2 + (k_q/r)^2). \end{aligned} \quad (154)$$

### INSIGHTS AND IMPLICATIONS

1. A distance measure,  $d$ , is an inverse (bijective) function of a volume,  $v$ , where  $v$  is the sum of subset volumes (Euclidean or non-Euclidean),  $g(s_{i,1}, \dots, s_{i,n})$ . Notionally:

$$\forall n \in \mathbb{N}, f, g : d \geq 0, d = f^{-1}(v) = f^{-1}(\sum_{i=1}^m g(s_{i,1}, \dots, s_{i,n})) : \quad (155)$$

- (a) shows the intimate relation between distance and volume that definitions, like inner product space and metric space, ignore [6, 7];
- (b) is a more simple and concise definition of a distance measure, having the metric space properties [6, 7].

2. The left side of the distance sum inequality (.7),

$$(\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}, \quad (156)$$

differs from the left side of Minkowski's sum inequality [11]:

$$(\sum_{i=1}^m (a_i^n + b_i^n)^{\mathbf{n}})^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad (157)$$

- (a) The two inequalities are only the same where  $n = 1$ .
- (b) The distance sum inequality (.7) is a more fundamental inequality because the proof does not require the convexity and Hölder's inequality assumptions required to prove the Minkowski sum inequality [11].
- (c)  $\forall n > 1, v > 0$ : The distance sum inequality term,  $v_i^n = a_i^n + b_i^n$ :  $d = v^{1/n} = (\sum_{i=1}^m v_i^n)^{1/n}$ , is the Minkowski distance, which makes it directly related to geometry. But the Minkowski sum inequality term,  $d = v^{1/n} = (\sum_{i=1}^m ((v_i^n)^{\mathbf{n}}))^{1/n} = (\sum_{i=1}^m v_i^{\mathbf{n}^2})^{1/n}$ , is *not* a Minkowski distance.

- (d) The distance sum inequality might be applicable to machine learning.
3. *Combinatorics.* The number of n-tuples,  $v_c = \prod_{i=1}^n |x_i|$ , was proven to imply the Euclidean volume equation (.1). And the notion of distance was derived as the inverse function of the sum of the number of n-tuples (.4). The Euclidean equation was derived without relying on the geometric primitives and relations in Euclidean geometry [18, 19], axiomatic geometry [20–24], trigonometry [25, 26], calculus [25, 27, 28], and vector analysis [3].
  4. *Combinatorics.* The commutative property of the operations defining volume and distance equations limits a domain set to  $n \leq 3$  unordered elements (.16). Other elements must have different types (are elements of different sets).
    - (a) If there are more than 6 dimensions  $\subseteq \mathbb{R}$ , then the additional dimensions are ordered, where each dimension has an intrinsic property having a fixed position in strict totally ordered set.
    - (b) If each type of quantum state is an ISCL, then there are at most 3 states of the same type: 3 orientations per dimension of space (+1,0-1), 3 quark color charges, {red, green, blue}, 3 quark anti-color charges, and so on.
  5. A discrete (point) valued state has measure 0 (zero-length interval). The ratio of a time or distance interval length to zero is undefined, which is the reason quantum entangled (discrete) state values exist independent of (non-deterministic with respect to) distance (position), time, mass, and charge. Therefore, a state value in space and time is undetermined until observed, like Schrödinger’s poisoned cat being both alive and dead until the box is opened [4].
  6. For each unit interval length,  $r_p$ , of a 3-dimensional distance,  $r \subseteq \mathbb{R}$ , there are unit interval lengths of other types of dimensions, forming unit ratios (117):  $c_t = r_p/t_p$ ,  $c_m = r_p/m_p$ ,  $c_q = r_p/q_p \Leftrightarrow$  the inverse proportion ratios (52):  $k_t = r_p t_p$ ,  $k_m = r_p m_p$ ,  $k_t = r_p q_p$ , where  $r_p$ ,  $t_p$ ,  $m_p$ , and  $q_p$  are the reduced Planck units (124).
  7. Many empirical equation were derived, in this article, using the perspective of mass and charge as dimensions. But there is empirical evidence of mass and charge confined to a point or volume and wave-particles moving through space. Quanta and wave behavior

are the results of the combination of direct and inverse proportion ratios, for example,  $\hbar = k_m c_t$ . The combined direct and inverse ratios and special relativity might be a framework to reconcile the two perspectives and might reduce the pantheon of particles in the Standard Model to a few simple parts.

8. An empirical law *describes* the relations between the variables in an equation. Deriving empirical laws from the ratios *explains* the reasons for relations between the variables in an equation. Further, all the derivations of the physics equations from the ratios are much shorter and simpler than other derivations, which shows that the ratios are an important tool for physicists and engineers.
9. As shown in the subsection deriving the Schwarzschild's gravitational time dilation and black hole metric (65) [14, 15] using ratios illustrates a simple way of finding exact solutions to Einstein's field equations.
10.  $c_t$ ,  $c_m$ , and  $c_q$  are the *maximum* ratios:  $r = \sqrt{r_1^2 + r_2^2 + r_3^2} \Rightarrow r_1, r_2, r_3 \leq r \Rightarrow \forall \tau \in \{t, m, q\} : r_1/\tau, r_2/\tau, r_3/\tau \leq r/\tau = r_p/\tau_p = c_\tau$ . For example, the speeds of gravity and light waves are limited to the speed,  $c_t$ .
11.  $c_t$  is a component of the constants:  $G = c_m c_t^2$ ,  $k_e = c_q^2 c_t^2 / c_m$ ,  $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2 / c_m)$ , and  $\hbar = k_m c_t$ . The only constants, derived in this article, that do not contain  $c_t$  are: vacuum permeability constant:  $\mu_0 = 4\pi k_e / c_t^2 = 4\pi c_q^2 / c_m$ , and the fine structure constants,  $\alpha = q_e^2 / q_p^2$  and  $\alpha_G = m_e^2 / m_p^2$ .
12. The fine structure electron coupling constant,  $\alpha$  was derived from the ratio of two subtypes of charge force that reduces to the unit-less ratio  $\alpha = q_e^2 / q_p^2 \approx 0.0072973526$  (106), which is the empirical CODATA value [16].
  - (a) The CODATA electron coupling version of the fine structure constant,  $\alpha$  is defined as:  $\alpha = q_e^2 / 4\pi\varepsilon_0 \hbar c = q_e^2 / 2\varepsilon_0 \hbar c$  [16]. The following steps show that the CODATA definition reduces to the ratio-derived equation:

$$\begin{aligned}
\varepsilon_0 &= 1/4\pi k_e = 1/(4\pi(c_q^2 c_t^2 / c_m)) \quad \wedge \quad \hbar = k_m c_t \quad \wedge \quad h = 2\pi \hbar \\
&\Rightarrow \varepsilon_0 \hbar c = 2\pi k_m c_t^2 / (4\pi(c_q^2 / c_m) c_t^2) = k_m / (2(c_q^2 / c_m)) \\
&= m_p r_p / (2((r_p / q_p)^2 / (r_p / m_p))) = q_p^2 / 2. \quad (158)
\end{aligned}$$

$$\alpha = q_e^2/2\varepsilon_0\hbar c \quad \wedge \quad \varepsilon_0\hbar c = q_p^2/2 \quad \Rightarrow \quad \alpha = q_e^2/2(q_p^2/2) = q_e^2/q_p^2. \quad (159)$$

(b) The fine structure electron coupling constant is the ratio of electron static charge force to the propagating quantum charge (photon/electromagnetic) wave force, caused by a moving charged particle.

(c) The gravitational fine structure electron coupling constant,  $\alpha_G$ , can be expressed as the ratio of two subtypes of Newton's gravity force, where  $m_e$  is the rest electron mass and  $m_p$  is the reduced Planck mass unit:  $\alpha_G = m_e^2/m_p^2$  (106).

The currently used gravitational fine structure constant reduces to the equation derived in this article:  $\alpha_G = Gm_e^2/\hbar c = (c_m c_t^2)m_e^2/(k_m c_t)c_t = c_m m_e^2/k_m = (r_p/m_p)m_e^2/m_p r_p = m_e^2/m_p^2$ .

13. The assumption that the old definition,  $\mu_0 = 4\pi k_e/c_t^2 = 1/\varepsilon_0 c^2$ , and the newer NIST/CODATA definition,  $\mu_0 = 2h\alpha/cq_e^2$ [16], are different because of slightly different values is *incorrect*! The following steps will show that both equations reduce to the same base equation,  $\mu_0 = 4\pi k_m/q_p^2$ , which shows that the old and newer NIST/CODATA definitions differ in value because of different measurement methods.

Using the ratios to reduce the old definition of  $\mu_0$  derived in this article:

$$\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m = 4\pi(r_p/q_p)^2/(r_p/m_p) = 4\pi m_p r_p/q_p^2 = 4\pi k_m/q_p^2. \quad (160)$$

Using the ratios to reduce the CODATA definition:

$$h = 2\pi k_m c_t, \quad \alpha = q_e^q/q_p^2 \Rightarrow \mu_0 = 2h\alpha/cq_e^2 = 4\pi k_m c_t(q_e^2/q_p^2)/c_t q_e^2 = 4\pi k_m/q_p^2. \quad (161)$$

14. Empirical and hypothesized laws of physics use an *opaque* constant to make the units in an equation balance. The opacity has led to the *incorrect* assumptions of those constants being fundamental (atomic) constants.

In this article, some opaque constants are derived directly from (composed of) the ratios: gravity,  $G = c_m c_t^2$  (57), charge,  $k_e = c_q^2 c_t^2/c_m$  (77), reduced Planck  $\hbar = k_m c_t$  (110),  $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2/c_m)$  (81), and  $\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m$  (87).

And the quantum extensions to: Schwarzschild's gravitational time dilation (147), Newton's gravity force (151), and Coulomb's charge force (154), show that, where the

quantum effects become measurable, the constants  $G$ ,  $k_e$ ,  $\varepsilon_0$ , and  $\mu_0$  no longer exist (are no longer valid).

Therefore,  $G$ ,  $k_e$ ,  $\varepsilon_0$ ,  $\mu_0$ , and  $\hbar$  are *not* fundamental constants.

15. Constants that use notions of temperature and pressure for example, the Boltzmann constant [16], might not be definable solely in terms of unit ratios.
16. The derivations of:  $\nabla \cdot \mathbf{g} = -4\pi G\rho$  from  $\mathbf{g} = 2\pi Gm/|\tilde{\mathbf{r}}|^2$  (58), and  $\nabla \cdot \mathbf{E} = -\rho/\varepsilon_0$  from  $\mathbf{E} = 2\pi k_e q/|\tilde{\mathbf{r}}|^2$  (81), show that:

- (a) Mass and charge density,  $\rho$ , as motivations are unnecessary complexities. Equation 55 used to derive Newton's law and equation 74 used to derive Coulomb's law relate linear acceleration,  $r/t^2$  to mass and charge. It is small step to recognize Gauss's  $\mathbf{g}$ , Poisson's  $-\nabla\Phi(\tilde{\mathbf{r}}, t)$ , and Gauss's  $\mathbf{E}$  relate the angular (orbital) acceleration,  $2\pi r/t^2$ , to mass and charge, where differentiation yields the gravity field and electric field laws.

The derivations are much simpler and shorter using angular (orbital) acceleration,  $2\pi r/t^2$ . The orbit length,  $2\pi r$ , corresponds to the Gauss's line integral length of a curve circumscribing a sphere containing the mass or charge.

- (b) Mass and charge density obfuscates the pattern,  $\nabla \cdot f(x, r) = -4\pi k_x x/|\tilde{\mathbf{r}}|^3$ , being derived from the inverse square pattern,  $f(x, r) = 2\pi k_x x/|\tilde{\mathbf{r}}|^2$ . The constant,  $G$  (derived from the inverse square relation (55)), is used in Einstein's constant,  $\kappa = 8\pi G/c^4$ , and the use of energy density in the stress energy tensor,  $T_{\mu,\nu}$ , in Einstein's field equations [3] also obfuscate the inverse square pattern.

17. The ratio-based derivation of Lorentz's law here (85) differs from other relativistic derivations, in that equations for the magnetic field,  $\mathbf{B}$ , and magnetic permeability,  $\mu_0$ , are derived in the process. The equation,  $\mathbf{B} = (2\pi k_e/c^2)vq/|\tilde{\mathbf{r}}|^2$ , allows a simple derivation of the Faraday's law (93) and the magnetic component of Maxwell's electromagnetic equations (103).
18. Maxwell's derivation of the electric and magnetic field wave equations required: Gauss's electric field law in a vacuum:  $\nabla \cdot \mathbf{E} = -\rho/\varepsilon_0$ , Gauss's magnetic field law in a vacuum:  $\nabla \cdot \mathbf{B} = 0$ , Faraday's Law:  $\nabla \times \mathbf{E} = -\partial\mathbf{B}/\partial t$ , and Ampere's Law:

$\nabla \times \mathbf{B} = \mu_0 \varepsilon_0 \partial \mathbf{E} / \partial t$  [13]. In contrast, the derivations (98) (103), in this article, did not require any of those laws and only required the ratio  $r = ct$  and the ratio-derived definitions:  $\mathbf{E}$ ,  $\mathbf{B}$ ,  $\varepsilon_0$ , and  $\mu_0$ . And the derivations were shorter and simpler.

19. Einstein's relativity equations: 1) assume the Lorentz transformations, 2) assume the laws of physics are same at each coordinate point, 3) assume the notion of light, and 4) assume that the speed of light is the same at each coordinate point [2, 3].

The derivations, in this article, were made without those assumptions (does not even require the notion of light). The unit ratios are ratios of Euclidean distance to time, mass, and charge. Therefore, the unit ratios (like the maximum speed ratio,  $c_t$ ), and all equations (laws) and constants derived from those ratios must be the same at each Euclidean-like coordinate point.

20. Using the quantum units,  $r_p$  and  $t_p$ :  $r_p/t_p^2$ , suggests a maximum linear acceleration for masses. And  $2\pi r_p/t_p^2$  suggests a maximum orbital or rotational acceleration.
21. The Einstein-Planck relations,  $E_\omega = \hbar\omega \text{ kg } m^2 \text{ radian } s^{-2}$  (115) and  $E_f = hf \text{ kg } m^2 s^{-1} Hz$  (116), are both instances of the ratio-derived base Einstein-Planck relation,  $E = \hbar/t \text{ kg } m^2 s^{-2}$  (114). And derivations of Schrödinger's wave equation (136) [4] and Dirac's wave equation (143) [5] from the ratio-derived base Einstein-Planck relation are shorter, simpler, and require fewer assumptions.
22. The quantum extensions to: Schwarzschild's black hole metric (73), Newton's gravity force (151), and Coulomb's charge force (154) make quantifiable predictions:

- (a) For gravity,  $\lim_{r \rightarrow 0} F = c^2 m_1 m_2 / k_m$ , and for charge,  $\lim_{r \rightarrow 0} F = 0$ . Finite maximum gravity and charge forces: 1) eliminates the problem of forces going to infinity as  $r \rightarrow 0$ , 2) might allow radioactivity without the need for a weak force, and 3) finite sloped energy well walls might explain quantum tunneling.
- (b) The quantum-extended relativity and classic equations reduce to the non-extended relativity and classic equations and constants, where the distance between 2 masses and between 2 charges is sufficiently large or the masses and charges sufficiently large that the quantum effects are not measurable. *Note* that

$G$ ,  $k_e$ ,  $\varepsilon_0$ ,  $\mu_0$ , and  $\kappa$  (Einstein's constant, which contains  $G$ ) do not exist (are not valid), where the quantum effects becomes measurable.

- (c) The covariant tensor components, in Einstein's field equations, that had the units  $1/\text{distance}^2$ , will now have the more complex units,  $1/\sqrt{(\text{distance}^4/c_\tau^2) + k_\tau^2}$ ,  $\tau \in \{t, m\}$ .
- (d)  $1/\sqrt{(\text{distance}^4/c_\tau^2) + k_\tau^2}$  implies that as  $\text{distance} \rightarrow 0$ , spacetime curvature peaks at the reduced Planck unit distance,  $r_p$  (124). The ratio of quantum units,  $m_p/r_p^3$ , might indicate a maximum mass density. A finite force (spacetime curvature) and finite mass density might imply that black holes have sizes  $> 0$  (are not singularities). It might allow Black hole evaporation. If there was a "big bang," then it might not have originated from a singularity.

- 
- [1] Copyright © 2026 George M. Van Treeck. All rights reserved.
  - [2] A. Einstein, *Relativity, The Special and General Theory* (Princeton University Press, 2015).
  - [3] H. Weyl, *Space-Time-Matter* (Dover Publications Inc, 1952).
  - [4] E. Schrödinger, *Collected Papers On Wave Mechanics* (Minkowski Institute Press, 2020).
  - [5] P. Dirac, *The Principles of Quantum Mechanics* (Snowball Publishing, 1957).
  - [6] R. R. Goldberg, *Methods of Real Analysis* (John Wiley and Sons, 1976).
  - [7] W. Rudin, *Principles of Mathematical Analysis* (McGraw Hill Education, 1976).
  - [8] W. Conradie and V. Goranko, *Logic and Discrete Mathematics* (Wiley, 2015).
  - [9] X. P. N. Guigui, N. Miolane, Introduction to riemannian geometry and geometric statistics: from basic theory to implementation with geomstats, *Foundations and Trends in Machine Learning* **16**, 182 (2023), <https://inria.hal.science/hal-03766900v2>, hal-00644459.
  - [10] Rocq, Rocq proof assistant (2025), <https://rocq-prover.org/docs>.
  - [11] H. Minkowski, *Geometrie der Zahlen* (Chelsea, 1953) reprint.
  - [12] I. Newton, *The Mathematical Principles of Natural Philosophy* (Patristic Publishing, 2019) reprint.
  - [13] R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics* (California Institute of Technology, 2010) <https://www.feynmanlectures.caltech.edu/>.



- [14] K. Schwarzschild, Über das gravitationsfeld eines massenpunktes nach der einsteinschen theorie, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften **7**, 198 (1916), <https://archive.org/stream/sitzungsberichte1916deutsch/page/188/mode/2up>.
- [15] S. Antoci and A. Loinger, An english translation of k. schwarzschild's 1916 article: On the gravitational field of a mass point according to einstein's theory, ARXIV.org (1999), <https://arxiv.org/abs/physics/9905030>.
- [16] CODATA, Physical constant values (2022), <https://physics.nist.gov/cuu/Constants/index.html>.
- [17] M. Jain, *Quantum Mechanics: A Textbook for Undergraduates* (PHI Learning Private Limited, New Delhi, India, 2011).
- [18] D. E. Joyce, Euclid's elements (1998), <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>.
- [19] E. S. Loomis, *The Pythagorean Proposition* (NCTM, 1968).
- [20] G. D. Birkhoff, A set of postulates for plane geometry (based on scale and protractors), Annals of Mathematics **33** (1932).
- [21] D. Hilbert, *The Foundations of Geometry (2cd Ed)* (Chicago: Open Court, 1980) <http://www.gutenberg.org/ebooks/17384>.
- [22] J. M. Lee, *Axiomatic Geometry* (American Mathematical Society, 2010).
- [23] O. Veblen, A system of axioms for geometry, Trans. Amer. Math. Soc **5**, 343 (1904), <https://doi.org/10.1090/S0002-9947-1904-1500678-X>.
- [24] A. Tarski and S. Givant, Tarski's system of geometry, The Bulletin of Symbolic Logic **5**, 175 (1999).
- [25] A. Bogomolny, *Pythagorean Theorem* (Interactive Mathematics Miscellany and Puzzles, 2010) <http://www.cut-the-knot.org/pythagoras/CalculusProofCorrectedVersion.shtml>.
- [26] J. Zimba, On the possibility of trigonometric proofs of the pythagorean theorem, Forum Geometricorum **9**, 275 (2009).
- [27] B. C. Berndt, Ramanujan-100 years old (fashioned) or 100 years new (fangled)?, The Mathematical Intelligencer **10** (1988).
- [28] M. Staring, The pythagorean proposition a proof by means of calculus, Mathematics Magazine **69**, 45 (1996).