

Some set and number properties underlying physics

George M. Van Treeck ORCID: 0000-0002-1729-2615

(Dated: 22 July 2026)

Euclidean volume and some distance equations are proved to be derived from ordered combinations (n-tuples) of elements from a list of countable sets. A linearly sequenceable property is proved to constrain a set of commutable elements to at most 3 elements: for example, a set of 3 distance values, {width, height, depth}, and a set of 3 non-distance values, {time, mass, charge}. The math proofs are verified in Rocq. Dividing Euclidean distance, time, mass, and charge into n number of constant-sized intervals, for i in 1 to n , there are 3 direct and 3 inverse constant proportion ratios of the i -th Euclidean distance constant interval length to the i -th constant interval lengths of time, mass, and charge. Shorter and simpler derivations of well-known gravity, charge, electromagnetic, relativity, quantum physics equations and related constants from the ratios are presented. The derivations eliminate the need for dimensional analysis. The inverse proportion ratios are used to add quantum extensions to the Schwarzschild metric, Newton's gravity force, and Coulomb's charge force, which predict that: 1) The gravity and charge constants do not exist where the quantum effects are measurable; 2) The forces between two point masses and between two point charges peak at the Planck unit length.

CONTENTS

I. Introduction	3
A. Insufficient reduction	3
B. First reduction step	4
C. Second reduction step	5
D. Third reduction step	6
II. Volume	6
A. Euclidean volume	6
III. Distance	7
A. Vector inner product	7
B. Minkowski distance (p -norm)	8
C. Distance inequalities	8
D. Metric Space	9
IV. Properties limiting a set to at most 3 elements	10
V. Derivation of physics equations and constants	12
A. 3 direct and 3 inverse proportion ratios	12
B. G , Newton, Gauss, Poisson gravity laws	13
C. Special relativity	14
D. Analytic general relativity solutions	14
E. Schwarzschild's time dilation and metric	15
F. k_e , ε_0 , E , Coulomb and Gauss laws	16
G. B , μ_0 , and Lorentz's law	17
H. Faraday's law	18
I. Maxwell's wave equations	19
J. The classic electron radius, r_e	19
K. The fine structure constants, α and α_G	20
L. The Rydberg constant, R_∞	20

M. $\hbar$, h , and Planck-Einstein relation	20
N. Values of the unit ratios and Planck units	21
O. Compton wavelength, λ	22
P. Schrödinger's wave equation	22
Q. Dirac's wave equation	23
R. Total of a type	23
S. Quantum extension to Schwarzschild's metric	23
T. Quantum extension to Newton's gravity force	24
U. Quantum extension to Coulomb's charge force	25
VI. Insights and implications	25
References	31

Keywords: mathematical physics, gravity, charge, electromagnetism, relativity, quantum.

I. INTRODUCTION

A. Insufficient reduction

The philosophy of reductionism posits that complex phenomena can be reduced to simpler objects and relations. But, the Einstein field equations (general theory of relativity) and Schrödinger's wave equation (quantum physics) are examples of complicated equations with difficulty in finding exact (analytic) solutions.²⁻⁴ And relativity and quantum theory make assumptions without mathematical motivation. For example both assume 3 dimensions of space. But integration, Hilbert spaces, linear algebra, (pseudo-)Riemann manifolds, and so on, have no restriction on the number of dimensions.^{3,5-8}

Further, many physics equations rely on volume or distance. But mathematical analysis defines Euclidean volume as the product of interval lengths $\subseteq \mathbb{R}^n$ and defines the Euclidean distance equation, dot product, inner product, and metric space.^{5,6} Justifications for the definitions use rigorous finger-pointing to Euclidean geometry, Cartesian geometry, and "It's an abstraction."

The first reduction step, in this article, proves that well-known volume and distance equations are reduced to (derived from) a few simple set and number properties. The

second step identifies three properties of the sets and corresponding volume and distance equations salient to physics. The third reduction step shows that many well-known physics equations, constants, and assumptions are reduced to (derived from) two consequences of those three properties.

B. First reduction step

If the underlying principles are simple, then the math proofs should also be simple, trivial even. But even apparently trivial proofs can have subtle flaws.

Therefore, each math proof, in this article, has a corresponding formal proof, which has been verified using the Rocq proof verification system,⁹ a verification tool used by mathematicians world-wide. The formal proofs are in the Rocq files, “euclidrelations.v” and “threed.v,” which are included as ancillary files.

Let $|x_i|$ be the cardinal of (number of elements in) the abstract, countable set, $x_i \in \{x_1, \dots, x_n\}$. And let the value, v_c , be the number of ordered combinations (n-tuples) of the elements of $x_1, \dots, x_n$. Notionally:

$$x_i \in \{x_1, \dots, x_n\}, \quad \bigcap_{i=1}^n x_i = \emptyset, \quad |x_i| \in \{0, \mathbb{N}\}, \quad v_c = \prod_{i=1}^n |x_i|. \quad (1)$$

The Euclidean volume equation will be proved to be the trivial case where each n-tuple corresponds to a scalar value, $\kappa \in \mathbb{R}$ and v_c is the total number of n-tuples:

$$v_c \kappa = \left(\prod_{i=1}^n |x_i|\right) \kappa \quad \Rightarrow \quad \exists v, s_i \in \mathbb{R} : v = \prod_{i=1}^n s_i. \quad (2)$$

For all $n > 1$, there are cases where different domain values, $|x_1|, \dots, |x_n|$, multiplied yield the same range value, v_c . Inferring a domain value, d_c , from v_c , requires an inverse (bijective) function, $d_c = f_n^{-1}(v_c)$ and $v_c = f_n(d_c)$. The trivial bijective case for all n is the cuboid case, $|x_i| = d_c$. And, where v_c is the sum of the cardinals of subset n-tuples, it will be proved that:

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} \quad \Rightarrow \quad d^n = v = \sum_{i=1}^m \left(\prod_{j=1}^n s_{i,j}\right). \quad (3)$$

The $n = 2$ case is the inner product, where d is the norm:⁶

$$d^2 = \sum_{i=1}^m \left(\prod_{j=1}^{n=2} s_{i,j}\right) \wedge \exists a_i, b_i \in \mathbb{R} : s_{i,1} = a_i, s_{i,2} = b_i \Rightarrow d^2 = \sum_{i=1}^m a_i b_i \stackrel{\text{def}}{=} \mathbf{a} \cdot \mathbf{b}. \quad (4)$$

And, where each summed volume, v_{c_i} , is also the bijective function, $d_{c_i}^n = v_{c_i}$:

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m d_{c_i}^n \Rightarrow d^n = \sum_{i=1}^m d_i^n. \quad (5)$$

$|d|$ is the p -norm (Minkowski distance),¹⁰ which will be proved to imply the metric space properties.^{5,6} The $n = 2$ case is the Euclidean distance (flat distance metric).^{11,12} Multiplying each summed area by a scalar function, $\alpha_i : d^2 = \sum_{i=1}^m (\alpha_i s_i^2)$, yields a Gaussian-like curved distance metric that motivated the metric, $g_{\mu,\nu}$, in Einstein's general relativity theory.²

C. Second reduction step

One property salient to physics: All distances derived here are bijective functions of a volume, $v \in \mathbb{R}$, which implies the dimension of each type of distance (for example, Euclidean distance) is always a $\subseteq \mathbb{R}$. *Note*: Euclidean distance is *not* a bijective function of the volume domain values (width, height, and depth).

A second property salient to physics: All the set elements, $x_1, \dots, x_n$ and volume domain values, $s_1, \dots, s_n$, are linearly sequenced via sequence functions, for example, sequenced via the successor and predecessor functions, $successor(x_i) = x_{i+1}$ and $predecessor(x_i) = x_{i-1}$.

A third property salient to physics: The commutative property of the union of subsets of n -tuples and the summing or multiplication operations defining the number n -tuples and corresponding Euclidean volume values (2) allow n number of values to be sequenced in all $n!$ permutations, for example, $v = width \times height \times depth = depth \times width \times height = \dots$.

Only a cyclic list, where $successor(x_n) = x_1$ and $predecessor(x_1) = x_n$, allows starting at each element and linearly sequencing a successor permutation and a predecessor permutation. For example, where $n = 3$, the volume cyclic successor and predecessor sequences are: $([s_1, s_2, s_3]$ and $[s_1, s_3, s_2])$, $([s_2, s_3, s_1]$ and $[s_2, s_1, s_3])$, $([s_3, s_1, s_2]$ and $[s_3, s_2, s_1])$.

The only cyclic lists that allows linear sequencing of n number of elements in all $n!$ permutations, is where each list element is sequentially cyclic adjacent (either an *immediate* cyclic successor or an *immediate* cyclic predecessor) to every other element, referred to, in this article, as an immediate symmetric cyclic list (ISCL). For example, the cyclic permutations, $[x_2, x_3, x_1]$ and $[x_2, x_1, x_3]$, require x_1 and x_3 be sequentially cyclic adjacent. An ISCL will be proved to contain $n \leq 3$ elements.

D. Third reduction step

Two consequences of the Euclidean distance dimension $\subseteq \mathbb{R}$ property, the linearly sequenced, and commutative properties of volume domain elements are:

1. **ISCL:** An ISCL of 3 distance dimensions, $\{r_1, r_2, r_3\}$, and an ISCL of 3 non-distance dimensions $\{t \text{ (time)}, m \text{ (mass)}, q \text{ (charge)}\}$, each dimension having the $\subseteq \mathbb{R}$ membership property.
2. **Corresponding lengths:** Where all Euclidean distance, time, mass, and charge are each divided into n number of constant-sized interval lengths, for i in 1 to n , the i^{th} constant unit interval length, r_p , of Euclidean distance corresponds to the i^{th} constant unit interval lengths: t_p of time, m_p of mass, and q_p of charge, where “unit” means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

The properties of $\mathbb{R}$ and the corresponding lengths imply 3 direct proportion unit ratios: $r/t = r_p/t_p$, $r/m = r_p/m_p$, and $r/q = r_p/q_p$, letting: $c_t = r_p/t_p$, $c_m = r_p/m_p$, and $c_q = r_p/q_p$. And the correspondence of unit interval lengths implies 3 inverse proportion unit ratios: $rt = t_p r_p$, $rm = m_p r_p$, and $r q = q_p r_p$. And let: $k_t = t_p r_p$, $k_m = m_p r_p$, and $k_q = q_p r_p$.

The gravity,^{13,14} special relativity,^{2,3} Schwarzschild black hole metric,^{15,16} charge equations,¹⁴ electromagnetism equations,¹⁴ and related constants are derived from the 3 direct proportion unit ratios. The Planck-Einstein relation,¹⁴ Compton wavelength,¹⁴ Schrödinger wave equation,⁴ Dirac wave equation,¹⁷ and related constants are derived from combining some of the direct and inverse proportion unit ratios. And the inverse proportion ratios are used to add quantum extensions to Scharwzchild’s metric, Newton’s gravity force equation, and Coulomb’s charge force equation.

II. VOLUME

A. Euclidean volume

Theorem II.1. *Euclidean volume,*

$$\forall v_c, d_c, |x_i| \in \{0, \mathbb{N}\}, x_i \in \{x_1, \dots, x_n\}, v_c = \prod_{i=1}^n |x_i| \Rightarrow v = \prod_{i=1}^n s_i, \quad s_i, v \in \mathbb{R}. \quad (6)$$

The formal proof, “Euclidean_volume,” is in the Rocq file, euclidrelations.v.

Proof.

$$\forall v_c, |x_i| \in \{0, \mathbb{N}\}, \quad \kappa \in \mathbb{R} : \quad v_c = \prod_{i=1}^n |x_i| \Leftrightarrow v_c \kappa = (\prod_{i=1}^n |x_i|) \kappa = \prod_{i=1}^n (|x_i| \kappa^{1/n}). \quad (7)$$

$$v_c \kappa = \prod_{i=1}^n (|x_i| \kappa^{1/n}) \quad \wedge \quad \exists v, s_i \in \mathbb{R} : \quad v = v_c \kappa \quad \wedge \quad s_i = |x_i| \kappa^{1/n} \Rightarrow v = \prod_{i=1}^n s_i. \quad \square \quad (8)$$

Lemma II.2. *The number of n -tuples, v_c , is the sum of the number of n -tuples, v_{c_i} , in each disjoint subset of n -tuples, implies a volume is the sum of volumes,*

$$v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v = \sum_{i=1}^m v_i, \quad v, v_i \in \mathbb{R}.$$

The formal proof, “sum_of_volumes,” is in the Rocq file, euclidrelations.v.

Proof. From the condition of this theorem and the Euclidean volume theorem (II.1):

$$\forall \kappa \in \mathbb{R} : \quad v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v_c \kappa = (\sum_{i=1}^m v_{c_i}) \kappa = \sum_{i=1}^m (v_{c_i} \kappa). \quad (9)$$

$$v_c \kappa = \sum_{i=1}^m (v_{c_i} \kappa) \quad \wedge \quad v = v_c \kappa \quad \wedge \quad v_i = v_{c_i} \kappa \Rightarrow v = \sum_{i=1}^m v_i. \quad \square \quad (10)$$

III. DISTANCE

Definition III.1. Bijective, countable domain value, d_c :

$$\exists v_c, d_c, |x_i| \in \{0, \mathbb{N}\}, \quad x_i \in \{x_1, \dots, x_n\}, \quad v_c = \prod_{i=1}^n |x_i| = \prod_{i=1}^n d_c = d_c^n. \quad (11)$$

A. Vector inner product

Theorem III.2. *Sum of volumes distance:*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} \Rightarrow d^n = v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}).$$

The formal proof, “sum_of_volumes_distance,” is in the Rocq file, euclidrelations.v.

Proof. Multiply both sides of the assumption by κ and apply the Euclidean volume proof (II.1) to v_i :

$$v_c = \sum_{i=1}^m v_{c_i} \Rightarrow v = \sum_{i=1}^m v_i, \quad v, v_i \in \mathbb{R}. \quad (12)$$

$$v = \sum_{i=1}^m v_i \quad \wedge \quad v_i = \prod_{j=1}^n s_{i,j} \Rightarrow v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad (13)$$

$$d_c^n = v_c \Rightarrow d_c^n \kappa = (d_c \kappa^{1/n})^n = v_c \kappa. \quad (14)$$

$$\exists d, v \in \mathbb{R} : d = d_c \kappa^{1/n} \quad \wedge \quad v = v_c \kappa \quad \wedge \quad (d_c \kappa^{1/n})^n = v_c \kappa \Rightarrow d^n = v. \quad (15)$$

$$d^n = v \quad \wedge \quad v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}) \Rightarrow d^n = v = \sum_{i=1}^m (\prod_{j=1}^n s_{i,j}). \quad \square \quad (16)$$

The $n = 2$ case is the inner product, where d is the norm:⁶

$$d^2 = \sum_{i=1}^m (\prod_{j=1}^{n=2} s_{i,j}) \wedge \exists a_{i,1}, a_{i,2} \in \mathbb{R} : s_{i,1} = a_i, s_{i,2} = b_i \Rightarrow d^2 = \sum_{i=1}^m a_i b_i \stackrel{\text{def}}{=} \mathbf{a} \cdot \mathbf{b}. \quad (17)$$

The inner product also allows a_i and b_i to be complex numbers, functions, and matrices.

B. Minkowski distance (p -norm)

Theorem III.3. *Minkowski distance (p -norm):*

$$d_c^n = v_c = \sum_{i=1}^m v_{c_i} = \sum_{i=1}^m d_{c_i}^n \Leftrightarrow d^n = \sum_{i=1}^m d_i^n.$$

The formal proof, “Minkowski_distance,” is in the Rocq file, euclidrelations.v.

Proof. From the sum of volumes distance theorem III.2:

$$d_{c_i}^n = v_{c_i} \Rightarrow d_i^n = v_i. \quad (18)$$

$$d^n = v = \sum_{i=1}^m v_i \wedge d_i^n = v_i \Rightarrow d^n = \sum_{i=1}^m d_i^n \quad \square \quad (19)$$

C. Distance inequalities

Theorem III.4. *Distance triangle inequality*

$$\forall n \in \mathbb{N}, v_a, v_b \geq 0 : (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}.$$

The formal proof, distance_inequality, is in the Rocq file, euclidrelations.v.

Proof. Expand $(v_a^{1/n} + v_b^{1/n})^n$ using the binomial expansion:

$$\begin{aligned} \forall v_a, v_b \geq 0 : v_a + v_b &\leq v_a + v_b + \sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^{n-i} (v_b^{1/n})^i + \\ &\sum_{i=1}^n \binom{n}{i} (v_a^{1/n})^i (v_b^{1/n})^{n-i} = (v_a^{1/n} + v_b^{1/n})^n. \end{aligned} \quad (20)$$

Take the n^{th} root of both sides of the inequality 20:

$$\forall v_a, v_b \geq 0, n \in \mathbb{N} : v_a + v_b \leq (v_a^{1/n} + v_b^{1/n})^n \Rightarrow (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n}. \quad (21)$$

□

Theorem III.5. *Distance sum inequality*

$$\forall m, n \in \mathbb{N}, a_i, b_i \geq 0 : (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}.$$

The formal proof, distance_sum_inequality, is in the Rocq file, euclidrelations.v.

Proof. Apply the distance triangle inequality (III.4):

$$\begin{aligned} \forall m, n \in \mathbb{N}, v_a, v_b \geq 0 : v_a = \sum_{i=1}^m a_i^n \quad \wedge \quad v_b = \sum_{i=1}^m b_i^n \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\ \Rightarrow \quad ((\sum_{i=1}^m a_i^n) + (\sum_{i=1}^m b_i^n))^{1/n} = (\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq \\ (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad \square \quad (22) \end{aligned}$$

D. Metric Space

All Minkowski distances (p -norms) imply the metric space properties. The formal proofs: triangle_inequality, symmetry, non_negativity, and identity_of_indiscernibles are in the Rocq file, euclidrelations.v.

Theorem III.6. *Triangle Inequality:*

$$\begin{aligned} d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k \\ \Rightarrow \quad d(u, w) = (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \end{aligned}$$

Proof. $\forall p \geq 1, \quad u = s_1, \quad w = s_2, \quad v = w/k:$

$$(u^p + w^p)^{1/p} \leq ((u^p + w^p) + 2v^p)^{1/p} = ((u^p + v^p) + (v^p + w^p))^{1/p}. \quad (23)$$

Apply the distance triangle inequality (III.4) to the inequality 23:

$$\begin{aligned} (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \quad \wedge \quad (v_a + v_b)^{1/n} \leq v_a^{1/n} + v_b^{1/n} \\ \wedge \quad v_a = u^p + v^p \quad \wedge \quad v_b = v^p + w^p \\ \Rightarrow \quad (u^p + w^p)^{1/p} \leq ((u^p + v^p) + (v^p + w^p))^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} \\ \Rightarrow \quad d(u, w) = (u^p + w^p)^{1/p} \leq (u^p + v^p)^{1/p} + (v^p + w^p)^{1/p} = d(u, v) + d(v, w). \quad \square \quad (24) \end{aligned}$$

Theorem III.7. *Symmetry:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \geq 1, \quad u = s_1, \quad v = s_2 \quad \Rightarrow \quad d(u, v) = d(v, u).$$

Proof. By the commutative law of addition:

$$\begin{aligned} \forall p : p \geq 1, \quad d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p} = (s_1^p + s_2^p)^{1/p} \\ \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} = (v^p + u^p)^{1/p} = d(v, u). \quad \square \quad (25) \end{aligned}$$

Theorem III.8. *Non-negativity:*

$$d(s_1, s_2) = (\sum_{i=1}^2 s_i^p)^{1/p}, \quad p \in N \quad \wedge \quad s_1, s_2 \geq 0 \quad \Rightarrow \quad d(u, v) \geq 0.$$

Proof.

$$\forall n \geq 1 \quad \wedge \quad s_1, s_2 \geq 0 \quad \Rightarrow \quad d(s_1, s_2) = (s_1^p + s_2^p)^{1/p} \geq 0. \quad (26)$$

$$\forall u = s_1, v = s_2 \quad \wedge \quad d(s_1, s_2) = (s_1^p + s_2^p)^{1/p} \geq 0 \quad \Rightarrow \quad d(u, v) = (u^p + v^p)^{1/p} \geq 0. \quad \square \quad (27)$$

Theorem III.9. *Identity of Indiscernibles:* $d(u, u) = 0$.

Proof. From the non-negativity property (III.8):

$$d(u, w) \geq 0 \quad \wedge \quad d(u, v) \geq 0 \quad \wedge \quad d(v, w) \geq 0 \quad (28)$$

$$\Rightarrow \quad \exists d(u, w) = d(u, v) = d(v, w) = 0. \quad (29)$$

$$d(u, w) = d(v, w) = 0 \quad \Rightarrow \quad u = v. \quad (30)$$

$$d(u, v) = 0 \quad \wedge \quad u = v \quad \Rightarrow \quad d(u, u) = 0. \quad \square \quad (31)$$

IV. PROPERTIES LIMITING A SET TO AT MOST 3 ELEMENTS

The following definitions and proof use first order logic. A Horn clause-like expression is used, here, to make the proof easier to read. By convention, the proof goal, in a Horn clause, is on the left side of the implication sign ($\leftarrow$) and supporting facts are on the right side. The formal proofs in the Rocq file `threed.v` are: `adj111`, `adj122`, `adj212`, `adj123`, `adj133`, `adj233`, `adj213`, `adj313`, `adj323`, and `not_all_mutually_adjacent_gt_3`.

Definition IV.1. Immediate Cyclic Successor of m is n :

$$\forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} :$$

$$\text{Successor}(m, n, \text{setsize}) \leftarrow (m = \text{setsize} \wedge n = 1) \vee (n = m + 1 \wedge n \leq \text{setsize}). \quad (32)$$

Definition IV.2. Immediate Cyclic Predecessor of m is n :

$$\forall x_m, x_n \in \{x_1, \dots, x_{\text{setsize}}\} :$$

$$\text{Predecessor}(m, n, \text{setsize}) \leftarrow (m = 1 \wedge n = \text{setsize}) \vee (n = m - 1 \wedge n \geq 1). \quad (33)$$

Definition IV.3. Adjacent: Element m is sequentially cyclic adjacent to element n if the immediate cyclic successor of m is n or the immediate cyclic predecessor of m is n :

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : \\ Adjacent(m, n, setsize) \leftarrow Successor(m, n, setsize) \vee Predecessor(m, n, setsize). \quad (34)$$

Definition IV.4. Immediate Symmetric Cyclic List (ISCL), where every set element is sequentially adjacent to every other element):

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : \quad Adjacent(m, n, setsize). \quad (35)$$

Theorem IV.5. An ISCL is limited to at most 3 elements.

$$\forall x_m, x_n \in \{x_1, \dots, x_{setsize}\} : \quad Adjacent(m, n, setsize) \Rightarrow setsize \leq 3. \quad (36)$$

Proof.

Every element is adjacent to every other element, where $setsize \in \{1, 2, 3\}$:

$$Adjacent(1, 1, 1) \leftarrow Successor(1, 1, 1) \leftarrow (m = setsize \wedge n = 1). \quad (37)$$

$$Adjacent(1, 2, 2) \leftarrow Successor(1, 2, 2) \leftarrow (n = m + 1 \wedge n \leq setsize). \quad (38)$$

$$Adjacent(1, 2, 3) \leftarrow Successor(1, 2, 3) \leftarrow (n = m + 1 \wedge n \leq setsize). \quad (39)$$

$$Adjacent(2, 1, 3) \leftarrow Predecessor(2, 1, 3) \leftarrow (n = m - 1 \wedge n \geq 1). \quad (40)$$

$$Adjacent(3, 1, 3) \leftarrow Successor(3, 1, 3) \leftarrow (n = setsize \wedge m = 1). \quad (41)$$

$$Adjacent(1, 3, 3) \leftarrow Predecessor(1, 3, 3) \leftarrow (m = 1 \wedge n = setsize). \quad (42)$$

$$Adjacent(2, 3, 3) \leftarrow Successor(2, 3, 3) \leftarrow (n = m + 1 \wedge n \leq setsize). \quad (43)$$

$$Adjacent(3, 2, 3) \leftarrow Predecessor(3, 2, 3) \leftarrow (n = m - 1 \wedge n \geq 1). \quad (44)$$

Element 2 is the only immediate successor of element 1 for all $setsize \geq 3$, which implies element 3 is not ($\neg$) an immediate successor of element 1 for all $setsize \geq 3$:

$$\neg Successor(1, 3, setsize \geq 3) \leftarrow \\ Successor(1, 2, setsize \geq 3) \leftarrow (n = m + 1 \leq setsize). \quad (45)$$

Element $n = setsize > 3$ is the only immediate predecessor of element 1, which implies element 3 is not ($\neg$) an immediate predecessor of element 1 for all $setsize > 3$:

$$\neg Predecessor(1, 3, setsize \geq 3) \leftarrow Predecessor(1, setsize, setsize > 3) \leftarrow \\ (m = 1 \wedge n = setsize > 3). \quad (46)$$

For all $setsize > 3$, some elements are not ($\neg$) sequentially adjacent to every other element (not immediate symmetric):

$$\neg Adjacent(1, 3, setsize > 3) \leftarrow \neg Successor(1, 3, setsize > 3) \quad \wedge \quad \neg Predecessor(1, 3, setsize > 3). \quad \square \quad (47)$$

The Symmetric goal matches the Adjacent goals 37 and fails for all “setsize” greater than three.

V. DERIVATION OF PHYSICS EQUATIONS AND CONSTANTS

A. 3 direct and 3 inverse proportion ratios

Application of the ISCL property of volume (IV.5) and the Euclidean distance dimension $\subseteq \mathbb{R}$ (III.3):

1. **ISCL:** $\{r_1, r_2, r_3\}$ is an ISCL of 3 “distance” dimensions, each dimension having the $\subseteq \mathbb{R}$ set membership property and $\{t \text{ (time)}, m \text{ (mass)}, q \text{ (charge)}\}$ is an ISCL of 3 “non-distance” dimensions, each dimension $\subseteq \mathbb{R}$. There is a total of 6 dimensions with the $\subseteq \mathbb{R}$ property: r_1, r_2, r_3, t, m, q .
2. **Corresponding lengths:** Where all Euclidean distance, time, mass, and charge are each divided into n number of constant-sized interval lengths, for i in 1 to n , the i^{th} constant unit interval length, r_p , of Euclidean distance corresponds to the i^{th} constant unit interval lengths: t_p of time, m_p of mass, and q_p of charge, where “unit” means a measurement type, for example, meters, seconds, kilograms, and Coulombs.

There are 3 constant direct proportion ratios from the properties of $\mathbb{R}$:

$$\forall r_p, t_p, m_p, q_p, k_1, k_2, k_3 \in \mathbb{R}, \exists r, t, m, q \in \mathbb{R} : \frac{r_p}{t_p} = \frac{r_p k_1}{t_p k_1} = \frac{r}{t} \quad \wedge \quad \frac{r_p}{m_p} = \frac{r_p k_2}{m_p k_2} = \frac{r}{m} \quad \wedge \quad \frac{r_p}{q_p} = \frac{r_p k_3}{q_p k_3} = \frac{r}{q}. \quad (48)$$

$$\begin{aligned} \exists c, c_t, c_m, c_q \in \mathbb{R} : \quad r_p/t_p = c_t = c \quad \wedge \quad r_p/m_p = c_m \quad \wedge \quad r_p/q_p = c_q \\ \Rightarrow \quad r/t = r_p/t_p = c_t = c \quad \wedge \quad r/m = r_p/m_p = c_m \quad \wedge \quad r/q = r_p/q_p = c_q. \end{aligned} \quad (49)$$

And there are 3 constant inverse proportion ratios from the properties of $\mathbb{R}$:

$$\begin{aligned} \forall r_p, t_p, m_p, q_p, k_1, k_2, k_3 \in \mathbb{R}, \exists r, t, m, q \in \mathbb{R} : \\ r_p t_p = r_p t_p \left(\frac{k_1}{k_1} \right) = \left(\frac{r_p}{k_1} \right) (t_p k_1) = r t \quad \wedge \quad r_p m_p = r_p m_p \left(\frac{k_2}{k_2} \right) = \left(\frac{r_p}{k_2} \right) (m_p k_2) = r m \\ \wedge \quad r_p q_p = r_p q_p \left(\frac{k_3}{k_3} \right) = \left(\frac{r_p}{k_3} \right) (q_p k_3) = r q. \end{aligned} \quad (50)$$

$$\begin{aligned} \exists k_t, k_m, k_q \in \mathbb{R} : \quad r_p t_p = k_t \quad \wedge \quad r_p m_p = k_m \quad \wedge \quad r_p q_p = k_q \\ \Rightarrow \quad r t = t_p r_p = k_t \quad \wedge \quad r m = m_p r_p = k_m \quad \wedge \quad r q = q_p r_p = k_q. \end{aligned} \quad (51)$$

B. G , Newton, Gauss, Poisson gravity laws

From equations 49, linear acceleration, r/t^2 , can be related to mass, m , and distance, r :

$$r = c_m m \quad \wedge \quad r = c_t t \quad \Rightarrow \quad \frac{r}{(c_t t)^2} = \frac{c_m m}{r^2} \quad \Rightarrow \quad \frac{r}{t^2} = \frac{(c_m c_t^2) m}{r^2} = \frac{G m}{r^2}. \quad (52)$$

Where r_p , t_p , and m_p are the length, time, and mass Planck units, $G = c_m c_t^2 = r_p^3 / m_p t_p^2 \approx 6.67430 \cdot 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}$, which is the empirical value of G and the SI units of G .¹⁸

Newton's law follows from multiplying both sides of equation 52 by m :

$$\frac{r}{t^2} = \frac{G m}{r^2} \quad \Leftrightarrow \quad F \stackrel{\text{def}}{=} \frac{m r}{t^2} = \frac{G m^2}{r^2}. \quad (53)$$

$$F = \frac{G m^2}{r^2} \quad \wedge \quad \forall m \in \mathbb{R} : \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \quad \Rightarrow \quad F = \frac{G m_1 m_2}{r^2}. \quad (54)$$

A new interpretation of Gauss's and Poisson's gravity laws is presented here. Equation 52 relates linear acceleration, r/t^2 , to mass and distance. Gauss's gravity vector field, $\mathbf{g}$, and Poisson's gravity scalar field, $-\nabla\Phi(\mathbf{r})$, is the angular acceleration, $2\pi r/t^2$. Applying equation 52, where the minus sign is the vector direction (attractive force):

$$\frac{r}{t^2} = -\frac{G m}{|\mathbf{r}|^2} \quad \wedge \quad \mathbf{g} = -\nabla\Phi(\mathbf{r}, t) = \frac{2\pi r}{t^2} \quad \Rightarrow \quad \mathbf{g} = -\nabla\Phi(\mathbf{r}) = -\frac{2\pi G m}{|\mathbf{r}|^2}. \quad (55)$$

$$\mathbf{g} = -\nabla\Phi(\mathbf{r}) = -\frac{2\pi G m}{|\mathbf{r}|^2} \quad \Rightarrow \quad \nabla \cdot \mathbf{g} = \nabla^2\Phi(\mathbf{r}) = \frac{4\pi G m}{|\mathbf{r}|^3} = 4\pi G \rho, \quad (56)$$

which are the Gauss and Poisson laws, where $\rho = m/|\mathbf{r}|^3$ is the mass density:

C. Special relativity

$$\begin{aligned} \forall \tau \in \{t, m, q\}, r^2 = r'^2 + r_v^2, \exists \mu_\tau, \nu_\tau : r = \mu_\tau \tau \quad \wedge \quad r_v = \nu_\tau \tau \quad \Rightarrow \quad (\mu_\tau \tau)^2 = r'^2 + (\nu_\tau \tau)^2 \\ \Rightarrow \quad r' = \sqrt{(\mu_\tau \tau)^2 - (\nu_\tau \tau)^2} = \mu_\tau \tau \sqrt{1 - (\nu_\tau / \mu_\tau)^2}. \end{aligned} \quad (57)$$

Local frame distance, r' , contracts relative to a distant observer frame distance, r , as $\nu_\tau \rightarrow \mu_\tau$:

$$r' = \mu_\tau \tau \sqrt{1 - (\nu_\tau / \mu_\tau)^2} \quad \wedge \quad \mu_\tau \tau = r \quad \Rightarrow \quad r' = r \sqrt{1 - (\nu_\tau / \mu_\tau)^2}. \quad (58)$$

A distant observer frame type, τ , dilates relative to the local observer frame type, τ' , as $\nu_\tau \rightarrow \mu_\tau$:

$$\mu_\tau \tau = r' / \sqrt{1 - (\nu_\tau / \mu_\tau)^2} \quad \wedge \quad r' = \mu_\tau \tau \quad \Rightarrow \quad \tau = \tau' / \sqrt{1 - (\nu_\tau / \mu_\tau)^2}. \quad (59)$$

Where τ is type, time, the space-like flat Minkowski spacetime event interval is:

$$\begin{aligned} dr^2 = dr'^2 + dr_v^2 \quad \wedge \quad dr_v^2 = dr_1^2 + dr_2^2 + dr_3^2 \quad \wedge \quad d(\mu_\tau \tau) = dr \\ \Rightarrow \quad dr'^2 = d(\mu_\tau \tau)^2 - dr_1^2 - dr_2^2 - dr_3^2. \end{aligned} \quad (60)$$

D. Analytic general relativity solutions

The Einstein field equation (EFE) is:

$$\mathbf{G}_{\mu,\nu} + \Lambda g_{\mu,\nu} = \kappa T_{\mu,\nu}, \quad \mathbf{G}_{\mu,\nu} = \mathbf{R} + \frac{1}{2} R g_{\mu,\nu}. \quad (61)$$

where: $\mathbf{G}_{\mu,\nu}$ is Einstein's tensor, $\mathbf{R}$ is the Ricci curvature, R is the scalar curvature, $g_{\mu,\nu}$ is the metric tensor, Λ is a cosmological constant, $\kappa = 8\pi G/c^4$, is Einstein's constant, and $T_{\mu,\nu}$ is the stress-energy-momentum tensor.^{2,3}

The EFE is a set of second order differential equation extending the previously derived second order differential equation, Poisson's law of gravity, $\nabla^2 \Phi(\mathbf{r}) = 4\pi G \rho$ (56), to include a time component. The Minkowski spacetime interval component for time is $c^2 g_{0,0} dt^2$.³ And from convention, $T_{0,0} = mc^2/|\mathbf{r}|^3$ is the energy density component of $T_{\mu,\nu}$.³

$$\frac{c^2}{2} (\mathbf{G}_{0,0} + \Lambda g_{0,0}) = \nabla^2 \Phi(\mathbf{r}, t) = 4\pi G \rho = -4\pi G \frac{m}{|\mathbf{r}|^3} = \frac{4\pi G}{c^2} \cdot \frac{mc^2}{|\mathbf{r}|^3} = \frac{4\pi G}{c^2} T_{0,0}. \quad (62)$$

$$\frac{c^2}{2} (\mathbf{G}_{0,0} + \Lambda g_{0,0}) = \frac{4\pi G}{c^2} T_{0,0} \quad \Rightarrow \quad \mathbf{G}_{0,0} + \Lambda g_{0,0} = \frac{8\pi G}{c^4} T_{0,0} = \kappa T_{0,0}. \quad (63)$$

The EFEs are nonlinear because gravity itself carries energy. *However:* 1) The laws of physics determine the amount of spacetime curvature and the distribution of stress, energy, and momentum. 2) The laws of physics are specified as the components of the metric, $g_{\mu,\nu}$. 3) $g_{\mu,\nu}$ can be determined independent of the EFE, for those cases where the laws (equations) expressing the interaction of gravity waves and gravity fields can be derived from the ratios. Derivation of the Schwarzschild metric from ratios, in next subsection, is an example.

Step 1) Use the unit ratios and equations (laws) derived from the ratios, for example, the ratio-derived special relativity equations (58) and the ratio-derived constant, G (52), to derive the scalar functions (laws) assigned to the components of $g_{\mu,\nu}$.

Step 2) The 4D flat spacetime interval equation is an instance of the 2D equation (60). Therefore, derive the metric first as a 2D metric and expand later to a 4D metric.

(Optional step 2d) The 2D metric tensor uses the much simpler 2D Ricci curvature (which, in 2D, is the Gaussian curvature). And the 2D tensors reduce the number of independent equations to solve. The 2D solutions can next be used to set constraints on the solutions in the 4D tensor, much like an Euclidean distance value places constraints on the coordinate distance values.

Step 3) Translate from 2D to 4D. For spherically symmetric cases, the simplest method to translate from 2D to 4D is to use spherical coordinates, where the first 2 diagonal components of $g_{\nu,\mu}$ remain unchanged and the other 2 diagonal components are functions of the longitude angle, θ , and latitude angle, ϕ .

E. Schwarzschild's time dilation and metric

From equations 58 and 49:

$$r = c_m m \quad \Rightarrow \quad c_m m / r = 1 \quad \Rightarrow \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - c_m m v^2 / r c^2}. \quad (64)$$

Where v_{escape} is the escape velocity and KE is the kinetic energy:

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} = \sqrt{1 - c_m m v^2 / r c^2} \quad \wedge \quad KE = m v^2 / 2 = m v_{escape}^2 \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2 c_m m v_{escape}^2 / r c^2}. \end{aligned} \quad (65)$$

For a photon, the escape velocity, $v_{\text{escape}} = c$.

$$\begin{aligned} \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2c_m m v_{\text{escape}}^2 / r c^2} \quad \wedge \quad v_{\text{escape}} = c \\ &\Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2}. \end{aligned} \quad (66)$$

Combining equation 66 with the derivation of G (52):

$$c_m c^2 = G \quad \wedge \quad \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2c_m m c^2 / r c^2} \Rightarrow \sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm / r c^2}. \quad (67)$$

Combining equation 67 with equation 59 yields Schwarzschild's gravitational time dilation:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Gm / r c^2} \quad \wedge \quad t' = t \sqrt{1 - (v^2/c^2)} \Rightarrow t' = t \sqrt{1 - 2Gm / r c^2}. \quad (68)$$

From equations 58 and 67, where Schwarzschild defined the black hole event horizon radius as $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$:^{15,16}

$$r' = r \sqrt{1 - (v/c)^2} = r \sqrt{1 - 2Gm / r c^2} \quad \wedge \quad \alpha \stackrel{\text{def}}{=} 2Gm/c^2 \Rightarrow r' = r \sqrt{1 - \alpha/r}. \quad (69)$$

Applying equation 69 to the time-like spacetime interval equation 60, where $ds^2 < 0$:

$$\begin{aligned} r' &= r \sqrt{1 - \alpha/r} \quad \wedge \quad ds^2 = dr'^2 - dt^2 \\ \Rightarrow \quad ds^2 &= (\sqrt{1 - \alpha/r} dr)^2 - (dt)^2 = (1 - \alpha/r) dr^2 - dt^2. \end{aligned} \quad (70)$$

$$\begin{aligned} ds^2 &= (1 - \alpha/r) dr^2 - dt^2 \quad \wedge \quad dr = d(ct) \quad \wedge \quad c = 1 \\ \lim_{ds \rightarrow 0} dr' &= \lim_{ds \rightarrow 0} dr \Rightarrow \lim_{ds \rightarrow 0} ds^2 = (1 - \alpha/r) dt^2 - dt^2. \end{aligned} \quad (71)$$

Using spherical coordinates to translate from 2D to 4D yields the $+- --$ form of Schwarzschild's black hole metric:^{15,16}

$$\begin{aligned} \lim_{ds \rightarrow 0} ds^2 &= (1 - \alpha/r) dt^2 - dr^2 \\ \Rightarrow \quad \lim_{ds \rightarrow 0} ds^2 &= (1 - \alpha/r) dt^2 - (1 - \alpha/r) dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \\ \Rightarrow \quad g_{\mu,\nu} &= \text{diag}[(1 - \alpha/r), -(1 - \alpha/r)^{-1}, -r^2, -r^2 \sin^2 \theta]. \end{aligned} \quad (72)$$

F. k_e , ε_0 , **E**, Coulomb and Gauss laws

From equations 49, linear acceleration, r/t^2 , can be related to charge, q , and distance, r :

$$r = c_q q \quad \wedge \quad r = c_t t \Rightarrow \frac{r}{(c_t t)^2} = \frac{c_q q}{r^2} \Rightarrow \frac{r}{t^2} = \frac{(c_q c_t^2) q}{r^2}. \quad (73)$$

$$r = c_m m \quad \wedge \quad r = c_q q \Rightarrow c_m m = c_q q \Rightarrow m = (c_q / c_m) q. \quad (74)$$

Combining equations 73 and 74 yields Coulombs's law:

$$\frac{r}{t^2} = \frac{(c_q c_t^2)q}{r^2} \quad \wedge \quad m = (c_q/c_m)q \quad \Rightarrow \quad F \stackrel{\text{def}}{=} \frac{mr}{t^2} = \frac{(c_q c_t^2/c_m)q}{r^2} = \frac{k_e q^2}{r^2}. \quad (75)$$

$k_e = c_q^2 c_t^2 / c_m = m_p r_p^3 / t_p^2 q_p^2$, conforms to the SI units: $kg \cdot m^3 \cdot s^{-2} \cdot C^{-2} \stackrel{\text{def}}{=} N \cdot m^2 \cdot C^{-2}$.¹⁴

$$\forall q \in \mathbb{R} \exists q_1, q_2 \in \mathbb{R} : |q_1 q_2| = q^2 \quad \wedge \quad F = \frac{k_e q^2}{r^2}. \quad \Rightarrow \quad F = \frac{k_e |q_1 q_2|}{r^2}. \quad (76)$$

A new interpretation of Gauss's electric field law is presented here. Equation 73 relates the linear acceleration, r/t^2 , to charge and distance. Gauss's electric field, $\mathbf{E}$, relates angular acceleration, $2\pi r/t^2$ to charge and distance. Applying equation 75:

$$F_C = \frac{mr}{t^2} = \frac{k_e q^2}{r^2} \quad \Rightarrow \quad \exists F_E \in \mathbb{R} : F_E = \frac{2\pi mr}{t^2} = \frac{2\pi k_e q^2}{r^2}. \quad (77)$$

$$F_E = \frac{2\pi k_e q^2}{r^2} = q \left(\frac{2\pi k_e q}{r^2} \right) \quad \wedge \quad \exists E \in \mathbb{R} : E = \frac{2\pi k_e q}{r^2} \quad \Rightarrow \quad F_E = qE. \quad (78)$$

The electric field, $E = 2\pi k_e q/r^2$, conforms to the SI units $kg \cdot m \cdot s^{-2} \cdot C^{-1} \stackrel{\text{def}}{=} kg \cdot m \cdot s^{-3} \cdot A^{-1}$.

$$\mathbf{E} = \frac{2\pi k_e q}{|\mathbf{r}|^2} \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -\frac{4\pi k_e q}{|\mathbf{r}|^3}. \quad (79)$$

$$\nabla \cdot \mathbf{E} = -\frac{4\pi k_e q}{|\mathbf{r}|^3} \quad \wedge \quad \varepsilon_0 \stackrel{\text{def}}{=} \frac{1}{4\pi k_e} \quad \wedge \quad \rho \stackrel{\text{def}}{=} \frac{q}{|\mathbf{r}|^3} \quad \Rightarrow \quad \nabla \cdot \mathbf{E} = -\rho/\varepsilon_0, \quad (80)$$

which is Gauss's electric field law in differential form.¹⁴

G. $\mathbf{B}$, μ_0 , and Lorentz's law

Applying the distance contraction equation, 58, to equation 78, where r' is the distant observer frame of reference and r is local (rest) frame of reference:

$$r' = r/\sqrt{1-v^2/c^2} \quad \wedge \quad F' = \frac{2\pi k_e q^2}{r'^2} \quad \Rightarrow \quad F = \frac{2\pi k_e q^2 (1-v^2/c^2)}{r^2}. \quad (81)$$

From equation 78:

$$E = \frac{2\pi k_e q}{r^2} \quad \wedge \quad F = \frac{2\pi k_e q^2 (1-v^2/c^2)}{r^2} \quad \Rightarrow \quad F = q \left(E - \frac{(2\pi k_e/c^2)q}{r^2} v^2 \right). \quad (82)$$

$$F = q \left(E - \frac{(2\pi k_e/c^2)q}{r^2} v^2 \right) \quad \wedge \quad \exists B : B = \frac{(2\pi k_e/c^2)vq}{r^2} \quad \Rightarrow \quad F = q(E - Bv), \quad (83)$$

where the magnetic field, $B = (2\pi k_e/c)q/r^2$, conforms to the base SI units: $kg \cdot C^{-1} \cdot s^{-1} \stackrel{\text{def}}{=} kg \cdot A^{-1} \cdot s^{-2} \stackrel{\text{def}}{=} T$. And assigning the constant portion to μ_0 :

$$B = \frac{(2\pi k_e/c^2)vq}{r^2} \quad \Rightarrow \quad \frac{\partial B}{\partial r} = \frac{(4\pi k_e/c)vq}{r^2} = \frac{\mu_0 vq}{r^3}. \quad (84)$$

where $\mu_0 = 4\pi k_e/c^2$ conforms to the SI units $kg \cdot m \cdot C^{-2} \stackrel{\text{def}}{=} kg \cdot m \cdot s^{-2} A^{-2} \stackrel{\text{def}}{=} N \cdot A^{-2}$.

Translating equation 83 to vector form:

$$F = q(E - Bv) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \mathbf{v}). \quad (85)$$

$$\mathbf{B} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{B}) \quad \wedge \quad \mathbf{F} = q(\mathbf{E} - \mathbf{B} \times \mathbf{v}) \quad \Rightarrow \quad \mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}), \quad (86)$$

which is Lorentz law in the local frame of reference.

H. Faraday's law

From the magnetic field equation 86, where the electric and magnetic fields are propagating at the speed, $v = c$:

$$B = \frac{(2\pi k_e/c^2)vq}{r^2} \quad \wedge \quad v = c \quad \wedge \quad r = ct \quad \Rightarrow \quad B = \frac{(2\pi k_e/c^3)q}{t^2}. \quad (87)$$

$$B = \frac{(2\pi k_e/c^3)q}{t^2} \quad \Rightarrow \quad \frac{\partial B}{\partial t} = -\frac{(4\pi k_e/c^3)q}{t^3}. \quad (88)$$

$$\frac{\partial B}{\partial t} = -\frac{(4\pi k_e/c^3)q}{t^3} \quad \wedge \quad r = ct \quad \Rightarrow \quad \frac{\partial B}{\partial t} = -\frac{4\pi k_e q}{r^3}. \quad (89)$$

Faraday's law is the case, where the electric field, $\mathbf{E}$, is a single radial vector, $\mathbf{E}_{\mathbf{r}_1}$ acting on some point, P . Therefore, at point, P , $\mathbf{E}_{\mathbf{r}_2} = 0$ and $\mathbf{E}_{\mathbf{r}_3} = 0$. From the Lorentz equation 86, the magnetic field, $\mathbf{B}$ at point P , is perpendicular to $\mathbf{E}_{\mathbf{r}_1}$ (aligned on $\mathbf{r}_2$). For a constant force, the Lorentz equation makes the electric field, $\mathbf{E}_{\mathbf{r}_1}$, a function of $\mathbf{B}_{\mathbf{r}_2}$. Combining with equation 79:

$$\begin{aligned} \mathbf{E}_{\mathbf{r}_1} &= 2\pi k_e q / |\mathbf{r}_2|^2 = f(\mathbf{B}_{\mathbf{r}_2}) \quad \wedge \quad \mathbf{E}_{\mathbf{r}_2} = 0 \quad \wedge \quad \mathbf{E}_{\mathbf{r}_3} = 0 \\ \Rightarrow \quad \nabla \times \mathbf{E} &= \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ \frac{\partial}{\partial \mathbf{r}_1^2} & \frac{\partial}{\partial \mathbf{r}_2^2} & \frac{\partial}{\partial \mathbf{r}_3^2} \\ \mathbf{E}_{\mathbf{r}_1} & 0 & 0 \end{vmatrix} = \left(\frac{\partial \mathbf{E}_{r_3}}{\partial \mathbf{r}_2^2} - \frac{\partial \mathbf{E}_{r_2}}{\partial \mathbf{r}_3^2} \right) \hat{\mathbf{i}} + \left(\frac{\partial \mathbf{E}_{r_1}}{\partial \mathbf{r}_3^2} - \frac{\partial \mathbf{E}_{r_3}}{\partial \mathbf{r}_1^2} \right) \hat{\mathbf{j}} + \left(\frac{\partial \mathbf{E}_{r_2}}{\partial \mathbf{r}_1^2} - \frac{\partial \mathbf{E}_{r_1}}{\partial \mathbf{r}_2^2} \right) \hat{\mathbf{k}} \\ &= 0\hat{\mathbf{i}} + 0\hat{\mathbf{j}} + \left(0 - \frac{\partial 2\pi k_e q / |\mathbf{r}_2|^2}{\partial |\mathbf{r}_2|^2} \right) \hat{\mathbf{k}} = \frac{4\pi k_e q}{|\mathbf{r}_2|^3} \hat{\mathbf{k}} \equiv \nabla \times \mathbf{E} = \frac{4\pi k_e q}{|\mathbf{r}|^3}. \end{aligned} \quad (90)$$

Combining equations 90 and 89 yields Maxwell-Faraday's law:¹⁴

$$\nabla \times \mathbf{E} = \frac{4\pi k_e q}{|\mathbf{r}|^3} \quad \wedge \quad \frac{\partial \mathbf{B}}{\partial t} = -\frac{4\pi k_e q}{|\mathbf{r}|^3} \quad \Rightarrow \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}. \quad (91)$$

I. Maxwell's wave equations

From the Gauss's electric field equations 80:

$$\mathbf{E} = \frac{2\pi k_e q}{|\mathbf{r}|^2} \Rightarrow \nabla^2 \mathbf{E} = \frac{12\pi k_e q}{|\mathbf{r}|^4}. \quad (92)$$

$$\mathbf{E} = \frac{2\pi k_e q}{|\mathbf{r}|^2} \wedge r = ct \Rightarrow \mathbf{E} = \frac{2\pi k_e q}{(ct)^2} \Rightarrow \frac{\partial^2 \mathbf{E}}{\partial t^2} = \frac{(12\pi k_e/c^2)q}{t^4}. \quad (93)$$

$$\frac{\partial^2 \mathbf{E}}{\partial t^2} = \frac{(12\pi k_e/c^2)q}{t^4} \wedge r = ct \Rightarrow \frac{\partial^2 \mathbf{E}}{\partial t^2} = \frac{12\pi k_e c^2 q}{|\mathbf{r}|^4}. \quad (94)$$

$$\nabla^2 \mathbf{E} = \frac{12\pi k_e q}{|\mathbf{r}|^4} \wedge \frac{\partial^2 \mathbf{E}}{\partial t^2} = \frac{12\pi k_e c^2 q}{|\mathbf{r}|^4} \Rightarrow \nabla^2 \mathbf{E} = \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2}. \quad (95)$$

$$\nabla^2 \mathbf{E} = \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} \wedge \varepsilon_0 \mu_0 = \frac{1}{4\pi k_e} \cdot \frac{4\pi k_e}{c^2} = \frac{1}{c^2} \Rightarrow \nabla^2 \mathbf{E} = \varepsilon_0 \mu_0 \frac{\partial^2 \mathbf{E}}{\partial t^2}, \quad (96)$$

which is Maxwell's electric field wave equation. And from equation (89):

$$\frac{\partial \mathbf{B}}{\partial t} = -\frac{4\pi k_e q}{r^3} \wedge r = |\mathbf{r}| \Rightarrow \nabla^2 \mathbf{B} = \frac{12\pi k_e q}{|\mathbf{r}|^4}. \quad (97)$$

$$\mathbf{B} = \frac{2\pi k_e q}{|\mathbf{r}|^2} \wedge r = ct \Rightarrow \mathbf{B} = \frac{2\pi k_e q}{(ct)^2} \Rightarrow \frac{\partial^2 \mathbf{B}}{\partial t^2} = \frac{(12\pi k_e/c^2)q}{t^4}. \quad (98)$$

$$\frac{\partial^2 \mathbf{B}}{\partial t^2} = \frac{(12\pi k_e/c^2)q}{t^4} \wedge r = ct \Rightarrow \frac{\partial^2 \mathbf{B}}{\partial t^2} = \frac{12\pi k_e c^2 q}{|\mathbf{r}|^4}. \quad (99)$$

$$\nabla^2 \mathbf{B} = -12\pi k_e q/|\mathbf{r}|^4 \wedge \frac{\partial^2 \mathbf{B}}{\partial t^2} = \frac{12\pi k_e c^2 q}{|\mathbf{r}|^4} \Rightarrow \nabla^2 \mathbf{B} = \frac{1}{c^2} \frac{\partial^2 \mathbf{B}}{\partial t^2}. \quad (100)$$

$$\nabla^2 \mathbf{B} = \frac{1}{c^2} \frac{\partial^2 \mathbf{B}}{\partial t^2} \wedge \varepsilon_0 \mu_0 = \frac{1}{4\pi k_e} \cdot \frac{4\pi k_e}{c^2} = \frac{1}{c^2} \Rightarrow \nabla^2 \mathbf{B} = \varepsilon_0 \mu_0 \frac{\partial^2 \mathbf{B}}{\partial t^2}, \quad (101)$$

which is Maxwell's magnetic field wave equation.

J. The classic electron radius, r_e

From the unit ratios 49:

$$r = c_m m \wedge r = c_q q \Rightarrow c_m m = c_q q \Rightarrow m = (c_q/c_m)q. \quad (102)$$

$$m = (c_q/c_m)q \wedge r = c_q q \Rightarrow mr = (c_q^2/c_m)q^2 = \frac{r_p m_p q^2}{q_p^2}. \quad (103)$$

$$m_e r_e = \frac{r_p m_p q_e^2}{q_p^2} \Rightarrow r_e = r_p \frac{m_p q_e^2}{m_e q_p^2} \approx 2.81794023 \cdot 10^{-15} m, \quad (104)$$

which is classic electron radius value,¹⁴ where m_e is the electron mass, q_e is the electron charge, and r_p , m_p , and q_p are Planck units.

K. The fine structure constants, α and α_G

Rearranging the classic electron radius equation yields the fine structure electron coupling constant, α , and fine structure electron gravity coupling constant, α_G :

$$r_e = r_p \frac{m_p q_e^2}{m_e q_p^2} \Rightarrow \frac{r_e m_e}{r_p m_p} = \frac{q_e^2}{q_p^2} \approx 0.0072973525643 = \alpha. \quad (105)$$

$$r_e = r_p \frac{m_p q_e^2}{m_e q_p^2} \Rightarrow \left(\frac{r_e q_p^2}{r_p q_e^2} \right)^2 = \frac{m_e^2}{m_p^2} \approx 1.751809 \cdot 10^{-45} = \alpha_G, \quad (106)$$

where m_e is the electron mass, q_e is the electron charge, and r_p , m_p , and q_p are Planck units.

L. The Rydberg constant, R_∞

Starting with the ratio-derived classic electron radius (104), rearrange the quantities so that electron properties are in the numerator and the wave (Planck unit) properties are in the denominator:

$$\begin{aligned} r_e = r_p \frac{m_p q_e^2}{m_e q_p^2} &\Rightarrow \frac{r_e}{r_p} = \frac{m_p q_e^2}{m_e q_p^2} \Rightarrow \frac{r_e}{r_p} \left(\frac{\alpha_G \alpha}{4\pi r_p} \right) = \frac{m_p}{m_e} \frac{q_e^2}{q_p^2} \left(\frac{\alpha_G \alpha}{4\pi r_p} \right) \\ &\Rightarrow \frac{r_e}{4\pi r_p^2} \frac{m_e^2 q_e^2}{m_p^2 q_p^2} = \frac{m_e q_e^4}{4\pi r_p m_p q_p^4} \approx 1.09737315 \cdot 10^7 \text{ m}^{-1} = R_\infty, \end{aligned} \quad (107)$$

where m_e is the electron mass, q_e is the electron charge, and r_p , m_p , and q_p are Planck units.

M. $\hbar$, h , and Planck-Einstein relation

Applying both the direct proportion ratio (49), and inverse proportion ratio (51):

$$m = k_m/r \quad \wedge \quad r = ct \quad \Rightarrow \quad m(ct)^2 = (k_m/r)r^2 = k_m r. \quad (108)$$

$$m(ct)^2 = k_m r \quad \Rightarrow \quad E \stackrel{\text{def}}{=} mc^2 = k_m r/t^2. \quad (109)$$

$$E = mc^2 = k_m r/t^2 \quad \wedge \quad r/t = c \quad \Rightarrow \quad E = mc^2 = k_m c/t. \quad (110)$$

Solving the fine structure constant equation 105 for $r_p m_p$, where $r_p m_p = k_m$:

$$\begin{aligned} \frac{r_e m_e}{r_p m_p} &= \frac{q_e^2}{q_p^2} \approx 0.0072973525705 = \alpha \\ \Rightarrow k_m = r_p m_p &= \frac{r_e m_e}{\alpha} \approx 3.5176728259 \cdot 10^{-43} \text{ kg } m. \end{aligned} \quad (111)$$

$$k_m c \approx 1.0545717829 \cdot 10^{-34} \text{ k} \cdot \text{m}^2 \cdot \text{s}^{-1} \approx \hbar. \quad (112)$$

$$E = mc^2 = k_m c/t \quad \wedge \quad k_m c = \hbar \Rightarrow E = mc^2 = \hbar/t \text{ kg } \text{m}^2 \text{ s}^{-2} = \hbar/t \text{ J}. \quad (113)$$

$$\begin{aligned} E = mc^2 &= \hbar/t \quad \wedge \quad \omega = 2\pi/t \quad \wedge \quad h \stackrel{\text{def}}{=} 2\pi\hbar \\ \Rightarrow 2\pi E &= 2\pi mc^2 = \hbar(2\pi/t) = \hbar\omega = (2\pi\hbar)/t = h/t. \end{aligned} \quad (114)$$

N. Values of the unit ratios and Planck units

Values of the direct proportion ratios, c_t , c_m , and c_q :

$$c_t = c = 2.99792458 \cdot 10^8 \text{ m} \cdot \text{s}^{-1}. \quad (115)$$

$$\begin{aligned} G &= c_m c_t^2 \quad \wedge \quad G \approx 6.67430(15) \cdot 10^{-11} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2} \Rightarrow \\ c_m &= G/c_t^2 \approx 7.426162 \cdot 10^{-28} \text{ m} \cdot \text{kg}^{-1}. \end{aligned} \quad (116)$$

$$\begin{aligned} k_e &= c_q^2 c_t^2 / c_m \quad \wedge \quad k_e \approx 8.9875517923(13) \cdot 10^9 \text{ N} \cdot \text{m}^2 \cdot \text{C}^{-2} \Rightarrow \\ c_q &= \sqrt{k_e c_m / c_t^2} \approx 8.6175182 \cdot 10^{-18} \text{ m} \cdot \text{C}^{-1}. \end{aligned} \quad (117)$$

From equation 113, the values of the inverse proportion ratios, k_t , k_m , and k_q :

$$k_m \approx 3.51767294 \cdot 10^{-43} \text{ kg} \cdot \text{m}. \quad (118)$$

$$r_p = \sqrt{r_p^2} = \sqrt{k_m c_m} \approx 1.616255 \cdot 10^{-35} \text{ m}. \quad \Rightarrow \quad (119)$$

$$k_t = r_p^2 / c_t \approx 8.71363 \cdot 10^{-79} \text{ s} \cdot \text{m}. \quad (120)$$

$$k_q = r_p^2 / c_q \approx 3.031361 \cdot 10^{-53} \text{ C} \cdot \text{m}. \quad (121)$$

Values of the Planck units, r_p , t_p , m_p , and q_p :

$$r_p = \sqrt{c_t k_t} = \sqrt{c_m k_m} = \sqrt{c_q k_q} \approx 1.616255 \cdot 10^{-35} \text{ m}. \quad (122)$$

$$t_p = r_p / c_t \approx 5.391246 \cdot 10^{-44} \text{ s}. \quad (123)$$

$$m_p = r_p / c_m \approx 2.176434 \cdot 10^{-8} \text{ kg}. \quad (124)$$

$$q_p = r_p / c_q = q_e / \sqrt{\alpha} \approx 1.875546038 \cdot 10^{-18} \text{ C}. \quad (125)$$

O. Compton wavelength, λ

^{14,19} From equations 51 and ??:

$$rm = k_m \quad \wedge \quad k_m c = \hbar \quad \Rightarrow \quad \lambda = 2\pi r = \frac{2\pi k_m}{m} = \frac{2\pi k_m}{m} \frac{c}{c} = \frac{2\pi \hbar}{mc} = \frac{h}{mc}. \quad (126)$$

P. Schrödinger's wave equation

Start with the previous unit ratio-derived “base” Planck-Einstein relation ??, where $V(r, t)$ is the potential energy in a volume. And multiply the kinetic energy component by mc/mc :

$$mc^2 = \hbar/t \quad \Rightarrow \quad \exists V(r, t) : \hbar/t = \hbar/2t + V(r, t). \quad (127)$$

$$\hbar/t = h/2t + V(r, t) \quad \Rightarrow \quad \hbar/t = (\hbar/2t)(mc/mc) + V(r, t). \quad (128)$$

And from the distance-to-time (speed of light) ratio (49):

$$\hbar/t = \hbar mc/2mct + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar mc^2/2mcr + V(r, t). \quad (129)$$

$$\hbar/t = \hbar mc^2/2mcr + V(r, t) \quad \wedge \quad \hbar/t = mc^2 \quad \Rightarrow \quad \hbar/t = \hbar^2/2mcrt + V(r, t). \quad (130)$$

$$\hbar/t = \hbar^2/2mcrt + V(r, t) \quad \wedge \quad r = ct \quad \Rightarrow \quad \hbar/t = \hbar^2/2mr^2 + V(r, t). \quad (131)$$

Multiply both sides of equation 131 by a probability density function, $\Psi(r, t)$.

$$\hbar/t = \hbar^2/2mr^2 + V(r, t) \quad \Rightarrow \quad (\hbar/t)\Psi(r, t) = (\hbar^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t). \quad (132)$$

$$(\hbar/t)\Psi(r, t) = ((\hbar)^2/2mr^2)\Psi(r, t) + V(r, t)\Psi(r, t) \quad \wedge$$

$$\forall \Psi(r, t) : \partial^2 \Psi(r, t)/\partial r^2 = (-1/r^2)\Psi(r, t) \quad \wedge \quad \partial \Psi(r, t)/\partial t = (i 1/t)\Psi(r, t)$$

$$\Rightarrow \quad i\hbar \partial \Psi(r, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t), \quad (133)$$

which is the one-dimensional position-space Schrödinger's equation.^{4,19}

$$i\hbar \partial \Psi(r, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(r, t)/\partial r^2 + V(r, t)\Psi(r, t) \quad \wedge \quad |\mathbf{r}| = r$$

$$\Rightarrow \quad \exists \mathbf{r} : i\hbar \partial \Psi(\mathbf{r}, t)/\partial t = -(\hbar^2/2m)\partial^2 \Psi(\mathbf{r}, t)/\partial \mathbf{r}^2 + V(\mathbf{r}, t)\Psi(\mathbf{r}, t), \quad (134)$$

which is the 3-dimensional position-space Schrödinger's equation.^{4,19}

Q. Dirac's wave equation

Start with the previous unit ratio-derived “base” Planck-Einstein relation 113:

$$\hbar/t = mc^2 \Leftrightarrow \hbar/2t + \hbar/2t = mc^2 \Leftrightarrow \hbar/2t = -\hbar/2t + mc^2. \quad (135)$$

$$\hbar/2t = -\hbar/2t + mc^2 \wedge qc_q/r = qc_q/ct = 1 \Rightarrow \hbar qc_q/2ct^2 = -qc_q/2rt + mc^2. \quad (136)$$

$$\hbar qc_q/2ct^2 = -qc_q/2rt + mc^2 \wedge r = ct \Rightarrow \hbar qc_q/2ct^2 = -qcc_q/2r^2 + mc^2. \quad (137)$$

$$\hbar qc_q/2ct^2 = -qcc_q/2r^2 + mc^2 \Rightarrow \hbar(qc_q/2ct^2)\psi = (-qcc_q/2r^2)\psi + mc^2\psi. \quad (138)$$

$$\psi = e^{-iqc_q((1/2ct)-(1/2r))} \Rightarrow \partial\psi/\partial t = (-iqc_q/2ct^2)\psi \wedge \nabla\psi = (iqc_q/2r^2)\psi. \quad (139)$$

$$\partial\psi/\partial t = (-iqc_q/2ct^2)\psi \wedge \nabla\psi = (iqc_q/2r^2)\psi \wedge$$

$$\hbar(qc_q/2ct^2)\psi = (-qcc_q/2r^2)\psi + mc^2\psi \Rightarrow i\hbar\partial\psi/\partial t = -i\hbar c\nabla\psi + mc^2\psi. \quad (140)$$

$$i\hbar\partial\psi/\partial t = -i\hbar c\nabla\psi + mc^2\psi$$

$$\Rightarrow \exists \alpha, \beta : i\hbar\partial\psi/\partial t = i\hbar c\alpha \cdot -\nabla\psi + mc^2\beta\psi, \quad (141)$$

which is the relativistic, free particle Dirac equation, where $\beta = \gamma^0$, $\alpha = \gamma^0\gamma^k$, $k \in \{1, 2, 3\}$, and each γ a matrix.¹⁷

R. Total of a type

Applying both the direct (49) and inverse proportion ratios (51) to the Euclidean distance:

$$\begin{aligned} r &= \sqrt{r_1^2 + r_2^2}, \quad r_1 = c_\tau\tau, \quad r_2 = k_\tau/\tau, \quad \tau \in \{t, m, q\} \\ &\Rightarrow r = \sqrt{(c_\tau\tau)^2 + (k_\tau/\tau)^2} \Leftrightarrow \tau = \sqrt{(r/c_\tau)^2 + (k_\tau/r)^2}. \end{aligned} \quad (142)$$

S. Quantum extension to Schwarzschild's metric

The simplest way to demonstrate how to add quantum physics to general relativity is by extending Schwarzschild's gravitational time dilation equation and black hole metric. Apply the total of a type equation 142 to the derivation of Schwarzschild's time dilation and metric

(64):

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - (v^2/c^2)(r/r)} \quad \wedge \quad r = \sqrt{(c_m m)^2 + (k_m/m)^2} = Q_m \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2 / r c^2}.\end{aligned}\tag{143}$$

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - Q_m v^2 / r c^2} \quad \wedge \quad KE = mv^2/2 = mv_{escape}^2 \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2 / r c^2}.\end{aligned}\tag{144}$$

For a photon, the escape velocity, $v_{escape} = c$.

$$\begin{aligned}\sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m v_{escape}^2 / r c^2} \quad \wedge \quad v_{escape} = c \\ \Rightarrow \quad \sqrt{1 - (v^2/c^2)} &= \sqrt{1 - 2Q_m c^2 / r c^2} = \sqrt{1 - 2Q_m / r}.\end{aligned}\tag{145}$$

Combining equation 145 with equation 59 yields Schwarzschild's gravitational time dilation with a quantum mass effect:

$$\sqrt{1 - (v^2/c^2)} = \sqrt{1 - 2Q_m / r} \quad \wedge \quad t' = t \sqrt{1 - (v^2/c^2)} \quad \Rightarrow \quad t' = t \sqrt{1 - 2Q_m / r}.\tag{146}$$

Schwarzschild defined the black hole event horizon radius, $\alpha \stackrel{\text{def}}{=} 2Gm/c^2$. The radius with the quantum extension is $\alpha \stackrel{\text{def}}{=} 2Q_m$. At this point the exact same equations 69 through 72 yield what looks like the same Schwarzschild black hole metric.

T. Quantum extension to Newton's gravity force

The quantum mass effect is easier to understand in the context Newton's gravity equation than in general relativity, because the metric equations and solutions in the EFEs are much more complex. From equations 142 and 49:

$$\begin{aligned}m / \sqrt{(r/c_m)^2 + (k_m/r)^2} &= 1 \quad \wedge \quad r^2 / (ct)^2 = 1 \quad \Rightarrow \quad r^2 / (ct)^2 = m / \sqrt{(r/c_m)^2 + (k_m/r)^2} \\ \Rightarrow \quad r^2 / t^2 &= c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2}.\end{aligned}\tag{147}$$

$$\begin{aligned}r^2 / t^2 = c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2} &\Rightarrow (m/r)(r^2 / t^2) = (m/r)(c^2 m / \sqrt{(r/c_m)^2 + (k_m/r)^2}) \\ \Rightarrow \quad F \stackrel{\text{def}}{=} mr / t^2 &= c^2 m^2 / \sqrt{(r^4/c_m^2) + k_m^2}.\end{aligned}\tag{148}$$

$$\begin{aligned}F = c^2 m^2 / \sqrt{(r^4/c_m^2) + k_m^2} \quad \wedge \quad \forall m \in \mathbb{R}, \exists m_1, m_2 \in \mathbb{R} : m_1 m_2 = m^2 \\ \Rightarrow \quad F = c^2 m_1 m_2 / \sqrt{(r^4/c_m^2) + k_m^2}.\end{aligned}\tag{149}$$

U. Quantum extension to Coulomb's charge force

$$q^2/((r/c_q)^2 + (k_q/r)^2) = 1 \quad \wedge \quad r^2/(ct)^2 = 1 \quad \Rightarrow \quad r^2/(ct)^2 = q^2/((r/c_q)^2 + (k_q/r)^2)$$

$$\Rightarrow \quad r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2). \quad (150)$$

$$r^2/t^2 = c^2 q^2/((r/c_q)^2 + (k_q/r)^2) \quad \wedge \quad r = c_m m$$

$$\Rightarrow \quad F \stackrel{\text{def}}{=} mr/t^2 = (c^2/c_m) q^2/((r/c_q)^2 + (k_q/r)^2). \quad (151)$$

$$\forall q \in \mathbb{R} : \exists q_1, q_2 \in \mathbb{R} : q_1 q_2 = q^2 \quad \wedge \quad F = (c^2/c_m) q^2/((r/c_q)^2 + (k_q/r)^2)$$

$$\Rightarrow \quad F = (c^2/c_m) q^2/((r/c_q)^2 + (k_q/r)^2). \quad (152)$$

VI. INSIGHTS AND IMPLICATIONS

1. A distance measure, d , is an inverse (bijective) function of a volume, v , where v is the sum of subset volumes (Euclidean or non-Euclidean), $g(s_{i,1}, \dots, s_{i,n}) \geq 0$. Notionally:

$$\forall n \in \mathbb{N}, \quad f, g \geq 0, \quad \exists d : \quad v = f(d), \quad d = f^{-1}(v) = f^{-1}(\sum_{i=1}^m g(s_{i,1}, \dots, s_{i,n})) : \quad (153)$$

- (a) shows the intimate relation between distance and volume that the, axioms defining inner product space and metric space lack,^{5,6}
- (b) is a more concise and useful (applied math) definition of a distance measure.

2. The left side of the distance sum inequality (III.5),

$$(\sum_{i=1}^m (a_i^n + b_i^n))^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}, \quad (154)$$

differs from the left side of Minkowski's sum inequality.¹⁰

$$(\sum_{i=1}^m (a_i^n + b_i^n)^{\mathbf{n}})^{1/n} \leq (\sum_{i=1}^m a_i^n)^{1/n} + (\sum_{i=1}^m b_i^n)^{1/n}. \quad (155)$$

- (a) The two inequalities are only the same where $n = 1$.
- (b) The distance sum inequality (III.5) is a more fundamental inequality because the proof does not require the convexity and Hölder's inequality assumptions required to prove the Minkowski sum inequality.¹⁰

(c) $\forall n > 1, v > 0$: The left side of the distance sum inequality, $v_i^n = a_i^n + b_i^n$: $d = v^{1/n} = (\sum_{i=1}^m v_i^n)^{1/n}$, is the Minkowski distance, which makes the distance sum inequality directly related to geometry. But the left side of the Minkowski sum inequality, $d = v^{1/n} = (\sum_{i=1}^m ((v_i^n)^{\mathbf{n}}))^{1/n} = (\sum_{i=1}^m v_i^{\mathbf{n}^2})^{1/n}$, is *not* a Minkowski distance.

(d) The distance sum inequality might be useful for comparing distances between concepts in machine learning.

3. *Combinatorics*. The number of ordered combinations (n-tuples) of the elements of disjoint sets was proven to imply the Euclidean volume equation (II.1). And the notion of distance was derived as the inverse function of a volume. The volume and distance equations were derived without relying on the geometric primitives and relations in Euclidean geometry^{11,12}, axiomatic geometry²⁰⁻²⁴, trigonometry^{25,26}, calculus^{25,27,28}, and vector analysis.³

4. *Combinatorics*. The commutative property of the operations defining volume allows n number of domain values to be sequenced in $n!$ number of permutations. But linear sequencing combined with the commutative property limits a set of domain elements to $n \leq 3$ elements (IV.5). Other elements must have different types (are elements of different sets).

(a) If there are more than 3 distance and 3 non-distance dimensions $\subseteq \mathbb{R}$, then the additional dimensions are either not commutative or not linear sequenceable.

(b) There cannot be an ISCL of more than 3 quantum states of the same type.

5. A discrete (point) valued state has measure 0 (zero-length interval). The ratio of a time, distance, mass, or charge interval length to zero is undefined, which is the reason discrete, state values exist independent of distance (position), time, mass, and charge. Therefore, discrete state values are undetermined with respect to space and time until observed.

6. For each constant unit interval length, r_p , of a 3-dimensional distance, there are constant unit interval lengths of 3 other types of dimensions, forming 3 direct constant

proportion unit ratios (49): $r = (r_p/t_p)t = (r_p/m_p)m = (r_p/q_p)q$ and 3 inverse constant proportion ratios (51): $r = (r_p t_p)/t = (r_p m_p)/m = (r_p q_p)/q$, where r_p , t_p , m_p , and q_p are the Planck units (122).

7. In this article, the following constants are derived directly from the ratios and Planck units: gravity, $G = c_m c_t^2 = (r_p/m_p)(r_p/t_p)^2 = r_p^3/m_p t_p^2$ (54), reduced Planck $\hbar = k_m c_t = (r_p m_p)(r_p/t_p) = r_p^2 m_p/t_p$ (113), charge, $k_e = c_q^2 c_t^2/c_m$ (76), vacuum permittivity $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2/c_m)$ (80), and vacuum permeability $\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m = 4\pi r_p m_p/q_p^2 = 4\pi k_m/q_p^2$ (84).

And the quantum extensions to: Schwarzschild's gravitational time dilation (145), Newton's gravity force (149), and Coulomb's charge force (152), show that the constants G , $\kappa = 8\pi G/c^4$, k_e , and ε_0 do not exist, decompose, where the quantum effects are measurable.

This provides the insight that the constants: G , κ , k_e , ε_0 , and μ_0 are *not* fundamental constants because 1) they were all shown to be composed of the Planck units and 2) G , κ , k_e , and ε_0 do not exist, decompose, where the quantum effects are measurable.

8. c_t , c_m , and c_q are the *maximum* proportionality constants: $r = \sqrt{r_1^2 + r_2^2 + r_3^2} \Rightarrow r_1, r_2, r_3 \leq r \Rightarrow \forall \tau \in \{t, m, q\} : r_1/\tau, r_2/\tau, r_3/\tau \leq r/\tau = r_p/\tau_p = c_\tau$. For example, the speeds of gravity and light waves are limited to the proportionality constant, c_t .
9. c_t is a component of the constants: $G = c_m c_t^2$, $k_e = c_q^2 c_t^2/c_m$, $\varepsilon_0 = 1/4\pi k_e = 1/4\pi(c_q^2 c_t^2/c_m)$, and $\hbar = k_m c_t$. The constants, derived in this article, that do not contain c_t are: vacuum permeability: $\mu_0 = 4\pi k_e/c_t^2 = 4\pi c_q^2/c_m$, the fine structures, $\alpha = q_e^2/q_p^2$, and $\alpha_G = m_e^2/m_p^2$, Rydberg, $R_\infty = m_e q_e^4/4\pi r_p m_p q_p^4$.
10. An empirical law and related constant *describes* the relations between the variables in an equation. Deriving an equation and related constant from the ratios *explains* the reasons for relations between the variables in an equation and explains the constant's units (dimensions).

The constant interval lengths, r_p , t_p , m_p , and q_p , have units (dimensions). Therefore, the ratios and all equations and constants derived from the ratios have dimensions, which eliminates the need for dimensional analysis. For example, $G = c_m c_t^2 = r_p^3/m_p t_p^2$, which conforms to the SI units: $m^3 \cdot kg^{-1} \cdot s^{-2}$.¹⁸

11. The ratio-based derivation of Lorentz's law here (86) differs from other relativistic derivations, in that equations for the magnetic field, $\mathbf{B}$, and magnetic permeability, μ_0 , are derived in the process. The ratio-derived equation, $\mathbf{B} = (2\pi k_e/c^2)vq/|\mathbf{r}|^2$, allows a simple derivation of the Faraday's law (91) and the magnetic component of Maxwell's electromagnetic equations (101).
12. Other derivations of Maxwell's electric and magnetic field wave equations require: Gauss's electric field law in a vacuum: $\nabla \cdot \mathbf{E} = -\rho/\varepsilon_0$, Gauss's magnetic field law in a vacuum: $\nabla \cdot \mathbf{B} = 0$, Faraday's Law: $\nabla \times \mathbf{E} = -\partial\mathbf{B}/\partial t$, and Ampere's Law: $\nabla \times \mathbf{B} = \mu_0\varepsilon_0\partial\mathbf{E}/\partial t$.¹⁴ In contrast, the derivations (96) (101), in this article, did not require any of those laws. Just the speed of light ratio, $r = ct$ and ratio-derived definitions of $\mathbf{E}$ and $\mathbf{B}$ are used.
13. As shown in the subsection V E, deriving the Schwarzschild's gravitational time dilation and black hole metric from ratios is a simple way, that in *some* cases, can be used to derive an analytic solution that does not require the complexity of manipulating Christoffel symbols, using determinants, etc.^{15,16}
14. Einstein's relativity equations: 1) assume the Lorentz transformations, 2) assume the laws of physics are same at each local Euclidean-like coordinate point, 3) assume the notion of light, and 4) assume that the speed of light is the same at each coordinate point.^{2,3}

The derivations, in this article, were made without those assumptions (does not even require the notion of light). The unit ratios are ratios of Euclidean distance to time, mass, and charge. Therefore, the unit ratios (like the maximum speed ratio, c_t), and all equations (laws) and constants derived from those ratios and Euclidean distance must be the same at each local Euclidean-like coordinate point.

15. The Planck-Einstein relation (113) was rigorously derived from the ratios, where: $E = mc^2 = k_m c_t/t = \hbar/t$. This shows that some common assumptions about the reduced and full Planck constants are incorrect!
 - (a) $\hbar$ was shown to be composed of the ratio (proportionality) constants, k_m and c_t , where: $k_m c_t = \hbar$. Therefore, the common assumption that the Planck constant is an "atomic" and fundamental constant is incorrect!

$$(b) \ E = mc^2 = k_m c_t / t = \hbar / t \Rightarrow 2\pi E = 2\pi mc^2 = \hbar(2\pi/t) = \hbar\omega = (2\pi\hbar)/t = h/t.$$

Therefore, the common assumption that $E = mc^2 = \hbar\omega = h/t$ is incorrect!

Further, the derivations, in this article, of Schrödinger's wave equation (134)⁴ and Dirac's wave equation (141)¹⁷ from the ratio-derived Planck-Einstein relation, $E = mc^2 = \hbar/t$ (113), are shorter, simpler, and require fewer assumptions.

Therefore, it would be more logical for CODATA to define the reduced Planck constant as $\hbar = k_m c_t$ and define full Planck constant in terms of the reduced Planck constant, $h = 2\pi\hbar$.

16. Algebraic manipulation of the classical electron radius, r_e (104), yields the fine structure constants, α (105) and α_G (106), and the Rydberg constant, R_∞ (107).

The derivations of α (105) and α_G (106) indicate the parsimonious explanation that the constants are each the ratio of two subtypes of force: Coulomb's charge force and Newton's gravity force:

$$\frac{F_{q_e}}{F_{q_p}} = \frac{k_e q_e^2 / r^2}{k_e q_p^2 / r^2} = \frac{q_e^2}{q_p^2} = \alpha. \quad (156)$$

$$\frac{F_{m_e}}{F_{m_p}} = \frac{G m_e^2 / r^2}{G m_p^2 / r^2} = \frac{m_e^2}{m_p^2} = \alpha_G. \quad (157)$$

:

17. The complicated and obtuse NIST/CODATA definition of the vacuum permeability constant, $\mu_0 = 2h\alpha/cq_e^2$,¹⁸ (derived from dimensional analysis) reduces to a simpler ratio-derived definition. Using the ratios to reduce the old definition of μ_0 derived in this article:

$$\mu_0 = 4\pi k_e / c_t^2 = 4\pi c_q^2 / c_m = 4\pi (r_p / q_p)^2 / (r_p / m_p) = 4\pi m_p r_p / q_p^2 = 4\pi k_m / q_p^2. \quad (158)$$

Using the ratios to reduce the CODATA definition, $\mu_0 = 2h\alpha/cq_e^2$:¹⁸

$$h = 2\pi k_m c_t, \ \alpha = q_e^q / q_p^2 \Rightarrow \mu_0 = 2h\alpha/cq_e^2 = 4\pi k_m c_t (q_e^2 / q_p^2) / c_t q_e^2 = 4\pi k_m / q_p^2. \quad (159)$$

18. The complicated and obtuse CODATA definition, $\alpha = q_e^2 / 4\pi\epsilon_0 \hbar c$, (derived from di-

dimensional analysis)¹⁸ reduces to the simpler, ratio-derived equation, in this article:

$$\begin{aligned}\varepsilon_0 &= 1/4\pi k_e = 1/(4\pi(c_q^2 c_t^2/c_m)) \quad \wedge \quad \hbar = k_m c_t \quad \wedge \quad h = 2\pi\hbar \\ \Rightarrow \quad \varepsilon_0 h c &= 2\pi k_m c_t^2 / (4\pi(c_q^2/c_m)c_t^2) = k_m / (2(c_q^2/c_m)) \\ &= m_p r_p / (2((r_p/q_p)^2 / (r_p/m_p))) = q_p^2/2. \quad (160)\end{aligned}$$

$$\alpha = q_e^2/2\varepsilon_0 h c \quad \wedge \quad \varepsilon_0 h c = q_p^2/2 \quad \Rightarrow \quad \alpha = q_e^2/2(q_p^2/2) = q_e^2/q_p^2. \quad (161)$$

19. The complicated and obtuse fine structure electron gravity coupling constant (derived from dimensional analysis) reduces to the simpler, ratio-derived equation:

$$\alpha_G = G m_e^2 / \hbar c = (c_m c_t^2) m_e^2 / (k_m c_t) c_t = (r_p/m_p) m_e^2 / m_p r_p = m_e^2 / m_p^2. \quad (162)$$

20. The obtuse NIST/CODATA definition of the Rydberg constant (derived from dimensional analysis)¹⁸ reduces to the more informative ratio-derived equation (107):

$$\begin{aligned}R_\infty &= \frac{m_e q_e^4}{8\varepsilon_0^2 h^3 c} = \frac{m_e q_e^4}{8(1/4\pi k_e)^2 (2\pi\hbar)^3 c_t} = \frac{m_e q_e^4 (4\pi k_e)^2}{8(2\pi\hbar)^3 c_t} = \frac{m_e q_e^4 (c_q^2 c_t^2/c_m)^2}{4\pi (k_m c_t)^3 c_t} \\ &= \frac{m_e q_e^4 (c_q^2/c_m)^2}{4\pi k_m^3} = \frac{m_e q_e^4 c_q^4}{4\pi k_m^3 c_m^2} = \frac{m_e q_e^4 (r_p/q_p)^4}{4\pi (r_p m_p)^3 (r_p/m_p)^2} = \frac{m_e q_e^4}{4\pi r_p m_p q_p^4}. \quad (163)\end{aligned}$$

21. The quantum extensions to: Schwarzschild's black hole metric (72), Newton's gravity force (149), and Coulomb's charge force (152) make quantifiable predictions:

- (a) For gravity, $\lim_{r \rightarrow 0} F = c^2 m_1 m_2 / k_m$, and for charge, $\lim_{r \rightarrow 0} F = 0$. Finite maximum gravity and charge forces: (1) eliminates the problem of forces going to infinity as $r \rightarrow 0$, (2) finite peak forces might account for radioactivity, and (3) finite sloped energy well walls might facilitate quantum tunneling.
- (b) The quantum-extended relativity and classic equations reduce to the non-extended relativity and classic equations and constants, where the distance between 2 masses and between 2 charges is sufficiently large or the masses and charges sufficiently large that the quantum effects are not measurable. *Note* that G , k_e , ε_0 , μ_0 , and κ (Einstein's constant, which contains G) do not exist, decompose, where the quantum effects becomes measurable.
- (c) The covariant tensor components, in Einstein's field equations, that had the units $1/\text{distance}^2$, will now have the more complex units, $1/\sqrt{(\text{distance}^4/c_\tau^2) + k_\tau^2}$, $\tau \in \{t, m\}$.

- (d) $1/\sqrt{(distance^4/c_\tau^2) + k_\tau^2}$ implies that as distance $\rightarrow 0$, spacetime curvature peaks at the Planck unit length (distance), r_p (122). The ratio of quantum units, $m_p/r_p^3 \approx 5.15484810^{96} \text{ kg m}^{-3}$, might indicate a maximum mass density. A finite force (spacetime curvature) and finite mass density might imply that black holes have sizes > 0 (are not singularities). It might impact whether black holes evaporate and whether a big bang might not have originated from a singularity.

REFERENCES

- ¹Copyright © 2026 George M. Van Treeck. All rights reserved.
- ²A. Einstein, *Relativity, The Special and General Theory* (Princeton University Press, 2015).
- ³H. Weyl, *Space-Time-Matter* (Dover Publications Inc, 1952).
- ⁴E. Schrödinger, *Collected Papers On Wave Mechanics* (Minkowski Institute Press, 2020).
- ⁵R. R. Goldberg, *Methods of Real Analysis* (John Wiley and Sons, 1976).
- ⁶W. Rudin, *Principles of Mathematical Analysis* (McGraw Hill Education, 1976).
- ⁷W. Conradie and V. Goranko, *Logic and Discrete Mathematics* (Wiley, 2015).
- ⁸X. P. N. Guigui, N. Miolane, “Introduction to riemannian geometry and geometric statistics: from basic theory to implementation with geomstats,” *Foundations and Trends in Machine Learning* **16**, 182 (2023), <https://inria.hal.science/hal-03766900v2>, hal-00644459.
- ⁹Rocq, “Rocq proof assistant,” (2025), <https://rocq-prover.org/docs>.
- ¹⁰H. Minkowski, *Geometrie der Zahlen* (Chelsea, 1953) reprint.
- ¹¹D. E. Joyce, “Euclid’s elements,” (1998), <http://aleph0.clarku.edu/~djoyce/java/elements/elements.html>.
- ¹²E. S. Loomis, *The Pythagorean Proposition* (NCTM, 1968).
- ¹³I. Newton, *The Mathematical Principles of Natural Philosophy* (Patristic Publishing, 2019) reprint.
- ¹⁴R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics* (California Institute of Technology, 2010) <https://www.feynmanlectures.caltech.edu/>.
- ¹⁵K. Schwarzschild, “Über das gravitationsfeld eines massenpunktes nach der einsteinschen theorie,” *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften* **7**, 198–196 (1916), <https://archive.org/stream/sitzungsberichte1916deutsch/page/188/mode/2up>.

- ¹⁶S. Antoci and A. Loinger, “An english translation of k. schwarzschild’s 1916 article: On the gravitational field of a mass point according to einstein’s theory,” ARXIV.org (1999), <https://arxiv.org/abs/physics/9905030>.
- ¹⁷P. Dirac, *The Principles of Quantum Mechanics* (Snowball Publishing, 1957).
- ¹⁸CODATA, “Physical constant values,” (2022), <https://physics.nist.gov/cuu/Constants/index.html>.
- ¹⁹M. Jain, *Quantum Mechanics: A Textbook for Undergraduates* (PHI Learning Private Limited, New Delhi, India, 2011).
- ²⁰G. D. Birkhoff, “A set of postulates for plane geometry (based on scale and protractors),” *Annals of Mathematics* **33** (1932).
- ²¹D. Hilbert, *The Foundations of Geometry (2cd Ed)* (Chicago: Open Court, 1980) <http://www.gutenberg.org/ebooks/17384>.
- ²²J. M. Lee, *Axiomatic Geometry* (American Mathematical Society, 2010).
- ²³O. Veblen, “A system of axioms for geometry,” *Trans. Amer. Math. Soc* **5**, 343–384 (1904), <https://doi.org/10.1090/S0002-9947-1904-1500678-X>.
- ²⁴A. Tarski and S. Givant, “Tarski’s system of geometry,” *The Bulletin of Symbolic Logic* **5**, 175–214 (1999).
- ²⁵A. Bogomolny, *Pythagorean Theorem* (Interactive Mathematics Miscellany and Puzzles, 2010) <http://www.cut-the-knot.org/pythagoras/CalculusProofCorrectedVersion.shtml>.
- ²⁶J. Zimba, “On the possibility of trigonometric proofs of the pythagorean theorem,” *Forum Geometricorum* **9**, 275–278 (2009).
- ²⁷B. C. Berndt, “Ramanujan-100 years old (fashioned) or 100 years new (fangled)?” *The Mathematical Intelligencer* **10** (1988).
- ²⁸M. Staring, “The pythagorean proposition a proof by means of calculus,” *Mathematics Magazine* **69**, 45–46 (1996).