

topology

Runi Malladi

June 17, 2023

1 metric spaces

Definition 1.1. Let (X, d) be a metric space. A *curve* in X is a continuous mapping $\gamma : [0, 1] \rightarrow X$.

Definition 1.2. Let γ be a curve in (X, d) . Let $P[0, 1]$ denote the set of partitions of $[0, 1]$. For each $\mathcal{P} \in P[0, 1]$, with components $p_0 < \dots < p_n$, define

$$\ell_{\mathcal{P}}(\gamma) = \sum_{i=1}^n d(\gamma(p_{i-1}), \gamma(p_i)).$$

The *length* of γ is

$$\ell(\gamma) = \sup_{\mathcal{P}} \ell_{\mathcal{P}}(\gamma).$$

Proposition 1.3.

$$\lim_{|\mathcal{P}| \rightarrow 0} \ell_{\mathcal{P}}(\gamma) = \ell(\gamma).$$

Proof. Fix $\epsilon > 0$. We need to find $\delta > 0$ such that $\ell(\gamma) - \epsilon < \ell_{\mathcal{Q}}(\gamma) \leq \ell(\gamma)$. By the definition of the supremum, there exists $\mathcal{P} \in P[0, 1]$ such that

$$\ell(\gamma) - \epsilon < \ell_{\mathcal{P}}(\gamma) \leq \ell(\gamma).$$

Let $\epsilon' > 0$ be arbitrary. Since σ is continuous on a compact space, it is uniformly continuous (REF), hence there exists $\eta > 0$ such that $|t - t'| < \eta$ implies $d(\gamma(t), \gamma(t')) < \epsilon'$ for all $t, t' \in [0, 1]$. Let $\delta = \min(\eta, |\mathcal{P}|)$.

Suppose $\mathcal{Q} \in P[0, 1]$ is such that $|\mathcal{Q}| < \delta$. Say it is given by $q_0 < \dots < q_m$. Let $\mathcal{R} = \mathcal{P} \cup \mathcal{Q}$. For each $i = 1, \dots, n$, there exists a finite set $\{r_{i,k}\}_{k=0}^{K_i} \subset \mathcal{R}$ partitioning $[p_{i-1}, p_i]$. Then

$$\begin{aligned} \ell_{\mathcal{R}}(\gamma) &= \sum_{i=1}^n \sum_{k=1}^{K_i} d(\gamma(r_{i,k-1}), \gamma(r_{i,k})) \\ &\leq \sum_{i=1}^n d(\gamma(\max_{q_j \leq r_{i,0}} q_j), \gamma(r_{i,0})) + \sum_{k=1}^{K_i} d(\gamma(r_{i,k-1}), \gamma(r_{i,k})) + d(\gamma(r_{i,K_i}), \gamma(\min_{q_j \geq r_{i,K_i}} q_j)) \\ &\leq \sum_{i=1}^n d(\gamma(\max_{q_j \leq r_{i,0}} q_j), \gamma(\min_{q_j \geq r_{i,K_i}} q_j)) \end{aligned}$$




Figure 1: This is an example of what $\mathcal{P} \cup \mathcal{Q}$ could look like. The green line extends from q_{j-3} to q_j and represents the domain of one summand, and the blue line extends from q_{j-1} to q_{j+1} and represents the next summand. The yellow line from q_{j-1} to q_j represents the overlap; this region is “double counted”.

$$\leq \ell_{\mathcal{Q}}(\gamma) + 2n\epsilon'.$$

Figure 1 demonstrates our strategy here.

Now

$$\begin{aligned} \ell(\gamma) - \epsilon &< \ell_{\mathcal{P}}(\gamma) \leq \ell_{\mathcal{R}}(\gamma) \leq \ell_{\mathcal{Q}}(\gamma) + 2n\epsilon', \\ \ell_{\mathcal{R}}(\gamma) - 2n\epsilon' &\leq \ell_{\mathcal{Q}}(\gamma). \end{aligned}$$

But our choice of ϵ' was arbitrary; if we pick, say,

$$\epsilon' = \frac{1}{2n} \cdot \frac{\ell_{\mathcal{P}}(\gamma) - (\ell(\gamma) - \epsilon)}{4}$$

then we can choose δ in such a way that that, whenever $|\mathcal{Q}| < \delta$, we have

$$\ell(\gamma) - \epsilon < \ell_{\mathcal{P}}(\gamma) - 2n\epsilon' \leq \ell_{\mathcal{Q}}(\gamma) < \ell(\gamma)$$

as desired. □

Proposition 1.4. Let $s(t) = \ell(\gamma|_{[0,t]})$ for $0 \leq t \leq 1$. Then s is continuous.

Proof. Let $\{t_n\}_1^\infty$ be a convergent sequence in $[0, 1]$ with limit t . Then

$$\begin{aligned} \lim_{n \rightarrow \infty} s(t_n) &= \lim_{n \rightarrow \infty} \ell(\gamma|_{[0,t]}) \\ &= \lim_{n \rightarrow \infty} \lim_{|\mathcal{P}_n| \rightarrow 0, \mathcal{P}_n \in P[0, t_n]} \ell_{\mathcal{P}_n}(\gamma) \\ &= \lim_{|\mathcal{P}| \rightarrow 0, \mathcal{P} \in P[0, t]} \ell_{\mathcal{P}}(t) = \ell(\sigma|_{[0, t]}) \\ &= s(t). \end{aligned}$$

□